them, however, the deformity has continued to progress more especially in the direction of elevation of the trochanters, which are now about half an inch above Nélaton's line, with the result that the left leg is in each case shorter than the one operated on.

3. *The Deformity on the Side Operated on has come to a Stand-still in both Cases.*—This is to me quite an unexpected result. Why it should have become arrested on the side operated on while it has progressed on the other is somewhat difficult of explanation. The trochanters on the limbs operated on still remain at the level of Nélaton's line; while, as has been said, on the other side they are half an inch at least higher. Consequently, the legs operated on are longer than the others. Two suggestions appear to me possible:

1. The process of repair and consolidation of the bone around the point of fracture may have extended upwards into the neck of the bone, and led to consolidation and arrest of the softening process, which was leading to the curvature.

2. The branches of the nutrient artery to the neck of the bone were divided in the operation, and, as a result of diminished vascularity, consolidation of the neck may have been hastened.

A CASE OF A TOOTH IMPACTED IN THE LEFT BRONCHUS: GANGRENE OF THE LEFT LUNG: DEATH.

By JAMES S. WARRACK, M.A., M.D.,
Demonstrator of Physiology in the University of Aberdeen.

A WOMAN, aged 26, was admitted to the Victoria Hospital, Burnley, on December 18th, and died on December 30th, 1897. She stated, on admission, that four days previously she was given gas for the purpose of having a tooth extracted. At the critical moment she appears to have taken a deep breath, the result being that the tooth, which must have escaped from the forceps, was inhaled into the air passages. Thereafter extreme cyanosis occurred, and she had a feeling of tightness in the throat, with dyspnoea, and a hard dry cough, aggravated by speaking or change of position. She had also a vague feeling of something being fixed inside the chest, but could not indicate the position. When I saw her she was lying on her back, with livid lips, and coughing in frequent paroxysms, but without expectoration. The coughing was much aggravated by attempts to speak, and by change of posture.

The left side of the chest scarcely moved at all on respiration. There were no breath sounds audible over the left side of the chest, with the exception of tubular breathing at the left apex, accompanied by small moist râles, and this was indistinct posteriorly. On the outside of the left nipple there was a small area over which a friction rub was audible. On the left side, from the level of the nipple down to an inch from the umbilicus and backwards as far as the mid-axillary line, there was a well-marked area of hyperæsthesia, which corresponded roughly to the distribution of the sixth, seventh, and eighth dorsal spinal roots. The breath sounds were harsh on the right side but otherwise normal. The pulse and respiration-rates were much quickened.

During the next few days there was little alteration in the patient's condition, but the temperature assumed the hectic type.

On December 27th, thirteen days after the accident, there were well-marked signs of gangrene of the left lung. The breath was offensive, and there was expectoration of a brown and foul-smelling sputum. The lower cervical glands on the left side were enlarged, and there was a well-marked friction rub in the left axilla. Her condition became gradually worse until death occurred on December 30th.

At the necropsy on December 31st, 1897, I explored the trachea up to the larynx, but no foreign body was found there. All over the left pleural cavity there were numerous recent adhesions, and on separating these some grumous fluid came away from the lung. The heart was flabby and rather smaller than usual, and its right side was dilated. On cutting open the left bronchus the missing tooth was found. It was very tightly wedged in the bronchus, its point being downwards and the crown upwards, so that it had

apparently acted as a "ball valve." It had almost ulcerated through the left bronchus, which above the situation of the tooth was congested and full of grumous fluid. On following the left bronchus into the lung there was found an irregular passage full of a brown foul-smelling fluid. The whole of the left lung was gangrenous and crepitating, and on squeezing the lung substance there exuded a fluid of the character described. The bronchial glands were enlarged, and there was also an enlarged gland at the root of the neck on the left side. Above the impacted tooth no bronchiole was given off, so that there must have been entire obstruction. The right lung was congested and oedematous, and showed commencing consolidation at the base. There were two cavities in the left lower jaw, and into the hinder one the tooth, which was carious, fitted exactly, its direction being forwards and inwards.

This case seems to illustrate the effects of obstruction to a bronchus, as Dr. Sevestre has recently pointed out in the *BRITISH MEDICAL JOURNAL*, the early stage being collapse of the lung, and the later, inflammation, leading sometimes to the entire disorganisation and gangrene of the lung. The tooth did not appear to have been tightly wedged in at first when the patient was seen, but seemed to have become so owing to the incessant attempts to expel the foreign body, and with each attempt the conditions must have been aggravated by the suction in during inspiration. The gangrene would be excited both by the retention of secretion, and by the introduction of septic material from the tooth.

CASE OF ACUTE TETANUS SUCCESSFULLY TREATED BY ANTITETANIN.

By J. GALLETLY, M.B., C.M.,
Bourne, Lincs.

G. F., a boy, aged 8, was seen by me for the first time at 10 P.M. on September 19th, 1898. He lay on his back in bed with his lower extremities rigid and fully extended, and his head, neck, and shoulders propped up with pillows. The abdomen had a flat, board-like appearance, due to contraction of all the muscles of its anterior wall. The head was somewhat retracted; he could open his mouth widely though slowly, and protrude his tongue. He had little difficulty in swallowing liquids, and could move his arms with considerable freedom. The temperature was normal and the pulse regular, with a rate of 90 beats per minute.

On inquiry, I found that the terminal phalanx of his right middle finger had been crushed in the cogwheel of a winnowing machine on August 28th. The wound had been somewhat roughly dressed, and examination showed that it was not soundly healed, the scar looking red and irritable. From the date of the injury up to September 17th—a period of nineteen days—the boy was apparently in his usual health. On that day it was noticed that he was a little stiff and awkward in his movements. On the 18th, the stiffness became more marked, and he was kept in bed.

On the 19th I was sent for. As the bowels had not acted for three days, a 2-gr. dose of calomel was given, and 10 gr. of bromide of potassium every three hours were ordered. He had a very restless night, and on the following morning more of the muscles were affected, the risus sardonicus was well marked, and the convulsive seizures were occurring with greater frequency and increased severity. Tetanus antitoxin was telegraphed for from the British Institute of Preventive Medicine, and in the evening of the same day (the 20th), chloroform having been administered, 20 c.cm. of the antitoxin were injected under the skin of the abdomen, and the terminal phalanx of the injured finger was amputated. Just before giving the chloroform he was seized every few minutes with attacks of marked opisthotonos. At 9 A.M. on the 21st, 10 c.cm. of antitoxin were injected, and again at 9 P.M. this dose was repeated. So far there was no improvement, the attacks recurring with great severity every few minutes. The tongue was badly bitten several times. He lay with his abdomen and lower extremities wholly uncovered, as the weight of the bedclothes—even of his night-shirt—was painful, and seemed to start a seizure. The next morning the attacks were less frequent, and he had snatches of sleep. A further dose of 10 c.cm. of antitoxin was injected, and this amount

was repeated twice a day till September 25th, when, as his condition seemed stationary, it was increased to 15 c.cm. On the 26th I injected 10 c.cm. twice, and this dose was continued twice a day till the 29th, when, as the improvement was becoming unmistakable, 10 c.cm. was given only once a day till October 5th, when it was omitted. Altogether 240 c.cm. of tetanus antitoxin were injected. The boy was sufficiently recovered by October 15th to be downstairs, and when I saw him for the last time on October 31st there was no perceptible stiffness in any of his muscles.

An urticarial eruption appeared on the fourth day of the injections in the neighbourhood of the needle punctures, and spread over most of the body, disappearing about the twelfth day.

The temperature was never much above normal; the pulse was usually somewhat quick; the bowels were obstinately confined, and the fæces very offensive. He was always able to take a fair amount of milk and beef-tea from a spoon. The amputation wound healed by first intention.

Acute traumatic tetanus, of which I take the above to be an example, is so commonly followed by a fatal result that I think this case worthy of record in helping to form an opinion as to the usefulness of antitetanin; for, with the exception of a few doses of bromide of potassium and chloral hydrate during the first two days, and an occasional aperient, no drug was administered.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

A CASE OF GANGRENE OF THE LUNG.

A SHORT note of a case, very similar to the first of the two cases described by Dr. Byrom Bramwell in the BRITISH MEDICAL JOURNAL of January 14th, 1899, will be, perhaps, of some interest.

On May 22nd, 1898, I was called to see a man aged 62. He had all the signs and symptoms of peritonitis. I could discover no cause for his illness. He said the attack came on quite suddenly, and that previously he had been quite well. His bowels had been quite regular; he was not vomiting. He said that he had suffered from chronic dyspepsia for years, but at no time had he had anything acutely wrong with his stomach or bowels. In about a week's time he was quite well, and during that time he had no sickness or any other stomach symptom. I treated him with hypodermic injections of morphine and atropine, hot fomentations, milk, and soda water. During this attack his lungs were absolutely healthy.

On June 2nd he reported himself as being quite well. I examined him on this day, and could detect nothing wrong with his lungs or abdomen except that his stomach was markedly dilated.

He continued quite well until July 11th, when I was called to see him in the evening. He told me that on the morning of July 11th, when jumping on to a tramway car which was in motion he felt a sudden pain a little above the crest of the left ilium, and that immediately after this he began to cough up a large quantity of a very badly-smelling fluid. I found some coarse crepitations at the base of the left lung posteriorly and vesicular breathing, but no dulness. The temperature was 102°, and the pulse 86. He had no abdominal symptoms. He was coughing frequently, and his sputum was very copious and intensely fetid. He said he had coughed up at least three large breakfastfuls of this material since the morning, and that he had no cough till after he felt sudden pain above described. During the next twenty-four hours he coughed up about twice the amount of the same kind of material. Dulness could now be made out; it extended from the base to almost the spine of the scapula. Breath sounds were now bronchial in type, and the crepitations were far more numerous at the base, and some could be detected scattered all over the dull area. The right lung was normal. In a few days the sputum diminished in quantity, and the lung began to clear up a little. Everything was going on well, when one day when turning in bed he had again a severe pain at the same spot as before. He now coughed up a great quantity of the same

fetid sputum, but this time it was tinged with blood. Crepitations were now found to be more numerous at the base than they had been before. A few days after this his motions were well mixed with blood, and about that time he passed by the bowel about an ounce of fluid which had exactly the same smell and appearance as his sputum. His sputum now rapidly diminished in quantity, and the lung cleared up. He is now (January, 1899) apparently quite well, the lung appears absolutely normal, and he has had no recurrence of the abdominal symptoms.

His sputum was examined three days after his lung symptoms began, and had the following characters:

It was a thin, greyish, purulent fluid, extremely fetid. Microscopically it presented: (1) Pus cells mostly presenting advanced fatty degeneration; (2) epithelial cells, squamous, cylindrical, and round, in considerable numbers; (3) a large amount of fat; (4) mucous shreds and blood pigment. No definite trace of lung tissue; no tubercle bacilli. Streptococci in large numbers, also diplococci pneumoniae. Starch corpuscles were not seen.

I think there is little doubt that the lung condition arose from something abdominal. I am inclined to think his abdominal condition and his lung condition arose from the same cause, but I am unable to say what the nature of that cause was. His stomach is markedly dilated, probably from chronic dyspepsia; but he says that he has had nothing at any time acutely wrong with his stomach or bowels. The apex beat is displaced several inches to the left. The urine was normal during his illness. I treated his lung condition by inhalations of eucalyptus oil and creosote.

Glasgow.

H. ARGYLL ROBERTSON, M.B., C.M. Edin.

A CASE OF INVERSION OF THE UTERUS.

INVERSION of the uterus is so rare that the following case seems worth recording. Only 1 case occurred in 190,000 at the Rotunda. Denham had 1 case in 100,000; C. Braun had no case in 250,000. Crosse in 1847 noted that in 28 per cent. of recorded cases of inversion death occurred in a few hours, that in 42 per cent. death followed later, and that in 7 per cent. the accident led to a fatal result more than a year after its occurrence.

Mrs. R. W., aged 37, is a well-developed woman of average size, who has not suffered from any illness since childhood. Her menstrual history was normal. She had had two previous labours but no miscarriage. Her first labour took place six and a-half years ago, and lasted fifteen hours; a female child presented by the vertex, and was safely delivered with forceps. The second labour occurred four and a-half years ago, and occupied only about an hour, a small female child being born safely before the arrival of the medical man. The third labour began at 1 A.M. on February 21st, 1898. The membranes were ruptured immediately. At 2 A.M. the pains ceased and did not return until noon. They continued strong and regular until 4 P.M., after which they diminished in frequency and intensity.

At 6 P.M. I was sent for, and at 6.45 P.M. found the os uteri fully dilated, the presentation vertex, and the position right occipito-posterior. The pelvic outlet was fairly roomy, and the abdomen was very large. Short weak pains occurred at intervals of about fifteen minutes until 8 P.M. without causing progress. At 9.30 P.M. I returned to the patient with Dr. F. L. Mackenzie, prepared to give an anæsthetic if required, and deliver with forceps. On re-examination it was deemed advisable to apply the forceps without an anæsthetic. This was successfully done, and at 9.45 P.M. a living male child, which was very large, weighing 10 lbs. and measuring 22 inches in length, was delivered "occiput to the back."

The patient now received a draught containing 3ss of liquid extract of ergot (B.P.). Ten minutes later the placenta was expressed with some difficulty. The descent of the placenta was estimated by withdrawing from the vagina as much of the cord as came down easily without pulling on it. Copious bleeding followed the birth of the placenta, and the hand being applied to the abdomen, the uterus could not definitely be made out. The hemorrhage continuing, a douche of hot water was prepared. The syringe, however, burst immediately, which was fortunate, as it led to the application of other remedies and the discovery of the true condition of things earlier than would have been the case had time been spent in giving the douche. The bleeding partially ceased, only to recommence a few minutes later. The patient had by this time lost about 5 pints of blood, and the pulse was becoming somewhat feeble, so it was decided to plug the uterus, although there were no symptoms of collapse.

On making a vaginal examination, a large quantity of clot and some fragments of membranes were removed. A soft mass was then felt by the fingers, a bag-like protuberance, round whose neck was a hard round ring. The cause of the post-partum hemorrhage was now seen to be total inversion of the uterus. The diagnosis was confirmed by pressing the inverted uterus upwards and forwards, when the os uteri could be distinctly felt by the external hand through the loose abdominal wall. This portion of the uterus was pressed downwards and backwards, while counter-pressure was made upon the fundus from the vagina. After the return to the normal condition, the remaining portion of the adherent membranes was scraped away with the fingers. The patient complained of feeling faint and the pulse was very feeble, the loss of blood by this time amounting to nearly 6 pints: 3j of spirit. ammon. arom. was given, followed later by 3ss of brandy with water. Bovril, milk, and brandy were given every two hours for the next twelve hours. The patient next morning felt quite comfortable except for the soreness of the womb and abdomen. She made an uninterrupted recovery.