more than mention in this place. I again must say that not all, though a very large number of curvatures, are due to pelvic deviation; it is, therefore, those only for which prevention is here proposed.

(To be continued)

A CONTRIBUTION TO THE TREATMENT OF PUS IN THE PELVIS.*

By J. FURNEAUX JORDAN, F.R.C.S.,

Surgeon to the Birmingham and Midland Counties Hospital for Women.

FOUR or five years ago, under the influence of the teaching of Lawson Tait and C. J. Cullingworth, we might have looked upon the treatment of pus in the pelvis and pelvic organs as settled for ever. The brilliant results achieved by these operators, especially by the former, are only equalled by the clear and decisive way in which they have described the cases, with their pathology and treatment. The treatment laid down by Cullingworth in his inaugural address on Pelvic Abscess delivered to the Midland Medical Society in the autumn of 1893, may be summed up as follows: For all cases of pyosalpinx, tubo-ovarian abscess, and ovarian abscess removal in its entirety of the organ containing the pus through an incision in the abdominal wall. For all cases of extra-peritoneal suppuration—that is, of pelvic cellulitis gone on to suppuration—incision and drainage through the point in the abdominal wall above Poupart's ligament where the abscess usually points. Mr. Tait in his book on abdominal surgery advocates abdominal section and drainage for cases of suppurating hæmatocele of the broad ligament, and even for cases of pus in the broad ligament.

In the Obstetrical Society's *Transactions*, 1892, vol. xxxiv, p. 254, is a most interesting paper by C. J. Cullingworth on the value of abdominal section in certain cases of pelvic peritonitis. Of 50 cases reported here in full there was suppuration in 30, chiefly tubal and tubo-ovarian; of these 30 cases 6 died, giving a mortality of 20 per cent.

In the last ten years of his work at the Women's Hospital, from 1883 to 1892 inclusive, Mr. Tait operated upon 71 cases of pyosalpinx with 11 deaths, a mortality of 15.6 per cent.

In the last ten years, 1888 to 1897 inclusive, in the Women's Hospital the appendages have been removed for pyosalpinx 105 times with a mortality of 15.2 per cent.

My own cases of pyosalpinx and tubo-ovarian abscess are 20 in number with 3 deaths, a mortality of 15 per cent.

All these figures refer to cases of pyosalpinx treated by abdominal section, and we may therefore take it that the mortality in these cases is on an average 15 per cent., or possibly, even a little higher. Up to the spring of 1897 I had operated upon 15 of these cases with 1 death, and I confess that I was quite satisfied that the method of operation as practised was the best, and should be adopted in all cases. During the summer of 1897 I was led to consider the question again. Two deaths which occurred in my practice at that time—1 at the hospital and 1 in private—impressed upon me the necessity, if possible, of some improvement in the method of treating a certain class of these cases. I say "a certain class of these cases" because I want it to be clearly understood that for the great majority of cases of pyosalpinx removal through an abdominal incision is undoubtedly the best method of treatment. I will even add that removal by this route—the treatment, in fact, first laid down by Mr. Tait—is the ideal method in every case. But if by some other method we can give the patient a greater chance of life and an equal relief from her suffering, then I maintain that it is our duty to adopt that method. This method is incision and drainage from the vagina, a method mentioned by Mr. Taylor in his recent paper to the Midland Medical Society.

Before dwelling in any detail upon the method, let me first of all say in what class of cases I should adopt it. I advise its adoption in cases of long duration in which the pyosalpinx, or an abscess communicating with it, can be felt filling up Douglas's pouch, in which the patient is exhausted both by pain and prolonged suppuration, and suffers from emaciation and night sweats. She is, in fact, in a hectic condition, and

absolutely unfit to stand a prolonged abdominal section. The intensity and the amount of the pain in these cases are an indication of the extent of the peritonitis and consequent adhesions and possible localised intrapelvic suppuration. The salpingitis itself does not cause intense pain; it is the peritonitis due to extension of inflammation or even leakage through the abdominal ostium of the tube that causes pain. The peritonitis may be suppurative, but, as a rule, the pus is shut off from the general peritoneal cavity by adhesions, and forms a localised intraperitoneal abscess communicating with the pus sac formed by the tube. It is in these cases that the tube or pus sac forms very dense adhesions with neighbouring organs. Adhesions are encountered in most cases of pyosalpinx; these, as a rule, can be separated easily and safely with the fingers or mop. It is not, however, always so; in the more severe cases they can only be separated with great difficulty, and with the risk every second of tearing the bowel. In my two fatal cases of the summer of 1897 I removed from one a double pyosalpinx, from the other a large tubo-ovarian abscess; in both the adhesions to bowel were extensive and strong. I separated them as gently as I could with a mop, and, as far as I could tell, without any injury to the bowel, but both patients died of perforative peritonitis. The abscess cavities in both of them could have easily been incised from the vagina, the adhesions would not have been disturbed, the cavities would have granulated up, and I believe I should have saved them both.

I have no doubt I shall be told that it is not good or scientific surgery to leave the pus sacs behind; that it is a reversion to the old method of tapping from the vagina; that suppurating sinuses will be left after convalescence, which will discharge for an indefinite time; and that it is quite possible that all the pus sacs will not be opened. I can only say that surgery which cures with the least possible risk to life is to my mind good surgery. That it is a reversion to the old method of tapping I absolutely deny. That was and always will be a step in the dark, and can never result in good drainage, without which the pus sac will never heal up.

As regards suppurating sinuses, I can only say that I have not had any experience of them in vaginal cases. Are they unknown in the cases treated by the abdominal route? My only experience of an obstinate sinus was in a case of large double pyosalpinx rising nearly up to the umbilicus, removed through the abdomen.

With regard to the last objection—that some pus sacs may be left unopened, I will deal with it when I speak of the technique of the operation itself.

I have operated upon six cases of pyosalpinx by the vaginal route; not a large number, I grant; but the results have been so satisfactory, and the convalescence has been so easy and quick that they are quite sufficient to convince me of the advantages of this route. They are all perfectly well now, they have no abdominal scar, no necessity for wearing a belt, and run no risk of getting a ventral protrusion.

Table of Cases.

Initials.	Married or Single.	Date.	Place of Operation.	Disease.	Operation.	Result.
A. C.	S.	Nov. 29, 1894	Newhall Street	Pyosalpinx and localised pelvic abscess	Incision and drainage from vagina	R.
H. M.	M.	Jan. 9, 1897	Women's Hospital	Double pyosalpinx	Removal per vaginam	R.
L. G.	M.	Jan. 18, 1897	Women's Hospital	Large left pyosalpinx with myoma uteri	Vaginal hysterectomy and drainage	R.
E. C.	M.	Mar. 4, 1898	Dudley	Large left pyosalpinx	Incision and drainage per vaginam	R.
A. F.	M.	Mar. 20, 1898	Women's Hospital	Large left pyosalpinx	Incision and drainage per vaginam	R.
E. M.	M.	July 6, 1898	Moseley	Double pyosalpinx	Removal per vaginam	R.

In addition to these there are 4 cases of pyosalpinx treated by my colleagues by this route on record in the Women's Hospital statistics.

The first of these cases was operated upon as far back as

* Read at a meeting of the Midland Counties Branch of the British Medical Association.

November, 1894. If I had known then as much as I do now I might at a much earlier date have looked upon this method as an operation of election. In that case it was one of necessity.

It was a case of pyosalpinx of long duration. The woman was extremely emaciated and pulled down by the disease. There was a large tender swelling above and behind the uterus. This I attempted to remove by abdominal section on November 21st, 1894, but on opening the abdomen I found the whole of the pelvic inlet shut off by coils of intestine densely adherent to one another. It was absolutely impossible to separate them so I closed up the abdomen. Four days later I tapped the mass from the vagina and drew off only about 2 drachms of pus. Her condition did not improve, and I felt certain there was more pus there. Under an anæsthetic on November 29th I made an incision behind the cervix, and cutting upwards behind the uterus I opened Douglas's pouch; with my fingers I could separate to a slight extent the adhesions between the floor of Douglas's pouch and the pyosalpinx. I then cut into the pyosalpinx and about 3 ozs. of pus escaped. I enlarged the opening and packed it with gauze. This I removed in 48 hours, and employed irrigation for the next few days. The cavity soon granulated up, and her recovery was complete.

I was so firmly impressed with the idea that the tube should have been removed through the abdomen that although she was cured I looked upon the case as one in which I had failed to perform the ideal operation. It was a failure on my part not to have seen that this very treatment to which I was driven as a matter of necessity was the very method that I ought to have adopted from the beginning. I have no doubt now that when I tapped her I simply tapped a small abscess in Douglas's pouch.

What we should do instead of insufficient tapping, is to make a free opening into Douglas's pouch behind the cervix. It is very easily done with scissors curved on the flat, keeping the point of the curve towards the uterus. If the pyosalpinx is very adherent to the floor of Douglas's pouch it will be opened straightway, if not the exact relations of the parts, using if necessary gentle bimanual examination, can be made out. When the pus sac has been defined it must be incised and the pus let out. Then the opening is enlarged and the cavity explored. If in doing this bimanual examination is employed I do not think any pus sacs will be left unopened. After this iodoform gauze packing is all that is required, followed afterwards by daily irrigation and in some cases by light repacking. This is the procedure I have adopted with the happiest results. The cases are few in number, but it does not necessitate a large number of cases to convince one of the advisability of a new procedure, if it did there would never be any attempt however slight at improvement. I do not pretend for one moment that this method is original. My first case was certainly done before I knew of such a proceeding having been adopted by anyone else, but as I have said I did not learn the lesson from it that I ought to have learnt. This method has been largely practised by French and American surgeons and to some extent in this country. I believe, however, that it is not sufficiently known that patients distinctly in a bad condition to stand a prolonged abdominal section can be not only relieved but cured by an operation that takes only a few minutes, that does not involve exposure or fouling of the general peritoneal cavity and which therefore must entail less risk of peritonitis and be followed by an easier convalescence.

I repeat that I do not set this up, nor has it been set up, except by a few enthusiastic Americans, as in any way a rival to removal by abdominal section as the treatment for pyosalpinx. It is to my mind a valuable auxiliary operation; one which will, I think, enable us to save lives which otherwise would be lost. This method of treatment may be extended even further, and in 2 out of my 6 cases I have succeeded in completely removing both pus tubes by the vaginal route. In these 2 cases removal would have been effected more easily from above. They were both rather stout, they both absolutely refused to have the abdomen opened, but consented to an operation by the vagina. I expected to have some difficulty in incising and draining the pus sacs in these cases, as they were not large enough to fill up Douglas's pouch. They were, however, low down, and after opening Douglas's pouch I found I could remove them. Removal of the appendages *per vaginam* is not an easy operation in any case, and where adhesions have to be separated and a soft, friable tube and ovary pulled down into the vagina great care is required.

I am not prepared to discuss now the performance of vaginal hysterectomy as part of the operative treatment in

cases of double pyosalpinx. I do not consider that we have sufficient evidence yet to show clearly whether the climacteric disturbances are greater after hysterectomy than after removal of the appendages or *vice versa*. As long as we can cure our patients either by removal of the appendages through the abdominal incision or by vaginal incision and drainage, or even by removal *per vaginam*, then I think myself that the cases which will call for vaginal hysterectomy are very few. I grant that I have one in my few cases treated by the vaginal route.

This was a case of large left pyosalpinx plus an interstitial myoma of the uterus. Before I performed vaginal hysterectomy I had attempted removal of the appendages from above, but in trying to separate the bowel adherent to the top of the pyosalpinx I nearly tore it across. As there were several other adherent coils of intestine, I contented myself with suturing the tear in the bowel and closing the abdomen. Some weeks afterwards I performed vaginal hysterectomy and drained the abscess cavity of the pyosalpinx through the opening made in the removal of the uterus. She made a good recovery.

Before concluding, I wish to repeat that it is only in a limited number of cases—in those which may be considered as of the severest kind, having as a rule a long-standing history of gonorrhœa—that I advise operation by the vaginal route. I may have to alter my views again. I can imagine that we shall find some day that all cases of pyosalpinx should be treated from the vagina. I can also imagine—though I hope it will not be so—that the closed pus sacs, tubal or tubo-ovarian, may some day give rise to further trouble, and we shall wish we had removed them entirely. If so, then I shall stand convicted of making a mistake now. But if I am told that they will fill up again with pus because the sacs which were tapped years ago refilled, I shall not believe it. This laying open of the sac and drainage is a very different thing to tapping.

Because I wish for this method of treatment to be adopted for the class of case that I have mentioned, I have not lost my admiration of the work done by the men who laid down the foundations on which the treatment of pyosalpinx is built. Without their experience, their teaching, and their boldness of treatment it would never have entered into the mind of anyone to remove these pus sacs through the vagina.

ON THE VALUE OF GLOVES IN OPERATIVE WORK, WITH A NOTE ON THE CLEANSING OF THE HANDS.*

By J. LYNN THOMAS, F.R.C.S. Eng.,

Assistant Surgeon to the Cardiff Infirmary; Consulting Surgeon to the Porth Hospital, to the *Hamadryad* Hospital, and to the Cardiff Provident Dispensary.

I PROPOSE in this short communication to place before the members of this Branch the impressions gained by the wearing of gloves in general operative surgery since February, 1898, and trust they will be of some interest and value.

"The use of gloves is of high antiquity," for it is recorded that Cyrus, King of the Persians, about 2,500 years ago, went without his gloves.

The end of the nineteenth century has seen the introduction of gloves into "certain kinds of manual labour" after the fashion of the ancient Greeks and Romans, but history does not relate what particular kind and form of glove were used by them.

The costumes which were worn by our forefathers to keep away the plague were not complete without the gloves, and the *fin de siècle* surgeon tries to ward off his plague—the pathogenic organism—by putting the final touch to his costume by wearing the microbeless glove.

When one of our great authorities on gloves stated that "yarn, thread, silk and cloth gloves are of comparatively little importance," he little dreamt that these despised articles of wearing apparel would so soon come to play an important part in the saving of human life.

The whole aim of the modern surgeon is to prevent his ubiquitous foe from establishing an "effective occupation" within his sphere of influence, and an old-fashioned custom

* Read at Aberdare before the South Wales and Monmouthshire Branch of the British Medical Association.