A LECTURE ON POLYMYOSITIS.

Delivered at the National Hospital for the Paralysed and Epileptic.

By SIR W. R. GOWERS, M.D., F.R.S.,

Physician to the Hospital; Consulting Physician to University College Hospital.

GENTLEMEN,—I desire to direct your attention to-day to a rare disease, which is important because it is a formidable malady and, though uncommon, has close relations with others which are more common and more familiar. It is a disease that has been termed "polymyositis." It consists in the simultaneous inflammation of many muscles and of some nerves—an inflammation that is symmetrical in its distribution. I am not sure that it has yet obtained a place in English descriptive medical literature or in the systematic teaching on which you chiefly depend for your knowledge; but the case I have to show you is an extremely well-marked example of the affection in a severe degree, and may therefore serve to impress on you its most characteristic features.

Let me first remind you of some facts relating to polyneuritis, with which this affection is closely allied, and which, indeed, forms part of it; the nerves suffer in association with the muscles, although, it would seem, less widely. You are all, no doubt, acquainted with "multiple neuritis," comparatively new to our nomenclature, but doubtless, beyond comparison, old in undiscerned familiarity. Its causes must have been always with us, and their effects cannot have been absent. The names we apply are not entirely appropriate. We call it "polyneuritis," "multiple neuritis," and "peripheral neuritis." These terms, however, are also applicable to other affections. We meet with inflammation of many nerves of quite different character and distribution. The designation "peripheral neuritis" has also been otherwise used, and it describes three features of inflammation of nerves—the fact that the affection is chiefly localised in their peripheral extremities; the fact that it affects chiefly the peripheral extremities of the limbs; and it has also been employed as a designation of affections of the nerve trunks in distinction from central disease. No term indicates that which is the chief characteristic of the disease, its bilateral symmetry. The motor nerves may be most conspicuously involved, or the sensory nerves may be primarily disordered—excited to spontaneous sensations or to hyperæsthesia—or they may be impaired so as to produce diminished sensibility, but in all cases the symptoms are bilaterally symmetrical. The symmetry is not always absolute; one side may be affected somewhat earlier, or in rather greater degree, than the other; but the correspondence of the affection of the two sides, whether it be in the arms or legs, or both, is the dominant feature. The two seldom present conspicuous differences, either in the area involved or in the character of the symptoms, and the correspondence is often remarkably precise.

This feature is not only the most characteristic; it is also, like most real characteristics of disease, the most significant. It indicates the nature of the cause of the affection. It shows the presence of a cause which is able to act upon structures of similar susceptibility on each side, that is, of a cause reaching all parts but acting only on those which present a special disposition to suffer from its influence. Bilateral symmetry, the correspondence on the two sides of the affected structures, is a feature which is far more than superficial. Not only does it involve correspondence of external conformation of the symptoms; it involves also correspondence of the incidence of the malady and of the vital characteristics of the corresponding structures that are involved. It implies a like susceptibility to morbid influences, and a correspondence in the effects of agents which are able to reach the two sides and to exert their influence, whatever that may be, on the similar structures. For any agent to act in the same manner on these distant but corresponding parts, it

must reach them by the only path to which they are equally accessible, namely, the circulation.

It is true we may have a bilateral correspondence in consequence of another process. Similar tissue elements on the two sides may fail in their nutrition because they possess an inherent defective vitality. That which we may call the vital endurance of the tissues is sometimes less in some structures than in others, and in consequence those which have less resistance slowly decay before the rest of the body, fail prematurely in their nutrition and in their function. Such failure is generally a senile change, although by a strange defect of local vitality it is sometimes met with soon after adult life is reached. In these cases, however, the failure of nutrition is an extremely gradual process, and the symptoms are correspondingly slow in their evolution. In the cases with which we are now concerned the symptoms are more or less rapid in their development. Such rapid failure of nutrition and disturbance of function in corresponding structures on the two sides may be safely taken as an indication that a toxic agent exists in the blood, an agent which is carried to all parts, but acts only on those which are specially susceptible to its influence.

Remember this great fact of the bilateral symmetry of symptoms, because it is of extreme practical importance. In the more common forms of the disease of the nerves, due to a toxic agent in the blood, the influence is exerted on the motor fibres, and the result is a symmetrical paralysis of the muscles, especially of the extremities, those that extend the wrist and fingers, or the homologous muscles which flex the ankle-joints and extend the toes, or of both. But not rarely the chief incidence of the toxic agent is on the sensory nerves, causing first subjective sensations in the extremities, and then, if severe, producing diminution and even loss of sensibility. In such cases there is commonly also loss of the knee-jerk. This indeed commonly results from the affection of the motor nerves, but in sensory neuritis it apparently depends on an early interference with the conduction of the afferent impressions through the sensory muscle nerves. There may even be definite ataxy, probably as a result of a greater degree of the same process.

The number of toxic agents which are capable of thus influencing the function of the nerves is great and their nature is most varied. Foremost among them are substances received from without—metallic poisons and some organic poisons, as alcohol. Some of the latter result from the growth of low organisms, and probably other substances of a more or less analogous origin, but of uncertain nature, having the same effect. A similar pathogenic influence may be exerted by poisons produced within the body. Some of these are the result of the growth of bacterial organisms, introduced from without, but generating the toxin, directly or indirectly, in the course of their development within the body. It seems to be in this manner that the polyneuritis occurs with which we have become familiar as the sequel of many acute specific diseases. But besides the poisons of this class, we have evidence of others. The disordered metabolism which takes place as the result of constitutional conditions, such as gout, seems occasionally to result in such toxic agents, the effect of which may be greater in consequence of some predisposition to suffer on the part of the individual, a state of general imperfect nutrition, and therefore of imperfect resistance to deleterious influences, whatever be their nature.

In other conditions there seems to be a state of the system which disposes it to react in a peculiar manner under external influences, so that a special toxic agent is produced under the influence of an external cause which we cannot even associate with a bacterial agency—such a cause as the exposure to cold. You know how various and grave may be the effects of this exposure. It may cause simply tonsillitis; it may cause rheumatic fever with all its manifestations; it may cause pericarditis alone. How the exposure produces its consequences we cannot yet surely discern, or indeed discern at all, beyond the fact that we have no reason to suspect the introduction of toxic agents into the system from without, and we are therefore thrown back on the assumption that the chill induces some disorder of the chemical processes within the body. These are always going on; they constitute the source of all the manifestations of vitality; but their normal course depends on a precise adjustment of many influences, some of

which seem readily to be disturbed. Remember, moreover, that the slightness of a deranging cause may be no measure of the degree of the derangement produced, nor of its extent. In a complex organism, such as our frame, one process set wrong acts upon others, and there seems to be a mutual action and reaction of the interrelated processes, so that a simple wetting of the feet may lead to a grave general disease.

It is a trite observation that but a narrow boundary separates life from death. An equally narrow one separates health from disease, and the same truth may be traced downwards in all the subsidiary processes, the sum of which makes up health and the condition of normal life. The change need often only be small which converts a material that subserves normal nutrition into one which is capable of arresting tissue life. You will remember the researches of Martin on diphtherial paralysis which have shown that a small quantity of a ferment produced by the diphtherial organisms in the throat will set wrong the chemical processes in the spleen and elsewhere concerned in the elaboration of the albumoses for the nutrition of the body. The change is probably one of slight degree, but it results in the production of a poison capable of paralysing the nerve structures of the body, and so leading to that sequel of diphtheria which is only less terrible than the disease itself. These considerations enable us, if not to understand, at least to conceive, that so simple a process as exposure to cold may result in the production of diseases of various and grave character, some of which we are compelled by all analogous facts to refer to toxic material produced in the body by deranged chemical processes.

Such toxic agents seem to have a more varied influence than the definite poisons received from without, and this is true of those that are due to exposure to cold. Among their effects is the polyneuritis of which I have spoken. One feature presented by this form is its frequent severity and acuteness in some cases, chronicity in others. Another important fact is that to which I desire especially to direct your attention to-day—that the inflammatory process may involve the muscles as well as the nerves, and even more widely than the nerves, producing the malady termed polymyositis. This result is met with almost exclusively as a result of exposure to cold. It seems to be a peculiar variety of the rheumatic poison produced in specially susceptible individuals, rendered such by some influence which depresses the general health. In such cases of multiple inflammation of both the muscles and the nerves, the affection presents the general distribution of polyneuritis so far as preponderant loss of power is concerned, but with a far wider implication of the muscles, of which many suffer that escape in ordinary polyneuritis. They are at first very tender, and afterwards undergo hardening and contraction, which may be extreme in degree; and indeed, after a time, may be insuperable, and resist all efforts to overcome it. It is an example of this disease that I have to show you to-day. It is the late stage of the disease which this patient presents, but its results are only too conspicuous, and its late stage may not render it the less instructive. She has been under treatment here for many months, and has improved very much in some parts, although in others we have been less successful. I had better give you the account of her history, which is contained in the hospital notes.

The patient is a married woman, now 36 years of age, the wife of a labourer. She was admitted here ten months ago with almost universal palsy. Her father had suffered twice from rheumatic fever, and his father was the subject of gout. This should be noted; it illustrates the fact that the majority of those who are liable to rheumatic fever are the children of a gouty parent. It is an important fact, and one that curiously illustrates the manner in which important causal facts elude discernment, until they are specially looked for. If you are on the look-out for it you cannot fail speedily to obtain evidence of its truth from your own experience; and yet few persons, surprisingly few, have noted it until their attention was directed to the sequence. It is one of the many points of connection between gout and articular rheumatism. Some change in the process of hereditary transmission seems to convert the gouty diathesis into what we regard as a purely rheumatic diathesis, which finds its expression in the most concentrated form of rheumatism, rheumatic polyarthrititis.

This by the way. The patient had the hard-worked life of women in her class. She occasionally took spirits, but close questioning failed to elicit evidence of excess. The affection came on after the death of her mother, whom she had nursed sedulously, with the result that she was constantly over-tired and often exposed to cold at night. The first symptoms were a constant tired feeling and aching pain in the back and around the loins. An irritable rash appeared over the hands and arms. Such a symmetrical eruption is always proof of a morbid state of the blood. Then she began to have pain in the hands and also in the ankles, and both hands and feet became weak. These symptoms slowly increased during two or three months, and in October, 1896, the feet had dropped and she could not get upstairs, and had difficulty in sitting upright. By the end of that year the arms and legs and trunk had become much weaker, and she was taken into a general hospital for seven weeks. At the end of that time she could not stand, and the limbs and trunk were getting stiff so that she could not bend forwards. In a sick asylum, to which she went next, and where she remained eight months, the stiffness steadily increased, and the arms and legs became rigid in the position of flexion, but the pain became slighter.

When admitted here she was so helpless as to be unable to sit up; she was continually perspiring, and complained of shooting pains in the shoulders. The only symptoms in the region of the cranial nerves were in the movements of the face and jaws. The facial muscles acted slowly and imperfectly, apparently from stiffness, and there was contraction of the masseters, which felt firm, and their rigidity prevented the mouth being opened widely. Rigid contracture existed in all the muscles of the neck, limiting its movements. The head was quite stiff upon the trunk, and hardly any flexion of the neck was possible, obviously from the muscular contraction. The same rigidity existed in the muscles of the back, and she could not bend the trunk either backwards or forwards. She had to be moved in bed as one stiff piece. The same condition of rigidity and contracture was most conspicuous in all the muscles of the upper limbs. Slight voluntary contraction could, however, be produced in most of them. But the arm was adducted by contraction of the pectoralis, and passive movement could not draw it from the trunk beyond an angle of 45° . The elbow was flexed, and contraction of the biceps prevented its extension beyond an angle of 60° . The hands were dropped at the wrist, and the fingers flexed and could not be extended completely. All the muscles of the arms and hands were much wasted, but although small they were extremely hard and firm; this was the case in both flexors and extensors. Voluntary movement was abolished in the extensors of the wrist and long extensors of the fingers, but only a very slight degree could be produced in the other muscles of the arms. All the muscles reacted very slightly to strong electrical stimulation, the faradic shock, or the interrupted voltaic current. The response to the latter was normal, so far as the poles were concerned, and it had the sharp, quick, character which indicates, like the faradic irritability, stimulation of the intramuscular nerves and not of the muscular substance itself. The latter contracts with a peculiar sluggishness, which presents a marked contrast with the instant response when even a few normal nerve fibres subserve. Sensation on the arms was quite normal. The rigidity prevented any reflex response to stimulation.

Both legs were extremely rigid, and all the muscles were small and hard. The feet could not be flexed. Neither the hip-joints nor the knee-joints could be passively extended, on account of the contraction of the flexors, and that in the extensors of the knees prevented flexion to more than 15° degrees from the position of complete extension. She had, however, just enough power over the flexors of the hip to raise each foot from the bed, but she could not flex the knee or move the foot at the ankle-joint, although the muscles could be felt to contract slightly on a voluntary effort. The contraction in the calf muscles was so great that the feet could not be passively brought nearly to a right angle with the leg.

In all the muscles of the legs and feet response could be obtained to both faradism and voltaism, but the contraction was slight which could be obtained even with a strong current. The polar reaction was normal in the legs also, except

in the muscles of the calves, in which the A.C.C. was obtained more readily than the K.C.C.

Sensibility in the lower limbs was everywhere normal. No pain or subjective sensation was complained of, but there was some pain in the back. The deep reflexes could not be anywhere obtained. There was no affection of the sphincters. A curious symptom was the excessive amount of perspiration, both on the face and limbs. Apart from exertion the sweat would often stand in beads on the surface.

That was her state on admission. Unfortunately the affection had then reached an advanced and chronic state. The inflammatory process in the muscles had attained its cicatricial stage, in which the newly-formed connective tissue, almost universally distributed, had already contracted. We know, from varied experience, in many places, and in many forms, how strong is the contracting tendency of this abnormal structure. In this patient we have striven to overcome the contraction by passive extension and by massage, and have endeavoured to increase the strength of the muscular fibres by movements and electricity as well as by therapeutic agents. I cannot say that much change has been produced in the arms. But if our success in improving the condition of the arms has been small, we have at least the satisfaction of having done good to the legs and back. Their condition is far better than when she came under our care. We have succeeded in overcoming the contraction of the hip and thigh muscles by passive extension, and so have enabled her to use the limbs, with the result that there has been a great increase of their power. The inability to stand has been exchanged for the ability not only to stand, but also to walk. There is still some contraction in the calf muscles, preventing full passive flexion of the ankle joint, but this residual amount will soon yield under perseverance in standing and walking, the most potent of all agents for the purpose. The state of the arms you can observe. In spite of variously contrived efforts to obtain the extension of the elbows, as you see, the contraction of the biceps prevents the extension of the forearm. That of the flexors of the wrist prevents the full extension of this joint, and the abduction of the upper arm is hindered by the contraction of the adductors, and especially of the pectoralis major. Note how the rigidity of this muscle prevents the movement of the arm far from the side. The deltoid, the chief abductor, is rigid and firm, and presents much less response to the faradic stimulus than does the pectoralis. You may observe the same relation between the voluntary power and the degree of shortening in other opposing muscles, in the biceps and triceps, and also in the forearm muscles. She can produce only very slight contraction by the will in the extensors of the wrist and fingers, while there is considerable voluntary power in the flexors. The electrical irritability agrees in the degree of response to the voluntary power. Accordingly, though both sets of muscles are small and hard, those which are least paralysed are most contracted. We may see in this the correspondence of this affection and polyneuritis. There is the same flexor contraction and extensor weakening in both sets of muscles, although all are involved in the cicatricial inflammatory changes. We must therefore infer that there is polyneuritis as well as polymyositis, that both nerves and muscles are involved in the morbid process; but the affection of the muscles is far more extensive than it is in the pure affection of the nerve. How widely and profoundly the former have been involved is shown not only by the degree of hardening and contraction, but by its extensive distribution. The affection of the facial muscles still exists; you may note not only its stiff aspect, but the small amount of movement which results from voluntary or emotional influence, for still more evident is the contraction of the masseters. As you can see, she cannot open her mouth to much more than half the normal extent, and if you presently feel the muscles as thus extended you will note how hard they are and how distinct it is that their contraction is the agency which prevents a greater depression of the jaw.

I do not doubt that her power of walking will improve to a useful degree. Its influence on the contraction of the calf muscles is, as I have said, steady and maintained. It is surprising how much contracture may be thus overcome in the course of time, even a degree that, at first, seems to permit the expectation of release only from tenotomy. But we have

no means to effect the extension of the arms of similar potency. I fear we shall not be able to secure her ability to use effectively what power she possesses in the hands without having recourse to division of the tendons of the biceps. Once developed, such contraction of the flexors of the elbow is a most formidable condition. The extensors act at great disadvantage when the joint is flexed, while the influence of the flexors is then at its maximum, while we have no habitual movement of which to avail ourselves, comparable to that of the weight of the body on the contraction of the calf muscles.

This therapeutic inability brings me to the last point which I desire to impress on you, and like many last words it is the most important. The scope for treatment, at this stage of the affection, is small. If you picture to yourselves the pathological condition, you must conceive the muscles as now containing only a small proportion of normal fibres, and that these are separated by fibrous tissue which has undergone cicatricial contraction. Many muscular fibres have perished in consequence of the acute inflammation and of the degeneration of their nerve fibres. The primary disease had already ceased when she came here, and only its effects remained. You know, in all diseases, what a different task it is to treat the primary affection and to treat its late results when these consist of grave structural damage. So far as recovery of the partially damaged muscles is concerned, these have probably regained as nearly a normal condition of the remaining fibres as they can reach. Some increase of power, however, may result from their voluntary use, if this can be increased by getting rid of the disabling shortening which interferes with all effective employment of the arms. But in this, as in so many diseases, the only opportunity for really effective treatment is that which is presented in the early period of the case, when the malady is in the stage of development. I cannot doubt that, in this patient, rest, diaphoretics, and salicylates, with perhaps small doses of mercury, would then have controlled the inflammation, and lessened very much the subsequent mischief which it has induced. Every degree of diminution of the intensity and duration of the active processes of disease is multiplied many times in the ultimate diminution which is thus caused in the later and permanent condition. I emphasise the fact, because to some of you may come the opportunity of treating such a case in the only stage in which treatment can be really effectual, and because the general lesson is applicable to many maladies of many kinds.

A CLINICAL LECTURE

ON

A CASE OF UNILATERAL (HEMIPLEGIC) SPASM AND CONTRACTURE WITHOUT PARALYSIS.

Delivered in the University of Edinburgh,

By SIR T. GRAINGER STEWART, M.D., LL.D.,

Professor of the Practice of Physic, Physician-in-Ordinary to the Queen for Scotland, and President of the British Medical Association.

(Concluded from page 5.)

THE outstanding features of the case as explained in the part of the lecture already published are:

A soldier, aged 25, had severe typhoid in India; on recovering from coma, found himself unable to use his left hand on account of stiffness of the fingers. Gradually they became flexed, and the arm, foot, leg, and left side of face became rigid with constant hypertonicity and frequent tonic spasms. This condition makes him resemble a case of hemiplegia with descending sclerosis, but there is no paralysis; and all the muscular disability is due to antagonistic action of opposing muscles. Otherwise at present no abnormality of any system or of the nervous system in any respect, except the unilateral spasm. The locomotory system is interfered with only in so far as the spasm, the contracture, and the consequent changes in joints and bones are concerned. He has, since his illness began, had two attacks of mania.

It should be particularly noticed, as to the state of the muscles, that the deep reflexes are exaggerated, and that, although the electric relations differ on the two sides of the body, there is no trace of the reaction of degeneration.