cxvii, p. 594. (iv) Da Costa, Large Fibroid Tumour of the Anterior Wall of the Vagina, *Medical News (Philadelphia)*, 1895, vol. lxvii, p. 463. (v) Caswell, Fibroma of the Vagina, *Boston Med. and Surg. Journ.*, 1883, cix, p. 151. (vi, vii) Gaye, *Berl. Klin. Woch.*, 1882, No. 43, p. 652. (viii) G. Braun, Gebrauch eines bei Extirpation eines Fibroms der vorderen Vaginalwand benützten Spekulum, *Wien. med. Woch.*, 1885, xxxv, pp. 1557-9. (ix) Lewers, *Obstet. Trans. (London)*, 1887, vol. xxix, p. 299. (x) Tillaux, Fibro-myome de l'Utérus, *Annales de Gynécologie*, 1889, vol. xxxii, p. 161. (xi) Herman, *Obstet. Trans. (London)*, 1880, vol. xxii, p. 44. (xii) Stumpf, Myofibrom der Vagina mit beginnender Verjauchung, *Münch. med. Woch.*, 1890, xxxvii, p. 694. (xiii, xiv) P. Strassmann, Zur Kenntnis der Neubildungen der Scheide, *Centrabl. f. Gynäk.*, 1891, p. 825. (xv) Hofmök, Entfernung eines Kindskopfgrossen Fibromyom der vorderen Vaginalwand durch elastische Ligatur, *Wien. med. Presse*, 1891, xxxii, p. 1229. Hasenbalg, Fibröser Polyp der Vagina, *Zeits. f. Geb. u. Gynäk.*, 1892, xxiii, p. 52. (xvii) A. Breisky, *Handbuch der Frauenkrankh.*, Bd. iii, p. 731. (xviii, xix, xx) Hastenpflug, Ueber Vaginal-Myome, *Inaugural Diss. Jena*, 1888. (xxi) C. Cleveland, *New York Journal of Gyn. and Obstet.*, 1893, vol. iii, p. 154. (xxii) Wernitz, *Centrabl. f. Gynäk.*, 1894, No. 26, p. 632. (xxiii) Thornton Parker, *Virginia Medical Monthly*, Richmond, 1886-7, xiii, p. 793. (xxiv) Alfred C. Godfrey, Fibro-myomata of the Vaginal Wall, *Colorado Medical Journal*, 1896, No. 9, p. 284. (xxv) J. C. Moore, *St. Louis Med. and Surg. Journal*, 1885, vol. xlvi, p. 125. (xxvi) R. Condamin, Note sur une Observation de Fibrome Juxta-urétral du Vagin, *Archives Provinciales de Chirurgie*, Paris, 1894. (xxvii) Paquet, Tumeur Fibreuse du Vagin, *Bulletin Médical du Nord*, 1882, No. 2, p. 65. (xxviii, xxix) John Phillips, present article.

AFTER-EFFECTS OF ABDOMINAL SECTION.

By MAY THORNE, L.S.A., M.D. BRUX.,

Assistant Anæsthetist (late Senior House Surgeon), New Hospital for Women.

At the meeting of the Obstetrical Society held in October, 1897, Dr. Herbert Spencer pointed out that intestinal obstruction from the adhesion of the intestine to the stump after ovariectomy was by no means rare, and a month later at the Harveian Society Mr. Doran drew attention to the fact that hernia of the abdominal cicatrix frequently followed laparotomies.

These two important statements are of such interest to all practitioners that it has been my endeavour to gather together as accurately as possible statistics of these complications in women patients from the records of general hospitals throughout the country. Thanks to the great courtesy of members of the staff and of the surgical registrars of 49 hospitals, the following information has been obtained. Several country hospitals were invited to send statistics, as it seemed probable that cases of acute intestinal obstruction would be treated in them, even though the primary operation had been done in a hospital in some large centre, as London, Edinburgh, or Dublin.

During the last four years 39 cases of ventral hernia and 12 cases of acute intestinal obstruction have been traced.

The cases of ventral hernia do not include any that may have been seen in the out-patient departments, but only those that came into the wards for treatment. The date of the occurrence of a ventral hernia or of acute intestinal obstruction after the original operation is of interest as showing approximately the length of time that the giving way of scar tissue or the formation of fibrous bands may take. It has not been possible to get all the data, but in 26 out of the 39 cases of ventral hernia they are as follows:

2 cases occurred 14 years after the original laparotomy

1	"	10	"	"
1	"	9	"	"
1	"	8	"	"
1	"	7	"	"
2	"	5	"	"
3	"	4	"	"
3	"	3	"	"
1	"	2½	"	"
2	"	2	"	"
2	"	1½	"	"
3	"	1	"	"
1	"	5	"	"
1	"	4	"	"
2	"	a few months	"	"

It would be interesting to learn in what manner the abdominal wound had been closed in these cases, whether in one or more layers, the number of cases in which a drainage tube was used, the length of rest in bed, and the use of an abdominal belt for months or years following the operation. Eighteen of the cases occurred, it will be noted, under five years, and seven in one year or under.

The dates of the cases of intestinal obstruction were as follows:

1	case occurred	14 years	after the original laparotomy
1	"	5	"
3	"	4	"
1	"	1	"
1	"	6 months	"
2	"	2	"
1	"	15 days	"
1	"	11	"
1	"	10	"

The two cases which occurred two months after abdominal section were (a) a case of perforated ulcer of the intestine, and (b) a case of sarcoma of the ovary. In neither case, therefore, was the peritoneal cavity in as sound a state as is found in the majority of cases of abdominal section, and one cannot but question whether sepsis was not responsible for the cases occurring within fifteen days of the operation.

Dr. John Phillips gives an account of 3 cases of acute intestinal obstruction after abdominal section, the obstruction occurring one, four, and seven years after the original operation.²

I am greatly indebted to Mrs. Scharlieb, M.D., M.S., and to Mrs. Stanley Boyd, M.D., for permission to look through the casebooks at the New Hospital for Women, in order that I might trace the after-histories of patients who had had abdominal section performed there.

Information has been received of 88 patients, and, amongst these, 3 have returned for further treatment—1 for acute intestinal obstruction and 2 for ventral hernia.

The history of these 3 cases is as follows:

Mrs. T., aged 22, was operated upon in 1892, when a mass was removed which consisted of the right ovary, considerably enlarged, and containing several abscess cavities and an irregularly distended long sac, which appeared to be the Fallopian tube. The patient was very ill, with high irregular temperature for some time, but eventually she completely recovered, and left the hospital on the fifty-fifth day after the operation. From that time to January, 1897, her health was good. On January 12th, 1897, she complained of pain in the abdomen; on the 15th she took to her bed at home, when sickness began and was frequent. No motion or flatus passed after the 18th. Her medical man advised her going into a hospital, and on January 21st she was brought to the New Hospital for Women with all the symptoms of acute intestinal obstruction. She was almost moribund on admission, and died two hours later, no operation being performed. At the post-mortem examination the intestines were found much distended and reddened, there were recent adhesions between the parietal and visceral layers of peritoneum, and also between the coils of large and small intestine. A firm white band attached the centre of the transverse colon to the back of the anterior abdominal wall, causing a bend and slight constriction in the colon. A loop of small intestine about 13 feet from the duodenum hung down into the pelvis, and was constricted by a narrow white fibrous band. The band passed round the loop of intestine, and was continuous at one end with the mesentery, and at the other with the peritoneal coat of intestine just beyond the part involved by the loop. The end of the ileum was found adherent to the abdominal wall in the neighbourhood of the iliac fossa and the stump of the right broad ligament was fixed at the same spot. The cæcum was bound down by dense pale adhesions. There were adhesions between the uterus and bladder. The left ovary and tube were fixed by adhesions to the pelvic wall in the neighbourhood of the iliac vessels.

The adhesions in this case were dense and numerous. The question arises as to what adhesions were formed before the operation in 1892 was undertaken. The tumour removed contained abscess cavities, and one might reasonably suppose that inflammatory conditions were present in the neighbourhood of the right ovary, which was removed. The patient's convalescence was long, and gave rise to much anxiety, and possibly it may have been during this time that adhesions were formed, which subsequently became dense and fibrous.

The two cases of ventral hernia are as follows:

Mrs. L., laundress, aged 45, very stout, was operated on for double ovariectomy in 1888; she made a good recovery, and kept in excellent health, and did her work till 1894, when she first noticed a swelling in the middle of the scar about the size of a walnut. This increased in size, and she suffered from gripping pain till she returned to the hospital in 1895, when she was readmitted. At the operation it was found that the omentum was extensively adherent to the parietal peritoneum under the scar. The patient made an excellent recovery.

A. D., a dwarf with contracted pelvis, aged 30, had Cæsarean section performed at full term in 1895. She recovered well. She was readmitted in 1896, suffering from profuse losses, when an abdominal hysterectomy was performed. In 1897 she returned with a ventral hernia in the middle of the scar, which she was anxious should be dealt with. At the operation several coils of intestine were found adherent to the parietal peritoneum and to one another. She made a third good recovery, and is now in excellent health, earning her own living.

The following tables give the after-history of the patients who have been traced:

REFERENCES.

¹ *Lancet*, Novem^r 27th, 1897. ² *Lancet*, 1892.

Single Ovariectomies.

No.	Date.	Imme- diate Result.	Ultimate Result.
1	1888	Excellent	Health very good.
2	1888	"	Returned with ventral hernia in 1895.
3	1893	"	"I have not had good health; never free from pain, especially in the lower part of my back."
4	1894	"	"I have had very good health since my operation."
5	1894	"	Very good health: earns her living by mangling.
6	1894	"	"I have no pain, but much general weakness, which makes one think of all kinds of imaginary diseases."
7	1894	"	"Since my operation I have had very good health."
8	1894	"	Much better since operation.
9	1894	"	"My health has been fairly good."
10	1895	"	"I have enjoyed the best of health since my operation."
11	1895	"	Has had excellent health since operation.
12	1895	Thrombo- sis of veins of left leg	"I am fairly well in health, but my leg still troubles me very much."
13	1895	Excellent	"I have had excellent health since the operation."
14	1895	"	Health excellent.
15	1895	"	A second tumour has formed.
16	1896	Thrombo- sis of veins of leg	Health excellent.
17	1896	Excellent	"My health has been quite good since the operation."
18	1896	"	Good health since the operation; birth of healthy child 9 months ago.
19	1896	"	Health before operation poor; now health excellent; 7 months pregnant.

Double Ovariectomies.

20	1892	Excellent	Excellent health since 1893.
21	1893	"	"I have had great relief since the operation, but I am not very strong."
22	1893	"	Died of cancer of the uterus in April, 1894.
23	1894	"	Has occasional numbness down the left side.
24	1895	"	"My health has been fairly good, and I continue to get stronger. I still have hot flushes rather frequently."
25	1895	"	Health excellent.
26	1896	"	Suffers from constipation, asthma, and occasional flushings.
27	1893	Excellent	<i>Salpingitis and Ovaritis.</i> "I have had years of life and freedom from pain from the agony I endured."
28	1893	"	"Owing to the splendid operation I have been quite cured."
29	1894	"	General health fair.
30	1894	"	"Before operation never free from pain; now no pain."
31	1894	Prolonged convales- cence.	"I still suffer very considerably from weakness in the abdomen."
32	1894	Excellent	Very good health now.
33	1894	"	Excellent health.
34	1895	"	Continued pain. Second operation, when left ovary was found bound down by adhesions; this was rejected. The pain continued, and as further operation was not advised, patient went to another hospital, where she was operated upon and died a few days later.
35	1895	"	"I now enjoy very good health."
36	1895	"	"Since the operation my health has much improved."
37	1895	"	In excellent health.
38	1895	"	"Since my operation I have enjoyed very good health."
39	1895	"	Health good but pains in head.
40	1895	"	Excellent health.
41	1895	"	General health fair.
42	1895	"	"My health is much better. I am growing quite stout."
43	1895	"	Has attacks of sickness, with headache and feet swell.
44	1896	"	General health fair.
45	1896	"	General health excellent.
46	1896	"	"I am better than formerly."
47	1896	"	General health good.

Abdominal Hysterectomy.

48	1893	Excellent	"My health was weak for two years, but now I am in good health, and am manageress in a busy hotel."
49	1893	"	General health fair.
50	1894	"	"I have better health now than I had for six years before the operation."
51	1894	"	Never feels well; worries about nothing.
52	1894	"	"My general health is fairly good."
53	1894	"	"Since the operation I have been perfectly well."
54	1895	"	"The operation was most satisfactory in every way. My health has been quite good since."
55	1895	"	General health good.
56	1896	"	"I feel very well indeed."
57	1896	"	Health good. Works hard in the Salvation Army.
58	1896	"	Health excellent.
59	1896	"	General health good. Returned with ventral hernia in 1897.
60	1896	"	General health poor. Much pain in back.

No.	Date.	Imme- diate Result.	Ultimate Result.
61	1894	Excellent	<i>Fibro-cyst of Uterus.</i> "I think my operation has been fairly successful, though I am not very strong."
62	1894	Excellent	<i>Exploratory Operations.</i> Excellent.
63	1894	"	Has been in very good health since the operation.
64	1895	"	Died in 1897. Cause unknown.
65	1895	Not relieved	Died of bronchitis a year later.
66	1896	Excellent	General health good.
67	1895	Excellent	<i>Hysterectomy.</i> "Since my operation my general health has been good."
68	1895	"	Excellent health. Birth of boy 3 months ago, excellent recovery.
69	1896	"	"I have not had so much pain, but I have not felt so strong."
70	1896	"	"My general health has much improved."
71	1895	Excellent	<i>Cæsarean Section.</i> "I feel as well now as ever I did in my life."
72	1895	"	Readmitted in 1896 for abdominal hysterectomy.
73	1894	Excellent	<i>Cyst of Corpus Luteum.</i> Health excellent.
74	1894	Excellent	<i>Umbilical Hernia.</i> "Excellent till the last four or five months, when I have had some abdominal pain."
75	1895	"	"The operation was quite successful."
76	1892	Protracted convales- cence	<i>Various Causes.</i> Readmitted in 1897 with acute intestinal obstruction; died soon after admission.
77	1892	Excellent	Health better than for six months prior to operation, but cannot say it is good.
78	1894	"	Health good.
79	1894	"	Had a second operation at another hospital a year later.
80	1894	"	Health excellent.
81	1894	"	Excellent health.
82	1894	"	"My health is very good."
83	1895	"	"My health has been quite as good since the operation."
84	1896	"	Excellent.
85	1896	Malignant of perito- neum	Died 3 months later.
86	1896	Excellent	General health very good.
87	1896	"	Excellent for one year, then contracted gonorrhœa, and developed double pyosalpinx. Operated on at Charing Cross Hospital, 1898.
88	1897	"	Health excellent.

ON LATERAL CURVATURE OF THE SPINE: ITS
PREVENTION AND TREATMENT.By RICHARD BARWELL, F.R.C.S.,
Consulting Surgeon, Charing Cross Hospital.

II.

In a previous communication on this subject¹ I defined four forms of pelvic deviations, pointed out their diagnosis, and ended with a strong recommendation—almost had I written exhortation—to watch lest the youth of either sex adopt one or other of these malpostures. On these same lines I must go a little further. The spine has for its basis the pelvis; every movement of which, even though momentary, is accompanied by a compensating change in position of vertebræ; were this not so we could neither walk, run, ride, or take any sort of active exercise, because without that adaptability of the spine, the body's centre of gravity could not be shifted from side to side, backward, or forward, so as to lie over the constantly changing point of support on the floor or saddle. That is to say, when in any of these exercises the pelvis assumes for a time an oblique, a sideways, or twisted position, the spine must during that time so curve or rotate as to adapt itself to the temporary posture of its base. If, however, such posture be not merely a momentary but a permanent condition, then that curve or rotation of the spine will be equally persistent. Therefore, if a child or adoles-