able and trustworthy, can dispense with this aid and pressure, but at first it is necessary. The thickness of the cushion must correspond to the severity of the deviation. Another exercise which may be described here, though I also use it as a remedial measure when spinal curvature has already appeared, may take two different forms. For left version (1) the patient lies on her left side, the thighs flexed at a right angle to the pelvis, she then rolls the upper part of the body over to the right so that while the pelvis is still quite on the side, the shoulders, chest, and as much of the loin as possible lie almost on the back. (2) The patient lies with thighs similarly bent on her right side and rolls the upper body-segment, so that this portion reposes on the front, not merely on the chest but also as much of the abdomen as she can manage. This is less comfortable than No. 1, but more potent, and its power may be increased temporarily if an assistant press gently with the hand the left lumbar region down towards the floor.

These various measures if taken alone must be considered only as prophylactic of curvature; they can, however, be also used for incipient curves.

NOTE AND REFERENCES.

¹ BRITISH MEDICAL JOURNAL, January 21st, p. 137. ² Occasionally such an injury resulting in increased vascularity and more rapid growth has a like effect as regards uneven length of limbs, although due to excess instead of to deficiency. ³ BRITISH MEDICAL JOURNAL, 1897, vol. ii. p. 335.

ON THE VALUE OF THE TREATMENT OF DIPHTHERIA BY ANTITOXIN.

By E. W. GOODALL, M.D. LOND.,

Medical Superintendent of the Eastern Fever Hospital of the Metropolitan Asylums Board.

(Concluded from p. 200.)

II.—DIPHTHERIAL PARALYSIS.

I now pass on to the consideration of the effects of the antitoxin treatment as it is shown by the behaviour of the cases of paralysis. This is an aspect of the question that has up to the present time been scarcely dealt with by those who have written on serum-therapeutics. But inasmuch as more than one important lesson is to be derived from a study of the facts, I bring them before you to-night.

It is well known that an attack of diphtheria is apt to be followed by a characteristic form of palsy. I say followed because this palsy is more a sequel than a complication of the disease. It does not often show itself before the end of a fortnight from the onset of the local affection of the mucous membrane. It has been shown that this palsy is the expression of structural changes in certain nerve tissues, chiefly nerve endings; and that these changes are brought about by the action of toxins produced by the diphtheria bacilli at the affected mucous surface, and absorbed therefrom into the system generally. If this be true then one would expect that the larger the quantity of toxin absorbed the more likely would paralysis be to follow. Now, from clinical evidence alone I have long believed this to be true. The cases of diphtheria which are severe are those in which the paralytic sequelæ supervene. By a severe case I mean one in which there is extensive and persistent exudation, accompanied usually by certain other symptoms indicative of toxic poisoning, one of the most important of which, from a prognostic point of view, is marked and persistent albuminuria.

For clinical evidence of this I must refer those interested to a paper published in *Brain* for 1895, based on a study of the cases of paralysis occurring at the Eastern Hospital during 1892 and 1893.

Paralysis therefore follows the severe cases, and the severe cases such as I have described them are those in which the toxins are absorbed in large quantities for a considerable time. Experimental evidence points the same way. Escherich, Loeffler, Ehrlich, and Woodhead²⁴ all concur in stating that the larger the dose of toxin injected into an animal the greater the liability to paralysis. The latter states: "This paralysis occurs more frequently in animals used for testing the strength of antitoxin serum, the greater the multiple of lethal units used in the test dose—that is, there are more cases of paralysis when a theoretically ten-times lethal dose is used than when one is used, and still more when the test dose is again multiplied by ten—that is, when it equals 100."

The incidence of paralysis varies considerably in different series of cases. This is well shown by the following table:

TABLE XII.—Percentage Incidence of Paralysis in the Hospitals of the Metropolitan Asylums Board.

	1893. (Non- anti- toxin).	1894. (Non- anti- toxin).	1895. (Anti- toxin).	1896. (Anti- toxin).	1897. (Anti- toxin).
Eastern	12.1	10.8	16.0	21.4	15.1
North-Western	14.0	11.1	18.9	14.1	12.8
Western	18.1	8.2	17.7	21.5	11.0
South-Western	14.3	18.3	22.0	20.6	20.5
South-Eastern	16.2	20.2	34.7	42.3	45.9
All the hospitals ...	14.3	13.2	20.7	21.3	20.3

The average incidence in non-antitoxin cases is about 14 per cent. It is quite clear from this table that in the Asylums Board hospitals this percentage has on the whole increased since the introduction of the serum treatment. But this has not been a universal experience, as indeed Table XII shows. Professor Baginsky²⁵ of Berlin, for example, stated in a discussion on diphtherial paralysis at the Edinburgh meeting of the British Medical Association that his own observations had led him to believe that the reverse was the case. I should like to point out that the incidence of paralysis will vary in different hospitals with the average duration of the patients' stay in hospital. In some of the Continental hospitals it is usual for the patient to be sent home very soon after the local exudation has cleared up, hence, doubtless, a low paralysis-rate. With patients kept in hospital five weeks or longer, experience shows that since the introduction of antitoxin the paralysis-rate has gone up. The reason usually given, a very probable one, is that more severe cases survive the early and fatal stages of the disease, and either recover or live sufficiently long to become paralysed. But while it appears from the above facts that under the prevailing conditions we have lost ground in our endeavours to prevent this serious sequel of diphtheria, I wish to bring especially before your notice some other facts which show that if the cases be looked at from another point of view, far from having lost ground, we have made considerable advance. For the following figures I am indebted to my colleague, Dr. Wool-lacott, who has been at great pains to extract them from the hospital records. They form part of a valuable paper written by him, at present unpublished, which I have had the opportunity of reading. Dr. Wool-lacott has abstracted the notes of 1,510 cases of diphtheria treated with antitoxin, and uncomplicated by the concurrence of any such disease as measles or scarlet fever. These cases were consecutive cases admitted during 1895, 1896, and 1897.

TABLE XIII.—Showing Incidence of Paralysis in 1,580 Cases of Diphtheria Treated with Antitoxin at the Eastern Hospital during 1895, 1896, and 1897.

Day of Disease upon which Antitoxin was first Injected.	Total Number of Cases.	Total Number of Cases of Paralysis.	Percentage Incidence of Paralysis.	Number of Severe Cases of Paralysis not Fatal.	Number of Fatal Cases of Paralysis.
1st	69	4	5.7	0	0
2nd	277	28	10.1	1	0
3rd	340	53	15.5	5	3
4th	323	61	18.8	8	7
5th and later ...	571	147	25.7	28	8
Total	1,580	293	18.5	42	18

It will be noted that of these cases 293 developed paralysis, an incidence of 18.3 per cent. But contrast the incidence of paralysis in the cases treated early with that in those brought

under treatment late. When the serum was employed on the first day of the disease only 5.7 of the cases became paralysed; of those when it was used on the second day, 10.1 per cent.; and, further than this, the important fact is brought out that only one case of severe paralysis occurred amongst cases treated on the first two days with no deaths (from paralysis), and only 9 such cases with 3 deaths amongst cases treated on the first three days. Whereas, on the other hand, 51 severe cases with 15 deaths arose amongst those which were not brought under treatment until the fourth day or later. These figures clearly indicate what was hinted at by the figures given concerning the post-scarlatinal diphtheria cases occurring in the Asylums Board hospitals in 1896. Of 483 such cases treated with antitoxin, 34, or 7.0 per cent., became paralysed, a low incidence compared with that amongst the cases admitted directly from outside—namely, 21.3. But 413, or 85.5 per cent., of the cases were treated on the first two days of illness. Only 1 of the 34 cases of paralysis proved fatal.

The only figures I have which show the incidence of paralysis according to the day of admission in non-antitoxin cases are the following, which refer to cases occurring at the Eastern Hospital for the year 1894, immediately before the introduction of the serum treatment.

TABLE XIV.—*Eastern Hospital: Non-antitoxin Cases, 1894: showing Incidence of Paralysis according to Day of Disease on which Patient was Admitted.*

Day of Disease on which Patient was Admitted.	Total Number of Cases.	Number of Cases of Paralysis.	Percentage Incidence of Paralysis.	Number of Severe Paralysis Cases not Fatal.	Number of Fatal Paralysis Cases.
1st	29	3	10.3	0	0
2nd	100	12	12.0	1	0
3rd	106	10	9.4	1	1
4th	73	4	5.4	0	0
5th and later ...	144	20	13.8	3	3
Total	452	49	10.8	5	4

Compared with Table XIII the above figures show that the date on which the patient is admitted to hospital has no influence on the occurrence of paralysis. But as far as these cases go they may all be treated as early cases; unless one supposes that the patients, before admission to hospital, received no treatment at all. I append, however, a table showing the paralysis cases only, arranged according to the day of disease when the patient was admitted. Only those cases are included in which the duration of the illness was obtained.

TABLE XV.

Non-Antitoxin Cases (1892-93-94).					Antitoxin Cases (1895-96-97).			
Day of Disease when Admitted.	Number of Cases of Paralysis.	Percentage on Paralysis Cases.	Number of Cases of Generalised Palsy (not fatal).	Number of Fatal Cases.	Number of Cases of Paralysis.	Percentage on Paralysis Cases.	Number of Cases of Generalised Palsy (not fatal).	Number of Fatal Cases.
1st	12	7.7	1	1	4	1.3	0	0
2nd	44	28.3	3	5	28	9.5	1	0
3rd	31	20.0	2	4	53	18.0	5	3
4th	21	13.5	1	2	61	20.8	8	7
5th	47	30.3	6	7	147	50.1	28	8
	155	95.8	13	19	293	95.7	42	18

The 155 non-antitoxin cases, with 19 others in which the duration of the disease was not stated, occurred among 1,523 cases of diphtheria admitted to the Eastern Hospital during 1892-1894.

The figures in these tables show that the cases treated early with antitoxin are less likely to develop paralysis than those treated early without antitoxin; and also that early treatment in non-antitoxin cases does not influence the character of the paralytic attack. From the two preceding tables it also appears that though paralysis occurs more frequently with the antitoxin treatment, yet the proportion of fatal cases has not altered. Of every 100 cases of diphtheria not treated with antitoxin 1.2 died of paralysis, while of every 100 treated with antitoxin 1.1 died.

THE IMPORTANCE OF EARLY TREATMENT.

The consideration of the paralysis cases alone shows the importance of beginning the serum treatment early, but the necessity for this is also exemplified by a study of the mortality according to the day of disease upon which the treatment was commenced. This time influence is well illustrated in the Asylums Board Reports. In 1894 (non-antitoxin cases), of 133 patients admitted to hospital on the first day of illness, 30 died, or 22.5 per cent.; and of 539 admitted on the second day of illness, 146 died, or 27.0 per cent.; while for the cases treated with antitoxin in 1895 and 1896, of 143 treated on the first day of their illness, 7 died, a mortality of 4.8 per cent., and of 809 treated on the second day, 121, or 14.9 per cent., died. For 1897, the figures of all the hospitals have not been published, but I have those for the Brook²⁶ and Eastern Hospitals, which combined are as follows: 66 cases treated on first day with 1 death, or 1.5 per cent.; 317 treated on the second day with 16 deaths, or 5.0 per cent.

In the annual reports of the Asylums Board are contained another series of figures which to my mind show conclusively the value of the antitoxin treatment and the necessity of its early application. For some years past cases of diphtheria have occurred annually in every hospital of the Board amongst the patients convalescent from scarlet fever. This post-scarlatinal diphtheria, as it has been termed, has been the source of much anxiety to the Committees of Management and Medical Superintendents of the different hospitals. The chief reason for this anxiety was the terrible fatality of the disease. It was rarely below 50 per cent. at the different institutions. Of 482 cases collected by Hume²⁷ from the reports of the various hospitals for the five years preceding 1895, the mortality was slightly over 50 per cent. At the Eastern Hospital for the years 1891-94 it was 61.2, 52.9, 60.0, and 42.8. Well, in 1895 the antitoxin treatment was used in these cases. What has been the result? For that year I am unable to give the results for all the hospitals, the figures not having been published; but at the Eastern Hospital it was 25.0 per cent., at the Northern 5 per cent., at the Western 8 per cent., at the South-Western 14 per cent., and at the Fountain 6 per cent. Not all the cases during 1895 were treated with antitoxin, but even then the reduction in the fatality was sufficiently noticeable. This reduction was continued during the years 1896 and 1897, when the fatality for all these cases in the Board's hospitals was 5.0 and 4.1 per cent. respectively, the majority of the cases during these years being treated with antitoxin.²⁸

I am aware that it will be objected that a large number of these were cases in which the diagnosis of diphtheria was made merely on bacteriological grounds, and that such cases were of a very mild character. But I am fully prepared to concede a large number of them: I will let you strike out three-quarters of those which recovered, and retain all the fatal cases, even though some of them were not treated with antitoxin.

This is by no means an unfair concession, for it reduces the incidence of diphtheria on the completed scarlet fever cases for the years 1896 and 1897 to 1.1 and 1.3 per cent. respectively, a lower percentage for any year since 1888 with the exception of 1893.²⁹

What, then, is the fatality of the remaining quarter of the cases? For 1896 it is 20.1 and for 1897 15.0 per cent., and that against a previous fatality of 50 per cent. I have no hesitation in saying that the chief reason for this marvellous reduction is the fact that most of these cases were treated early, on the first or second day. The patients were inmates of hospitals and under the strict observation of medical men and nurses at the time they were taken ill with diphtheria, so that the treatment was undertaken early; and I would like to add

that whatever doubts there may be concerning the statements made about the duration of illness of patients admitted with diphtheria from outside the hospitals, for the reason I have just mentioned there can be no doubt on that score with respect to these cases of post-scarlatinal diphtheria.

EFFECTS OBSERVED CLINICALLY.

So far I have dealt almost entirely with the statistical proofs of the value of antitoxin, though the clinical fact of its action in arresting the local affection has also been demonstrated. I must very briefly state the effects of antitoxin as exhibited in individual cases. A much larger experience has confirmed the opinions expressed in a paper read by Dr. Washbourn, Mr. Card, and myself before the Clinical Society in December, 1894,³⁰ when we recorded the results of the treatment in the first series of cases treated in this country. Not only is the extension of existing and the formation of fresh exudation stopped, but also that which is present clears off more rapidly than in cases where no antitoxin is used. When the nasal passages are affected, the foul discharge from the anterior nares that nearly always exists very quickly ceases, greatly to the comfort of the patient and those in attendance upon him. In many of the fatal cases the fauces and nose are absolutely clean before the termination of the case. The lessening of the faucial inflammation enables the patient to breathe and take nourishment without discomfort. The enlargement of the cervical glands disappears and the inflammation of the cellular tissue of the neck subsides. The visible improvement in the general condition of the patient reflects the changes that have taken place in the local lesions. Pulse-rate and temperature fall, appetite returns, and the patient is soon convalescent. Even in cases which end fatally, when antitoxin has been given at too late a period to save life, the patient's last moments are made easeful, and the death-bed is robbed of all the loathsomeness we were accustomed to associate with fatal diphtheria.

The list of drugs in our *Pharmacopœia* is long. For all it is claimed that they are of value in the treatment of one and the other of the ills to which our flesh is heir. Yet I do not think I exaggerate when I say that few of even the most useful have been subjected to the same fierce criticism as has the antitoxin of diphtheria, and still fewer have borne the ordeal as triumphantly.

REFERENCES.

- ¹ Sanné, *Traité de la Diphthérie*, 1877, p. 457. ² Met. Asyl. Board, Statistical Committee's Report for 1895, p. 132. ³ Henoch, *Lectures on Children's Diseases*, New Sydenham Society's Translation, p. 308. ⁴ Sanné, *loc. cit.*, p. 468, 469. ⁵ Quoted by Welch in paper entitled the Treatment of Diphtheria by Antitoxin, *Trans. Assoc. Amer. Physicians*, vol. x, 1895. ⁶ Report of Antitoxin Committee of Clinical Society, 1898, p. 12. These cases were treated at University College, Guy's, St. Thomas's, Middlesex, London, and St. Bartholomew's Hospitals for ten years preceding the antitoxin treatment. ⁷ Met. Asyl. Board Statistical Committee's Reports for years mentioned. The cases are reported in some of the reports of the medical superintendents of the Fever Hospitals. ⁸ Ditto, Report for 1895, p. 131. ⁹ BRITISH MEDICAL JOURNAL, October 23rd, 1897. ¹⁰ *Gaz. des Hôp.*, 1864, p. 99. Quoted by Sanné, *loc. supra cit.*, p. 469. ¹¹ *Transactions of Eighth International Congress of Hygiene*, vol. ii, p. 192. ¹² Cadet de Gassicourt, *Traité Clinique des Maladies de l'Enfance*, 2nd ed., vol. iii, p. 250. ¹³ Roux, *Trans. Cong. Hygiène*, vol. ii, p. 192. ¹⁴ Welch, *loc. cit.* ¹⁵ Henoch, *loc. cit.*, p. 309. ¹⁶ Trousseau, *Clin. Med.*, Syd. Soc. Trans., vol. ii, p. 595. ¹⁷ Amer. Pediatric Society's Report on Antitoxin, See BRIT. MED. JOURN., May 29th, 1897. ¹⁸ Reports on Antitoxin Treatment, Stat. Com. Rep., 1895-97. ¹⁹ Belin, *Minch. med. Woch.*, October, 10th, 1897; and EPITOME, BRITISH MEDICAL JOURNAL, December, 11th, 1897. ²⁰ Roux, *loc. supra cit.* ²¹ Zuppinger, *Lancet*, April 3rd, 1897. ²² Sidney Martin and Bertram Hunt, BRITISH MEDICAL JOURNAL, September 3rd, 1898. ²³ Cuno, *Deut. med. Woch.*, December 24th, 1896, and BRITISH MEDICAL JOURNAL EPITOME, January 23rd, 1897. ²⁴ See a paper by Sims Woodhead on Post-Diphtherial Paralysis, BRITISH MEDICAL JOURNAL, September 3rd, 1898. ²⁵ BRITISH MEDICAL JOURNAL, September 8th, 1898, p. 595. ²⁶ MacCombie, M.A.B. Stat. Com. Rep., 1897, p. 63. ²⁷ Hume, M.A.B. Stat. Com. Rep., 1895, p. 78. ²⁸ Reports on Post-scarlatinal Diphtheria, M.A.B. Stat. Com. Rep., 1896 and 1897. ²⁹ See a paper on this subject in the Epidemiological Society's *Transactions*, vol. xv, p. 68. ³⁰ Washbourn, Goodall, and Card, Clinical Society's *Transactions*, vol. xxviii.

INDIAN EXCHANGE COMPENSATION.—It has been ruled that medical officers appointed by the Secretary of State for plague duty are not entitled to draw exchange compensation allowance.

Dr. N. H. CHOKSEY, medical officer in charge of the Arthur Road Hospital, Bombay, has by the order of King Humbert been invested with the Order of the Crown of Italy in recognition of services rendered to Italian savants during the prevalence of plague in 1897.

PERIPHERAL NEURITIS FOLLOWING THE SOFT SORE (? SYPHILIS).

By C. B. MAITLAND, M.R.C.P.E., Etc.,
Major, Indian Medical Service, Poona.

I HAVE lately read in the BRITISH MEDICAL JOURNAL a statement to the effect that syphilis can give rise to the so-called tertiary symptoms years after the initial sore, without there having been any intermediate symptoms whatever. More recently I read in the JOURNAL a question by Dr. Buzzard in speech at the Annual Meeting, asking whether it is right to assert that the bacterium of the soft sore can have no part in producing fine changes in sensory neurons, and whether such effect must be due to syphilis. The idea embodied in Dr. Buzzard's question struck me first in 1888, when, during a campaign, I attended a young English officer for a phagedænic venereal sore on the penis. There was no history of previous syphilis, and the sore recovered under local treatment only, but in three months he was a helpless cripple from multiple peripheral neuritis. I have a vivid recollection of how the poor fellow had to be lifted on to his orderly's back, and how his feet flapped helplessly about with the swing of the man's walk. No symptoms of syphilis ever appeared, and in time he recovered completely without treatment. When I say completely, I mean sufficiently to be still in the army. Whether his strength is exactly as it was before, or his knee-jerks returned, I am unable to say.

In 1895 I admitted eight young native cavalrymen into hospital for what was then called "ulcer of the penis." Whether these sores were due to one woman I cannot say, but they all were followed by "rheumatic" pains and weakness of limbs, especially of legs, that kept the patients in hospital many weeks after the sores were healed. None of these men showed any signs whatever of syphilis, or had any specific history. Four were treated with full doses of mercury and potassium iodide without any apparent benefit. They all recovered in course of time.

On December 16th, 1896, a native cavalryman had connection, and came to me on March 25th, 1897, with a large collar sore on the inside of the prepuce, painful and slightly phagedænic. There was no history of syphilis, so the only mercury he took was a 5-gr. blue pill daily for thirteen days. The sore healed on April 25th. By May 11th multiple peripheral neuritis was well marked. Mercury and iodide were then pushed, but seemed to do no good. Strychnine pleased both the patient and myself most.

On August 11th, 1897, he was able to be sent on sick leave, and returned on December 8th, as he said, quite well; but the right hand, which had all along been more affected than the left, was still a little weak, and the knee-jerks were absent. He did duty till June, 1898, when he was admitted with a large venereal sore and suppurating bubo. These soon healed, but he has remained in hospital with what I can only call general paresis, the sole symptoms being increasing weakness of the limbs without loss of size or, as far as I can judge with the machine at my disposal, of electrical excitability of muscles, and some pain in joints of limbs. His grasp is like that of a small child, and his legs are just strong enough to let him creep along with the help of a stick. There are no sensory troubles except the pain mentioned. The man is not shamming, as he only wants a year to complete his pension service. No treatment has done any good.

Another native cavalryman acquired a venereal sore about the middle of January, 1897, and came for treatment on March 30th, 1897, with the whole foreskin in a phimotic and ulcerating condition. This was removed and a sore found on the glans. Amongst other treatment he had a 5-grain blue pill daily for twenty-eight days. There were no symptoms of syphilis. By May he was a good example of a severe case of multiple peripheral neuritis. He was treated chiefly with strychnine, and on September 26th was able to go on sick leave. He returned to duty on March 25th, 1898; the knee-jerks were still absent, and slight weakness of the right leg remained. This latter symptom has by now quite gone.

Of these eleven cases I saw nine from the first appearance of the sores and two after the sores had been in existence three months. None of the men had any previous history of