appeared. The threadworms persisted in spite of all treatment for two years.

It seems probable that in this case, as in the preceding, the symptoms may have been due to the presence of threadworms in the appendix, which would also account for the obstinacy of the case, if the view I have suggested be correct. That pain in the right iliac fossa was due to threadworms in the appendix has seemed possible in other cases, but it can only rarely happen that actual evidence can be obtained as in the first case mentioned. There was, however, ample evidence in the series of 200 cases examined *post mortem* that the presence of threadworms may cause swelling and thickening of the vermiform appendix, especially when the worms are present there in large numbers. Such a condition was found several times, and, indeed, it was sometimes possible to predict the presence of threadworms in the appendix before opening it from its swollen appearance. The following cases may serve as typical instances:

W. M., aged 2½ years; empyema, suppurative pericarditis, and commencing peritonitis. The vermiform appendix had a markedly swollen and thickened appearance: the rest of the intestine appeared quite normal. The appendix was found to contain 20 threadworms, nearly all of these being in the distal inch; of this number, only 5 were full-grown females containing ova; the rest were very minute, measuring from about 3 mm. upwards; many of them were females, but some were males. Only 4 threadworms, all full-grown, were found in the large intestine; no young ones were seen with the naked eye, nor in portion of the mucus examined with the microscope. One full-grown female was found 4 inches above the ileo-cæcal valve.

J. L., aged 3½ years; tuberculous meningitis. The vermiform appendix had the typical swollen appearance suggesting at once the presence of threadworms; it was therefore examined carefully. In the distal ½ inch, close to the tip, 26 were found; in the rest of the appendix 33, or 43 including 10 at the mouth of the appendix—that is, 69 in all. Most of those in the appendix were very young females. The mucus of the lower part of the ileum near the ileo-cæcal valve was examined microscopically, but showed no ova and no embryos. There were numerous threadworms in the colon and rectum.

A. A., aged 4 years; spinal caries, tuberculosis. The vermiform appendix was much swollen—"nearly as thick as one's little finger." The appearance suggested the presence of threadworms. The mucus from the appendix was examined with a low power of the microscope, and showed 43 threadworms, chiefly very minute young females. There were many threadworms in the cæcum but only a few in the rest of the colon.

The above observations suggest some practical points with regard to treatment. The idea that the rectum is the chief habitat of the threadworm is erroneous and misleading; it is here that the worm, perhaps, most often causes symptoms—namely, itching and worry, but it is not here that the worm is most commonly found. It is most often present in the cæcum, and, as I have shown, in two-thirds of the cases it is present also in the vermiform appendix; moreover, in some cases it is found in the small intestine.

Any injection therefore which is only large enough to reach the top of the rectum or the lower part of the colon is not likely to do much good. Some have even gone so far as to say that injections are useless, on the ground that it is not possible to reach the cæcum by this means. Such a statement is no doubt perfectly true if the injection consist of only 3 to 4 ounces of fluid. With such a quantity no part higher than the descending colon is likely to be reached. But with larger quantities—12 to 18 ounces or more—it is quite possible to fill not only the cæcum, as is apparent in the treatment of intussusception by injection, but also the vermiform appendix, if one may judge from *post-mortem* experiments in which I have repeatedly demonstrated this. Injections, therefore, are of the greatest importance, but it is essential that large quantities of the fluid should be used—a child of 6 to 12 years will often tolerate as much as 16 to 20 ounces—and experience has shown that large injections of salt and water or of infusion of quassia are often effective.

Treatment by injection alone, however, often fails, and this is easily intelligible, for the large intestine varies so much in length, and therefore in capacity, in children of about the same age, that it is practically impossible to know what amount of fluid is required in any given case. Thus, in one child, aged 15 months, the large intestine was 28 inches long, in another, aged 15 months, it was 34 inches, while in another, aged 23 months, it was 44½ inches long, and similar differences were found in older children.

Moreover, as the above observations have shown, threadworms may be present, though not very commonly, in the small intestine; and the chance of reaching these by any injection is very slight, even when they are situated, as is usually the case, only just above the ileo-cæcal valve. In the

post-mortem experiments alluded to it was found that occasionally the fluid did pass up into the small intestine, but it is obvious that the conditions after death are so different from those during life that any conclusions drawn therefrom must be of very doubtful value. There is, however, some evidence that this occasionally happens during life, and in one case where an enema of olive oil followed by a large enema of soap and water, was given as an aperient to a boy, aged 4½ years, twenty-four hours before death, I found the oil as high up as 6 inches above the ileo-cæcal valve. This, however, is probably quite exceptional, and for practical purposes one may say that the small intestine cannot be reached by injections.

It is necessary, therefore, to give drugs by the mouth also, and no drug has proved more effectual in my experience than *santonin*. At the same time the prevention of any possible conveyance of ova to the mouth must be attempted, for, as shown above, this is probably one mode of keeping up the supply of worms, and the cutting short of finger nails, and the prevention—by mechanical restraint if necessary—of scratching the anus, are no unimportant adjuncts to treatment.

The conclusions to which the above observations would seem to point, are:

1. That the appendix vermiformis is a common habitat of *oxyuris vermicularis* in childhood.
2. That the generally accepted view that every single ovum of *oxyuris vermicularis* must be swallowed before it can be hatched, is at least open to doubt, and there is a strong probability that the appendix vermiformis serves in some cases as a breeding place for threadworms.
3. That the presence of threadworms in the appendix may cause a catarrhal condition therein, as shown *post mortem* by a swollen appearance, due to thickening of its wall.
4. That this swollen condition of the appendix due to threadworms is associated in some cases with pain in the right iliac fossa which may simulate ordinary appendicitis.
5. That in the treatment of threadworms large injections must be used, and in view of the difficulty of dislodging the worms from the appendix, and their possible presence in the small intestine, the injections should be combined with the administration of drugs by the mouth.

REFERENCE.

¹ *Zoologie Médicale* Paris, 1889, i, 713.

NEUROTIC ULCERS OF THE MOUTH

(STOMATITIS NEUROTICA CHRONICA).

By W. KNOWSLEY SIBLEY, M.A., M.D., M.R.C.P.,
Assistant Physician, North-West London Hospital.

IN 1894, Jacobi of New York published in the *Transactions of the Association of American Physicians* a paper on the above subject, in which he described three cases that had come under his notice. He collected all the cases he could find in literature, namely those reported by Kaposi, Kopp, Flatau, and Mandelstamm, and came to the conclusion "that such cases must be very rare."

The cases I am about to describe I have met with during the last few years—two in private, and one in hospital practice, all being women. The patients were rather more of the hysteroid or neurotic than the insane type, as those described by Jacobi, two of whose cases were men. I am also inclined to believe that the disease is much commoner than he supposes, but that it is frequently overlooked. Such cases are often classed as simple catarrhal or dyspeptic ulcers, whereas I consider they are the direct result of a tropho-neurosis, of which dyspepsia in some form or other may be, and often is, another expression.

CASE I.—The first case is that of a widow, aged 74, who has had good general health. There is no known mental disease in the family. She had ten children before she was 35, and several miscarriages. Three children died from diphtheria, a fourth (13) from tuberculous peritonitis, and another at 53 from carcinoma of the kidney. The other children are living and well. She has lived an extremely active life.

The first ulcer came on the gums in 1875 during great domestic trouble, and it lasted a few weeks. During a visit the country the ulcer healed up. Six months later she came back to London, and the ulcers soon returned. At this time she had a perfect set of teeth. The ulcers usually came between the teeth, and there was great difficulty in applying local applications to them.

During 1880 there were numerous ulcers on the tongue, gums, and buccal mucous membrane.

During 1882 she had all her teeth extracted, but the ulcers continued as severe and as distressing as before.

In 1883 the patient had an attack of eczema, after which she went to the Pyrenees, but the ulcers continued.

In 1884 the mouth remained in the same state of ulceration, and during the months of September and October she was yachting a good deal, but without any improvement.

During the whole of 1885 she had ulcers.

In 1886 she was free from ulcers from February to the middle of March; after this the disease was very severe, and continued so during the next four years. She was practically never free from ulceration.

In 1890, during October, November, and December, during a visit to Scotland, the mouth was almost well.

In 1891 the ulcers continued and caused great pain, though the patient was yachting a good deal. She was at Malta for three months, but continued in great misery till December, 1893.

In January, 1894, she was free for a time while in London. During February, March, and April the ulcers were moderate, but after this she suffered tortures while yachting and while at Braemar from May till November.

In October, 1894, she broke her leg while in Scotland, and during November and December she was free from ulcers. After this she went to Cannes and Grasse, and was away for eight months, during this time the mouth improved considerably.

In 1895, the ulcers were severe, though they were less painful during the winter months.

In March, 1896, an improvement again took place.

In 1897 ulcers were present during April, May, and September.

In 1898 the patient had bad ulcers during January, May, July, and September.

Thus it is seen that often for many weeks the patient's existence was made almost unbearable from ulcers which gave rise to the most excruciating pain; the tongue and often the lips were much inflamed and swollen, so that any movement either for speech or mastication was almost intolerable. Though naturally very active, she was often compelled for days together to shut herself up and see no one. The mucous membrane of the mouth and lips was very glossy, thin, and of a bluish tint, and covered with cicatrices from old ulcers. The tongue was usually covered with a thick brown fur. There were no distinct dyspeptic symptoms.

CASE II.—The second case is that of a single woman, aged 50, whose father died from the exhaustion of neuralgia, following herpes zoster: her mother died from carcinoma of the rectum. One sister died, at 45, from carcinoma, and another about the same age from tuberculous disease. The patient had pleurisy and bronchitis when eight years old, and has always been thin, pale, and rather delicate. She has a faint systolic mitral murmur, but there is no enlargement of the heart or anything unusual in its action. She suffers from constipation and dyspeptic symptoms. The urine is of low specific gravity, but otherwise normal. She suffers from headaches, hot and cold sensations, perspirations, flushings, and various pains. Recently the patient had an attack of erythema multiforme. She has had several attacks of hystero-mania (climacteric), lasting usually only a few days, occasionally two or three weeks, and gets perfectly well again. At the present time she presents all the signs of climacteric hysteria, and is very easily upset.

She has suffered from ulcers of the mouth at intervals almost as long as she can remember. She formerly suffered a great deal from conjunctivitis, and since this has been better the mouth has been worse. At times she is very irritable, and this is always accompanied or followed by a bad mouth. The attacks usually coincide with domestic worries. She develops a series of usually small but very painful greyish ulcers; sometimes only one, but more frequently two, three, or more at a time, or immediately following one another, on the sides of the tongue, or frequently beneath the tongue, either at the tip or sides; occasionally on the buccal mucous membrane, pharynx, palate, or mucosa of the lips; usually accompanied by more or less stomatitis, with general swelling of the organ, which is considerably coated with a thick pale fur, and associated with salivation and offensive breath, and slight enlargement of the submaxillary glands. This condition lasts from a few days to two or three weeks, and then for a few days or even weeks she is free from discomfort. A few years ago she generally had an outbreak of ulcers about the catamenial periods, and when these ceased or became irregular, hysterical symptoms accompanied by ulcers occurred at these times. On one occasion she lost the use of one arm for some hours.

When the mouth is very bad and the attack is a severe one, all movements of the tongue, either for speech, mastication, or deglutition, are extremely painful, and the patient frequently for days goes without taking proper nourishment.

It not infrequently happens that when the mouth is ulcerated there is also some irritation of the conjunctiva, usually on one side, and very occasionally there has been slight superficial ulceration of the external genitalia, apparently commencing as a folliculitis.

CASE III.—The third case is that of a widow, aged 64, whose father died at 70, and mother at 24 from consumption. The catamenia ceased about 53. It is stated that she had inflammation of the kidneys five years ago, and was in bed three weeks. She has had thirteen children, and a great deal of trouble during the last few years, and has had to work hard with the needle for her living. Three years ago (1896) she felt a small pimple at the tip of her tongue, which soon became an ulcer. She has had sores on the tip and sides of her tongue and gum at intervals ever since. The whole tongue is evenly coated with a thickish, grey fur. Three years ago she had several teeth out, but the ulcers were not improved. The gums and buccal mucosa are smooth, and rather glossy in appearance. She frequently complains of a burning sensation in different parts of the tongue, even when no obvious ulceration is present.

PATHOLOGY.

This disease has usually been described as of the nature of pemphigus, although its distribution is sometimes sug-

gestive of herpes. It, however, differs from pemphigus in its persistence through many years. Pemphigus may recur several times at intervals of a few weeks or months, but during the intervals it ceases altogether. With the onset of pemphigus there is usually a rise of temperature, but I have not observed this in my cases. An actual vesicle or bleb is not by any means common in this disease, but it generally commences as a crack or streak, or from the beginning as a small superficial bright red ulcer. Occasionally in the tongue it begins in an inflammatory localised thickening just beneath the mucous membrane, which rapidly breaks down and forms an ulcer, usually with a slough in the centre and considerable inflammatory redness around. It sometimes happens that the ulceration is preceded for a day or two by a heaping up of the epithelium, often forming a pale, gelatinous-looking ridge fitting in the spaces between the teeth. At other times the ulcers are preceded by small gelatinous-looking bodies about the size of millet seeds, and occasionally by small vesicles; accompanying the ulcers is usually a considerable desquamation of the surface of the tongue. There is usually a good deal of burning sensation and great distress, accompanied by profuse salivation, and, if the ulcer is very indolent, with oedema of the parts around. If the lesion is situated in the mucous membrane of the lips, these may become so swollen as to hardly permit of the mouth being opened and the tongue protruded. I consider these ulcers are produced by a distinct trophoneurosis, and that they are quite different from the common catarrhal or dyspeptic ulcer.

ETIOLOGY.

Mental trouble appears to be the cause of the ulcers in these subjects.

The first case was a woman who managed a large household, and maintained a very prominent position in society on a very limited income, which was an incessant cause of great anxiety to her. The ulceration commenced at a time of unusual mental strain, and continued through many years; in fact, during the whole period that the anxiety continued. Of recent years her position has been rather less embarrassing, and concurrently the ulceration has been less severe.

In the second case the patient's attacks almost always correspond with some more or less trivial worry, but one which affects her out of all proportion to its importance.

In the third case, trouble connected with money matters appears to have been the cause of the complaint.

TREATMENT.

No medicinal treatment appears to give any very obvious result; possibly some nerve tonics may do good indirectly, and local applications may relieve the distress and pain for the time, and assist in the more rapid healing of the sores. Better results have been obtained from painting the ulcers with tincture of iodine than the application of nitrate of silver. Many cases accompanied by severe pain are greatly relieved by cocaine and borax, carbolic or peroxide of hydrogen mouth washes. I believe in giving opium, especially at night, in moderate doses.

There is no doubt that the only curative treatment is complete removal from home and its accompanying worries, and it appears that mountain air is preferable. No good has been derived from sea voyages; in fact, the condition during them is sometimes worse.

Removal of the teeth has no beneficial effect except that the seat of the ulceration is more easily reached by local application.

LITERATURE.

Boer, *Archiv f. Dermatol. u. Syph.*, 1890. Kopp, *Trophoneurosis Hysterica (Trophoneurosen der Haut)*, *Inaug. Diss.*, Würzburg, 1891. Flatau, T. S., *Chronic Recurrent Herpes of the Oral Cavity*, *Deut. med. Woch.*, 1891. Mandelstamm, B., *Pemphigus of Buccal Mucous Membrane*, *Berl. klin. Woch.*, 1891. Landgraf, *Berl. klin. Woch.*, 1891. Chiari, *Wien. klin. Woch.*, 1893. Jacobi, *Transactions of the Association of American Physicians*, vol. ix., 1894. Rosenbach, O., *Vesiculating Affections of the Oral Mucous Membrane*, *Deut. med. Woch.*, 1894.

THE Council of the Medical Faculty of Bucharest has expressed its approval of a scheme for the establishment of a new institute of bacteriology and experimental medicine in that city. It is proposed to place Dr. Cantacuzino, at present assistant in the Institut Pasteur of Paris, at the head of the new institute.