THREE CASES OF CYST OF THE PANCREAS TREATED BY INCISION AND DRAINAGE THROUGH THE ANTERIOR ABDOMINAL WALL.

By BILTON POLLARD, B.S., F.R.C.S.,

Surgeon to University College Hospital, and Consulting Surgeon to the North-Eastern Hospital for Children.

THE diagnosis of pancreatic cysts by means of the clinical signs presented by them is admitted to be very uncertain, and, as the condition is rather uncommon, it is desirable that cases should be reported; and there is the further reason for this course that surgeons have not yet decided which is the best way of dealing with them. Pancreatic cysts have been completely removed by some surgeons; by others they have been incised and drained through an incision in the anterior abdominal wall; and by others they have been drained through the loin by an incision below the last rib. The cases recorded below were all treated by drainage through an anterior abdominal wound. It has been objected to this method, on the one hand, that the adhesion which must of necessity be formed between the anterior and posterior walls of the abdomen might be injurious, and, on the other hand, that possibly simple drainage might not effect a permanent cure of the disease. I have delayed publication of these cases in order that, in regard to them, time might show the validity or otherwise of these two objections. It is now nearly five years since my first case was operated upon; the second case was operated upon three years ago, and the third case two years ago.

CASE I.

J. N., male, aged 34, a signalman, was admitted to University College Hospital, under the care of Dr. Ringer, on March 10th, 1894.

History.—According to his account, about three weeks previous to his admission he was awakened one night by a sharp pain in his epigastrium, and he began to vomit. After two days the vomiting ceased, and the severe pain was replaced by a burning sensation in the epigastrium, whence it radiated to his throat. This pain, associated with occasional attacks of vomiting, had persisted since. During the last six years the patient had suffered from some half-dozen similar, though less severe, attacks.

Condition on Admission.—The patient, though not emaciated, had evidently lost flesh. His temperature ranged from 99.6° to 101.2°. He took food well, and suffered no discomfort from it. There was no jaundice. The motions were of good colour, and contained no excess of undigested fat. The urine was normal. The upper part of the abdomen and the lower part of the chest on the left side were bulged, owing to the presence of a mass which could be felt to occupy the whole of the left hypochondriac region and a large portion of the epigastro-umbilical and left lumbar regions. The lower and right borders of this swelling could be defined with ease, forming a well-marked curved edge, which extended to the right for 2 inches beyond the middle line and downwards to a point 2 inches below the level of the umbilicus; above and to the left the mass extended beneath the costal margin; it occupied the lumbar region, but it did not fill up the loin in the interval between the last rib and the iliac crest. Fluctuation and thrill were easily obtained. The mass was only slightly movable. It could not be moved on the deeper parts, but the relation of its lower border to the skin shifted about half an inch during respiration. A ridge which was thought to be formed by the transverse colon could be felt crossing the lower part of the swelling obliquely. The percussion note over the whole area of bulging was deficient, and where the prominence was most marked the note was quite dull. In the upward direction the dullness was continuous with that of the liver and heart: in the left axilla the percussion note was impaired as high as the sixth rib.

First Step of Operation.—On the second day after admission Mr. Pollard operated. The abdomen was opened by an incision 6 inches in length in the left linea semilunaris. The stomach, the great omentum, and the transverse colon presented in the wound lying in front of the tumour, by which they were pushed forwards. The hand was passed into the abdomen, and the liver, kidneys, and spleen explored and found to be unconnected with the tumour. It was determined to open the cyst in the interval between the stomach and the transverse colon, and to do the operation in two stages. Some vessels were ligatured, and then the great omentum was divided transversely between the stomach and the transverse colon, and the small bag of the peritoneum opened. The tumour was still covered by a layer of peritoneum. A strip of gauze was packed into the wound between the stomach and the transverse colon, and the abdominal wound above and below this spot was closed. A dressing of cyanide gauze and salicylic wool was applied.

Second Step.—After the operation the temperature rose gradually till the fifth day, when it reached 102°. On the sixth day the gauze plug was removed, and the cyst was tapped with a hydrocele trocar, and about 1,400 c.cm. of fluid were removed.

Characters of the Contents of the Cyst.—The fluid was turbid, of a dirty drab colour, alkaline reaction, and of a specific gravity of 1.010. Microscopical examination showed the presence of leucocytes, some large granular cells, and some fat globules. A portion of the fluid was acidulated and boiled, with the result that an abundant precipitate of coagulated proteid was deposited. The filtrate gave the biuret reaction for peptones, and contained a small quantity of a copper-reducing substance. The action of the fluid on starch fibrin and fats was tested. It was found to

possess active amylolytic properties; in its presence fats were emulsified and fatty acids liberated; fibrin also was digested.

Third Step.—On the eighth day an anæsthetic was given, and after the removal of about 1,000 c.cm. of fluid by tapping the cyst was freely opened. Its wall was composed of dense fibrous tissue about $\frac{1}{8}$ inch in thickness. Some bleeding followed the incision into the cyst, and so the wound was packed with cyanide gauze. On the following day the gauze was removed; 500 c.cm. of dark-red, blood-stained fluid escaped, and a large drainage-tube was inserted. The temperature, which had been falling after the first tapping, rose again after the incision into the sac, but as soon as free drainage was established it fell rapidly, and subsequently remained normal.

After-History.—The man's general condition improved greatly after the evacuation of the cyst, and the farther history of the case is one of gradual recovery, uninterrupted except for a transient attack of jaundice some three weeks after the cyst was opened. This lasted two days only and was slight in degree. The discharge, which had been gradually lessening, increased in quantity with the onset of jaundice, and became tinged with green; it gave the reaction of bile-pigment with nitric acid. With the subsidence of the jaundice the discharge again diminished in amount, and changed to a clear, thin, serous fluid. Six weeks after the operation the man left the hospital; he was in very good health, but he was still wearing a tube 4 inches long.

In July, 1895, the patient was well-nourished and in excellent health, and had been at his work as signalman at Finsbury Park Railway Station for more than a year. He was still wearing a short tube, through which a small amount of clear discharge escaped. The tube was removed and in less than a week the sinus closed permanently. This patient was last seen in December, 1898. He said that since the operation he had had occasional attacks of burning pain at the epigastrium. The last attack was a year and nine months ago; since then he had enjoyed excellent health and had felt much better than he had done for several years previous to his operation. Nothing could be felt in the regions previously occupied by the cyst, and the patient had no discomfort there; the scar had yielded, and the patient was wearing a belt.

CASE II.

J. W., male, aged 56, was admitted to University College Hospital, under the care of Dr. Ringer, on January 17th, 1896, complaining of a tumour in his abdomen.

History.—On December 18th, 1895, he began to feel unwell. On the 22nd his abdomen felt distended, and he had slight abdominal pain, and felt short of breath. During the following night he awoke with severe abdominal pain, was repeatedly sick, and felt very short of breath. The attacks of pain and vomiting recurred, but were milder in character during the following days. On January 3rd a tumour was detected in the abdomen. During the illness there had been nothing abnormal in the characters of the urine, and the patient had not been jaundiced. The patient could not assign any cause for his illness. He had not had any injury to his abdomen. He had been losing flesh for two months. He had had frequent attacks of epistaxis, and four years ago "he broke a vessel" in the left side of his scrotum.

Condition on Admission.—The abdomen was greatly enlarged, and particularly so at the upper part. There was marked bulging in the epigastric and left hypochondriac regions. A large tumour could be felt in the abdomen filling the left upper quadrant of the cavity, and extending to the right nearly to the nipple line and downwards to just below the umbilicus. It extended upwards beneath the left costal margin, and gave rise to dullness on percussion in the left hypochondriac region, which reached to the level of the sixth intercostal space midway between the nipple and mid-axillary lines. Overlapping the right upper portion of the tumour in the epigastrium, and separating the dull area due to the liver from that due to the tumour, there was an area of resonance, the extent of which became considerably increased when the stomach was distended with gas. The surface of the tumour was smooth, and it had a tense elastic feel. There was well-marked pulsation in the tumour, but this was only communicated, for it disappeared when the patient rested on his hands and knees or turned on to his left side. The tumour moved very little if at all with respiration. It was slightly movable from side to side. It could be felt readily from the loin, but it did not fill the loin, and did not cause any bulging there. On percussing the tumour lightly an absolutely dull note was obtained, but on deep percussion there was some resonance in the note. A mass, which appeared to be the lower end of the right kidney, could be felt behind the tumour just below the last rib. When air was injected up the rectum a well-marked tympanic note was obtainable in the left flank and at the upper part of the abdomen. This resonant area was behind the tumour. The urine occasionally contained a slight trace of albumen, but in other respects it was normal. The stools had a natural appearance. The patient was emphysematous, and expectorated slightly blood-stained mucus.

Operation. The patient was kept under observation for fourteen days before operation. During this time the tumour became tenser, but in other respects it did not change. The patient lost 5 pounds in weight. On the fourteenth day after admission Mr. Pollard operated. The abdomen was opened by a longitudinal incision, 4 inches in length, placed a little to the left of the middle line, and over the most prominent part of the tumour. The great omentum was exposed; it was adherent to the parietal peritoneum below and to the left, but above and to the right there were no adhesions, and there the stomach and the lower border of the liver were displayed. The great omentum was torn through and the cyst wall was exposed. The cyst was punctured with a hollow needle, and a brownish fluid withdrawn. After sufficient fluid had been withdrawn to render the cyst walls lax, an incision, about one inch long, was made into the cyst, and its contents were thoroughly removed; 4,600 c.cm. of fluid were withdrawn. The abdominal wound was sutured at each end, and the margins of the incision into the cyst were secured by several stitches to the margins of the central portion of the abdominal wound. The cyst occupied the smaller bag of the peritoneum, and from its interior the left kidney could be felt of normal size behind it.

Characters of the Contents of the Cyst.—The fluid removed from the cyst had a greenish-brown colour; its specific gravity was 1.012; its reaction was alkaline; it contained half albumen. Its digestive properties were

investigated by Dr. Vaughan Harley, who reported that it contained the three ferments—trypsin, steapsin, and amylpsin, of pancreatic juice. Some fluid which was withdrawn from the cyst on the fourth day after the operation had no digestive properties.

After-History.—The patient made an uninterrupted recovery. The highest temperature recorded was 100° F. on the day after the operation; with that exception it was normal. There was very little discharge, and the cavity contracted up rapidly. The patient got up on the sixteenth day after the operation, and he was discharged on the twenty-fifth. There was a tube 3 inches in length in the sinus, and the patient was told to keep it in as long as he could. He however removed it as soon as he left the hospital, and the wound healed in a few days. This patient was seen in December, 1898. He was quite well, and had not had any abdominal trouble since his operation. A small ventral hernia had protruded through an aperture in the centre of the scar at the place where the drainage tube had been. No recurrence of the cyst could be detected.

CASE III.

T.A.B., male, aged 36, was admitted to University College Hospital on January 30th, 1897, under the care of Dr. Roberts.

History.—Towards the end of October, 1896, he began to suffer from pain in his back. On November 11th he was seized with a sudden pain in his left hypochondrium, which spread up into his chest, and, as he says, "nearly choked him." He kept his bed for a month, during which time the severe pain subsided, but the pain in his back continued until about a fortnight previously. About two months before his admission a tumour was noticed in his abdomen.

Condition on Admission.—The patient was well-nourished and looked healthy. There was fullness in the epigastric and the left hypochondriac and lumbar regions. A large tumour could be felt in the left loin and extending thence upwards beneath the ribs, downwards to the level of the umbilicus and across the epigastric and upper part of the umbilical regions to 2 inches to the right of the middle line. Though occupying the left loin it did not really fill it and could not be made to bulge out the space between the last rib and the iliac crest. The lower end of the left kidney could be felt behind the tumour just below the last rib. The borders of the tumour, with the exception of the lower, were not well-defined. It moved a little with respiration, and on grasping the tumour with both hands, one in front and one in the loin, it could be moved a little from before back. It felt distinctly elastic. The percussion note was deficient in resonance, but not dull. The stomach resonance and the succussion splash could be detected in front of the tumour in the epigastric region, over the abdominal mass, but percussion over the chest elicited a definite dull note, extending in the nipple line to about the fifth rib. The urine contained a trace of albumen, but was otherwise normal. There was no excess of fat in the stools. Temperature normal.

Operation.—On February 27th, 1897, Mr. Pollard operated. The abdomen was opened by an incision 4 inches long, passing through the rectus about an inch to the left of the middle line. The hand was introduced, and the tumour explored. It was free from the kidney, and it was found to be presenting between the stomach and the transverse colon. The great omentum was torn through, and the wall of the cyst was exposed. A large aspirator needle was then plunged into the tumour, and over a quart of brownish fluid was withdrawn. The cyst was then freely opened and emptied. The margins of the incision were stitched to the skin, and the upper and lower parts of the abdominal wound were closed. A large tube was passed into the cyst, and a dressing of cyanide gauze and salicylic wool was applied.

Characters of the Contents of the Cyst.—The fluid removed had a dirty brown colour and a rather offensive smell, and it deposited a large sediment. It had a faintly alkaline reaction, and leucin drops and crystals were seen with the microscope. The fluid was examined by Dr. Vaughan Harley, and was found to possess all the properties of pancreatic juice.

After-History.—After the operation the temperature gradually rose, and reached 103° F. on the third day; it then fell, and reached the normal on the sixth day. The dressing was changed daily, and sometimes twice a day, as there was a copious, foul-smelling discharge. Many large black sloughs presented at the wound and were removed. The patient left the hospital six weeks after the operation. He had a small sinus then, but that healed in a fortnight. This patient was seen in December, 1898. He had had no further trouble from the cyst. The upper and lower parts of the abdominal wound were firmly united, but at the centre, where the drainage tube had been, there was a weak spot through which a hernia protruded. Nothing abnormal could be detected in the regions which had been previously occupied by the cyst. The patient's general health was good, and he had gained flesh.

Cysts of the pancreas have frequently developed after abdominal injuries, and it has been suggested that their formation is due to extravasation of blood, and subsequently of pancreatic secretion from the lacerated gland. There was no history of injury in any of my cases. In two of them the onset of the illness was marked by severe pain in the epigastrium and by frequent vomiting; in the other case the patient had a sudden severe pain, which he located in the left hypochondriac region, and which he stated spread up to his chest and nearly choked him. Severe epigastric pain has been frequently noted as an antecedent of pancreatic cysts. Two of the patients had lost a good deal of flesh, and one of them lost 5 lbs. in fourteen days. In all the cases the urine was free from sugar. In two there was an occasional trace of albumen. The stools appeared natural, and there was no excess of undigested fat. There was no jaundice before the operation in any of the cases. In one there was a slight attack after the operation.

All the three cases presented very similar clinical signs. In each there was a large tumour occupying the epigastric

and left hypochondriac regions, and portions of the umbilical and left lumbar regions. The tumours moved very little with respiration, but they could be moved slightly from before backwards and from side to side between the two hands. In two of the cases stomach resonance could be obtained by light percussion in the epigastric region, although the tumours appeared to fill that region, and the area of stomach resonance was increased by distending the stomach with gas. In one case the percussion note was dull over the most prominent part of the tumour in the epigastric region, although at the operation this part of the tumour was found to be covered by the stomach. In this case the transverse colon could be traced over the lower part of the tumour. In one case there was good resonance in the flank behind the tumour, and this was increased by injecting air up the rectum. In each case the tumour gave rise to a dull area in the left hypochondriac region, and it also occupied the lumbar region, but it could not be made to fill the loin in the way that a hydronephrotic kidney either does or can be made to do. In two of the cases the lower end of the left kidney could be felt behind the tumour just below the last rib. The fact that the tumours moved very little, if at all, with respiration, and the presence in two cases of the stomach and in one of the transverse colon in front of the tumour showed that the tumours were not situated in the liver, and suggested that they were located either in the small bag of the peritoneum or in the retroperitoneal tissue. In one case distinct fluctuation and thrill could be elicited, and in all three the tense and elastic feel of the swellings suggested that they were cystic. In the first case the tumour was diagnosed as a cyst which was not connected with either the liver or the kidney, and it was thought to be either a pancreatic cyst, or a collection of fluid in the lesser bag of the peritoneum. The diagnosis of pancreatic cyst was made in each of the other two cases on the grounds stated above and on their resemblance to the first case; and in all three the diagnosis was verified after operation by an examination of the fluid removed from the cysts. In each case the fluid removed at the operation contained the three digestive ferments of pancreatic juice—amylpsin, trypsin, and steapsin; but after they had been drained for a few days the discharge from the cysts was found to have lost its digestive powers over starch and fibrin; its fat-splitting properties were not tested.

Pancreatic cysts have come forward in at least three different situations, namely, above the stomach, opposite the small omentum; between the stomach and the transverse colon, opposite the great omentum; and below the transverse colon, opposite the descending layer of the transverse mesocolon. The second of these positions is the most common.

In the first of my cases the cyst lay behind the stomach and the transverse colon, but it had not disturbed their relations; in the third case the cyst presented between the stomach and colon, and separated them from one another; and in the second the cyst presented below the stomach, and had displaced the transverse colon downwards considerably, but as there were adhesions between the omentum and the abdominal wall below, it was thought unwise to disturb them sufficiently to determine the exact position of the colon. In all three cases the cyst was covered in front by omentum. In the first and third cases the great omentum and the ascending layer of the meso-colon had to be divided to reach the cyst wall; in the second case the tissues covering the cyst wall were so blended together that their component parts could not be identified.

The practice of aspirating abdominal tumours either for the purpose of diagnosis or treatment merits strong condemnation. In some cases undoubtedly it may be safely done, but it is not possible to determine which by physical signs. In the first of these cases if the most prominent part of the tumour, which was dull on percussion, had been aspirated the needle would have passed through the stomach. In the second case the most prominent part of the tumour had separated the stomach and colon so widely that aspiration could in all probability have been done without injury to the abdominal viscera. In the third case it would have been a lucky chance if the needle did not transfix either the stomach or the colon.

In the first case, which was operated upon nearly five years

years ago, the operation *à deux temps* was performed, but in the other two the cysts were opened at once, after the greater part of their contents had been removed by aspiration. The margins of the openings into the cysts were stitched up to the skin, and the cysts were drained for some time. It was quite easy to prevent leakage into the peritoneum when the cysts were opened at once, and on other grounds that method is preferable to that *à deux temps*.

The method employed for the treatment of these cases was the simplest possible, and I think it was the best. The removal of the cysts has been recommended, and the operation has been performed successfully in some cases, but in several there has been a fatal termination. In my cases extirpation of the cysts would have involved most difficult and severe operations. And since it is now clear that drainage alone will suffice for the cure of pancreatic cysts, I think it is no longer justifiable to attempt their removal. A point more worthy of consideration is the route by which drainage of the cysts should be effected. The advisability of draining these cysts through a posterior incision has been urged, in order to avoid the necessity of making an adhesion between the cyst and the anterior abdominal wall. In the cases above reported no harm has resulted from the adhesions. If the lumbar operation could be performed safely without making an anterior incision at all, I should be in favour of it; but in all cases I should think it would be advisable to make an anterior incision in order to verify the diagnosis; and possibly, too, in order to determine the spot at which the cyst could be best reached by the posterior incision. The posterior route would certainly be a good one in cases where the cyst is small, and in which it would be difficult to bring it up to the anterior abdominal wall. It must not be forgotten that important structures and large blood vessels may be encountered in attempting to reach the cyst through an incision below the last rib, and that owing to want of space it might be difficult to recognise and avoid them. The only objection to anterior drainage which has been apparent in my cases is the occurrence of hernia at the spot where the presence of the tube prevented primary union of the abdominal wound.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

OPERATION FOR ELEPHANTIASIS OF THE VULVA.

THE interesting cases reported in the BRITISH MEDICAL JOURNAL by Dr. Renner and Surgeon-Captain Smith of operation for elephantiasis of the vulva, remind me of one I treated when practising in Jamaica in the year 1865.

A black girl, aged 25, complained that she had for many years suffered from great enlargement of the external genitals. On examination, I found the parts greatly hypertrophied, and presenting all the well-known appearances of elephantiasis. The right labium formed a large tumour $19\frac{1}{2}$ inches in circumference, which, when the patient was standing, extended to within 2 inches of the knee. The left labium was the size of a large turnip. The nymphæ and prepuce of the clitoris were also hypertrophied but not to a great extent.

To avoid the risk of hæmorrhage in removing the diseased mass a local mechanic made for me a clamp composed of two parallel bars of lancewood united at one end by a hinge, and on the inside edge of each bar were placed several short sharp metallic pins, the object of these being to prevent the tissues slipping from the grasp of the clamp when they became slippery with blood.

The patient being under chloroform, the clamp was applied around the base of the right labium, as close to the pubes as possible, and tightly held. A slightly-curved incision on each side, about half an inch below the clamp, was made, and a few more strokes of the knife from above downwards separated the mass. Not more than 3 ounces of blood were lost, and this came entirely from the tumour, as, on wiping the cut surface controlled by the clamp with a sponge, it was completely white and bloodless. The pressure of the clamp was gradually relaxed, and one by one as they bled ten large vessels were tied. The cut edges were brought together by

several sutures. The redundant tissues of the nymphæ and prepuce were pared away with only a slight amount of bleeding. After a few days the left labium was removed in a similar manner with the aid of the clamp, and five vessels were tied. The patient recovered without a bad symptom.

After a considerable amount of fluid had drained away and the weight thereby lessened, the diseased mass on the right side weighed 3 lbs. 10 ozs., that on the left side nearly $\frac{3}{4}$ lb. The effect of the clamp in preventing hæmorrhage was very marked, and the sharp pins acted admirably in preventing the slipping which must have occurred had they not been used. As a curious circumstance it may be mentioned that, three years previous to my seeing her—and when, as she assured me, the parts were exactly in the same condition—she was delivered of a full-term child after an easy labour.

Eastbourne.

IZETT ANDERSON, M.D. Edin.

THE TREATMENT OF PNEUMONIA BY IRON PERCHLORIDE.

I CAN add my testimony to that of Dr. King in the BRITISH MEDICAL JOURNAL of November 12th, 1898, as to the great value of large doses of perchloride of iron in the treatment of pneumonia—a remedy far more effective here than in diphtheria or erysipelas. I have to thank a correspondent who some four years ago drew attention to it in the JOURNAL.

The late Sir Andrew Clark, in the discussion in London on the treatment of this disease, spoke very highly of it as appearing to have the effect of preventing consolidation. I have, however, of late combined large doses of digitalis, and consider the results eminently satisfactory in comparison with the usual treatment by expectorants. It is not merely an empirical, but a scientific, remedy. With children it is not so successful, because the administering of large doses of steel is so apt to cause vomiting. As to the exhibition of stimulants, I am cautious, believing that many a patient is poisoned by alcohol.

Ballarat, Victoria.

A. TEEVAN, M.R.C.S. Eng., etc.

A CASE OF OBSTINATE HICCUGH.

A PATIENT aged 50 years, of a highly nervous temperament, suffering from hæmorrhoids had them replaced under gas and ether. About two hours afterwards he commenced to hiccough. Pressure on the epigastric region checked it for a short time, but it recommenced and continued practically all night, sleeping or waking, the spasm being regular in character. Crushed ice was given by the mouth, without effect. Bromide of ammonia and soda, half drachm of each, with 20 grains of chloral, merely caused the hiccough to be slightly less violent during sleep. Mustard to the epigastrium was tried without any good result; neither had calomel (4 gr.) followed by a saline draught.

After consultation injections of morphine were given in quarter-grain, and later in half-grain, doses. These merely gave slight relief during the sleep which followed. Inhalations of ammonia were next tried, but with equal want of success. Laughing temporarily arrested it.

On the night of the ninth day I prescribed 2 minims of a 1 per cent. solution of nitroglycerine, with a drachm of the spirit of chloroform in half an ounce of water, to be given every hour. The first dose was taken at 9 p.m. The hiccough shortly began to diminish in frequency and violence, and at 2 a.m. finally ceased.

Whether this satisfactory condition would have resulted from natural causes it is difficult to say, but apparently the medicine last used produced the good effect, and as such cases are very troublesome, both to the patient and his medical attendant, I would suggest the above as a possible remedy.

R. W. S. CHRISTMAS, L.R.C.P., M.R.C.S.

Streatham, S.W.

THE new medical faculty established in the University of Odessa will be formally opened in August next.

MEDICAL GRADUATION IN RUSSIA.—During 1898 the degree of Doctor of Medicine was conferred by the University of Moscow on fourteen, by the University of Charkoff on five, and by the University of Kieff on three candidates. The number of students of medicine in the three universities severally was on January 1st, 1899, as follows: Moscow, 1,294; Kieff, 895; Charkoff, 709.