the empty sac from which a stone has been removed. There is no doubt that many of these sacs, especially when prolonged attacks of inflammation have destroyed their epithelial lining, tend to contract and cicatrise of their own accord if their orifice is of sufficient size to preclude the possibility of pus stagnating in their interior. If on the other hand the orifice of the sac is small and contracted and the sac itself of large size, some means must be adopted for its independent drainage, and the route which is selected must depend on the condition of the patient and the position of the sacculi. As most of these sacculi are situated near the base of the bladder, it will probably be advisable if they demand drainage to drain them through the perineum, and to make a temporary channel between the bladder and rectum if this is possible.

Table of Cases.

Number.	Patient.	Surgeon.	Prominent symptoms.	Result.
1	W. M., 55	Bruce Clarke	Stones removed on two occasions successfully. Died three years later	R.
2	E. H., 67	"	Stricture. Encysted stone found at post-mortem examination	D.
3	H. H., 57	"	Stricture; stones removed from cysts on several occasions	R.
4	—	"	{ All dependent on pieces of bone that	D.
5	—	"	{ had protruded into bladder and be-	R.
6	—	"	{ come nuclei for stones. One died }	R.
7	J. J., 62	Bruce Clarke	Frequent micturition; some pyuria	R.
8	J. B., 67	"	Pain; hæmaturia; foul urine	D.
9	J. P., 66	"	Cystitis; enlarged prostate	R.
10	J. P., 68	"	Pain and cystitis two years	R.
11	A. C., 64	"	Prostatic trouble several years	R.
12	J. G., 71	"	Symptoms fourteen years; lithotripsy on two previous occasions	D.
13	Gen. C., 60	"	Repeated slight attacks of cystitis. No operation	D.
14	Col. C., 80	"	Enlarged prostate seven years; catheter	R.
15	J. S., 69	"	Enlarged prostate four years; castration	D.
16	W. J. T., 57	Self, and Mr. Langton	Stricture six years. No operation	D.
17	R. W., 64	Self and Mr. Butlin	Severe hæmaturia	R.
18	C. S., 64	Bruce Clarke	Two years cystitis off and on	D.
19	J. P., 67	"	Three years cystitis. No operation	D.
20	J. E., 71	"	Stone crushed twenty-two years before death; cystitis one year. No operation	D.
21	F. P., 49	Mr. Langton and Mr. Cripps	Stricture; recurrent attacks of cystitis	R.
22	W. W., 60	Mr. Butlin	Recurrent attacks of cystitis	D.
23	W. B., 57	"	Stricture; cystitis some months and epithelioma of bladder. No operation	R.
24	G. E., 61	"	Cystitis. Left hospital	*
25	J. B., 40	Mr. Walsham	Lithotripsy three years previously; cystitis three months	R.
26	R. W., 38	"	Hæmaturia and cystitis for several months	R.
27	H. B., 14	Mr. Willett and Mr. Bowlby	Lithotripsy twice; lithotomy twice within eighteen months. Stone removed from retroprostatic pouch	R.

* The result of Case No. 24 is unknown.

The cases Nos. 7 to 27 are referred to in the present paper.

On looking over the symptoms which these cases presented it is difficult to single out any one sign which might be held to be of considerable diagnostic value. Cystitis of longer or shorter duration appears to have been present in every one of the cases, but as the great majority of them—namely, 16 out of 21—were above 60 years of age, and had in addition some evidence of enlargement of the prostate gland, the cystitis was naturally in the first instance ascribed to that condition, and not to the presence of a calculus. In several undue tenderness of the prostate gland when examined *per rectum* and rectal tenesmus were marked symptoms. Only 3 of the cases were under 40 years of age, one of them being but 14. The only two cases in the list between the ages of 50 and 60 had suffered from stricture; and it is noteworthy that 2 of my first 6 cases also had suffered from stricture, and were likewise between 50 and 60 years of age.

The difficulties of diagnosis were also enhanced by the fact that in several instances washing out the bladder was quite sufficient temporarily to restore the urine to a healthy condition. It is, perhaps, also worthy of note that in several instances the stones when extracted proved to be so thickly

coated with mucus, that even when they lay upon the table no audible note could be obtained when they were struck with the beak of the sound. This affords a ready explanation of the difficulty which may be experienced in detecting a stone in the bladder under certain conditions.

In 6 of the 21 cases no operation was performed in most instances, because the patient's condition was too bad to admit of any chance of alleviating his symptoms. Of the remaining 15 that were submitted to operation, 4 died and 11 recovered.

The fact that in less than half the cases was the existence of the stone known before the operation was undertaken, certainly affords an additional argument for exploration of the bladder in those cases of cystitis which do not yield to ordinary treatment.

In one instance (namely, Case xv) the patient had been castrated to relieve his prostatic symptoms, and the stone was only detected after death. It is quite possible that if his bladder had been explored at an earlier period he might have retained his testes and had his life prolonged after removal of the stone.

I have since that time made it a rule to explore the bladder in obscure cases of enlarged prostate which remain irritable after repeated washing, and have never had occasion to regret this plan of treatment, which I have no doubt largely accounts for this series of encysted calculi. Even if no stone is detected, the patient usually derives great benefit from the operation, owing to the rest which is afforded to the urethra.

REFERENCES.

¹ Latent Calculus in the Bladder. *St Bartholomew's Hospital Reports*, vol. xxvii, p. 117. ² See previous paper, *op. cit.*

ON ANTISTREPTOCOCCIC SERUM IN THE TREATMENT OF SMALL-POX.

By W. J. LINDSAY, B.A., M.B., B.C.

WHILE Resident Medical Officer at the Middlesbrough Isolation Hospital during the small-pox epidemic in that town and district during the earlier months of last year, I had opportunities of treating a large number of patients suffering from every variety of variola. It was soon evident that medicinal treatment did not influence the clinical course of the disease, and often did not alleviate symptoms. Attention was chiefly devoted to confluent cases, in which the prognosis is always doubtful, frequently most grave, since in mild cases no medicinal treatment was needed, and in hæmorrhagic none was of any avail. Drugs therefore giving unsatisfactory results, the clinical course run in these cases was closely observed in the hope of finding what, if any, form of treatment would be beneficial.

PRESENCE OF A PYOGENIC ELEMENT.

It was found that death most frequently occurred during a period of from three to five or seven days after the pustules began to dry and be absorbed. During this period the pustular "crusts" had disappeared from the face, the vesicles on the body had either ruptured or, the contents being absorbed in part, lay crinkled and dry on the skin as a sort of scab, the mouth and larynx, having lost much of their epithelium, exuded foul-smelling, blood-stained mucus. There was delirium, frequently violent, especially at night, universal coarse tremor, aphonia, often almost complete inability to move the limbs, great restlessness, refusal of food, the pulse, very rapid, weak, and generally failing, varied from 120 to 140 or more, the respiration was quick and shallow, varying from 30 to 40, or even 60 per minute. Death, as a rule, occurred apparently from sudden cardiac failure, which frequently gave no signs of its immediate approach. No organic disease of heart or lungs was found. Observing that this critical condition always occurred after the absorption of the pustules, it seemed reasonable to suppose that the specific virus of variola was reinforced by toxins of pyogenic organisms absorbed from the pustules, and that if this element of pyogenic septicæmia or pyæmia could be eliminated, the normal resistance of the tissues might, in certain cases, be able to cope with the effects of the variola virus successfully.

PROCESS OF PUSTULATION.

It was found that the purulent change occurred, first on

the surface of the vesicle, and not in its contents. Thus, as found by Klein,¹ and also by Copeman,² the clear serum of the vesicle yielded no growth on ordinary media (though it was in this serum the above observers found the bacillus which they believe to be specific for the disease) so the coagulation necrosis commences superficially, and at an early stage, if the thick cap of necrosed tissue of the vesicle be removed aseptically, the layer of clear serum beneath yields no growth. Finally, in the earliest stages of purulent change, the surface of the vesicle yielded growths, while the deeper portions of the same vesicle did not. The conclusion drawn, therefore, was that the purulent change was due to pyogenic organisms entering the vesicle from without, not through the lymphatic system, and that the pyogenic toxic element, referred to above, might be eliminated by two methods: (1) Prevention of suppuration in the vesicle; (2) by injecting an antitoxin.

PREVENTION OF SUPPURATION.

The possibility of preventing suppuration in the vesicles was tested through a series of 20 to 30 cases by the following methods. In the papular or early vesicular stage the forehead was well scrubbed with soap and water, washed with a solution of hydrarg. perchlor. (1-1000), and finally with ether. On this surface was painted collodion containing iodoform in suspension. The vesicles in this area did not undergo purulent change, though the coagulation or "creamy" stage was not prevented, also minor changes in the shape of the vesicles and depth of the resulting mark took place. Again, sterilisation of the arm was attempted, as in the forehead, except that liq. potassæ was also used, and a compress of a solution of hydrarg. perchlor. (1 in 1,000) was left on for six to twelve hours. Then the arm was carefully dressed in cyanide gauze, sal alembroth wool, and bandaged. The dressings were not disturbed till the pustules had well developed on the opposite arm; when removed it was found that purulent change had not taken place, but that the coagulation "creamy" change was present in the vesicles. The individual vesicle in the sterilised arm was rather smaller than on the exposed arm, the zone of inflammation surrounding each vesicle was less intense, and the patients asserted that the bandaged arm was less irritable. Success depended on the thoroughness of the preliminary sterilisation, the eruption should also be in an early stage. The difficulty of applying these methods clinically is manifestly great, and the results at best must be very uncertain.

SELECTION OF ANTITOXIN.

The view that pustulation is due to the access of ordinary pyogenic organisms was supported by the fact that convalescence from confluent attacks is generally interrupted, death occasionally brought about, by the large number of abscesses which develop all over the body, a condition resembling chronic pyæmia, and that crops of boils—the "acute furuncular diathesis" of Trousseau—were constantly present. The frequent presence of extensive cellulitis seemed to point to the streptococcus pyogenes aureus as the predominant organism, though the furuncles also indicated the presence of the staphylococci. Hence I determined to try what effect, if any, anti-streptococcic serum would have on this toxic condition. The serum was obtained from the Jenner Institute.

SELECTION OF CASES.

Those cases only were chosen in which the prognosis was gravest, and in which, judging by analogy from similar attacks in other patients, a fatal termination was to be expected. In these cases confluence occurred either just before or early in the pustular stage. When confluence occurred extensively in the coagulation stage, and was not followed by marked pustulation; death invariably, in the writer's experience, followed. The serum was injected aseptically into the subcutaneous tissue of the abdominal wall, either just before or coincident with pustulation, those injections being usually given at intervals of twelve to twenty-four hours. In addition, digitalis, with either tinct. ferri perchlor. or liq. strychninæ, was usually given four-hourly.

CASE I.—H. W., aged 29, a furnaceman, previously healthy. Admitted on April 15th with a very close-set papular eruption over face, and much hyperæmia of intervening skin, papules closely set on arms and legs, less so on trunk. Patient very restless during succeeding four days, violently delirious at night, trying to get out of bed.

April 19th. Temperature rising; vesicles large, confluent on face and hands

April 20th. Condition very grave, face and hands covered with a mass of confluent pustules, difficulty in swallowing, trunk thickly covered with eruption. Pulse 120, regular; respiration 25, slight tremors present on April 21st. Temperature, after a restless night, rapidly rose to 103.2°, and remained about the same level throughout the day. Serum, 10 c.cm., injected at 3.30 P.M. The condition throughout the day was critical.

April 22nd. Slightly quieter night after morphine; condition unchanged. Temperature did not rise above 102° through the day.

April 23rd. Delirious during night; temperature tending to rise. Serum, 10 c.cm., injected at 2 P.M.; pustules drying; general condition rather better.

April 25th. General condition better; quieter nights; seems stronger; face clearing. Pulse 108; respiration 22.

April 26th. Much worse; pulse 115, weak and irregular; respiration very shallow; temperature rose from 100.4° to 103°. Serum, 10 c.cm., injected at 6 P.M.

April 27th. Had a quiet night after morphine; seems better; clear-headed. He said he did not feel very ill, but excessively weak. From this date his temperature did not rise above 101°, and five days later reached normal. Here the temperature rather than the general condition was taken as an indication for serum, hence the long intervals between the injections. The rapidity of convalescence in this case was very remarkable, no complications of any kind arising.

Chart of Case I.

CASE II.—W. T., male, aged 6. Admitted May 3rd with a close set eruption of large papules; face very flushed; eruption very copious on arms, legs, and trunk. Patient progressed well during three days.

May 6th. Vesicles very large and pustular, delirious, refuses food, very restless, temperature 101.2°.

May 8th. Very restless and delirious, refuses nourishment, pulse 118, weak, respiration 25, cardiac sounds feeble, pustules confluent, condition very grave. Temperature in the evening rose to 102.2°. Serum, 10 c.cm., injected at midnight.

May 9th. Had rather quieter night; seemed better in morning but in afternoon became very restless. Pulse 130, weak; respiration 35, very shallow; temperature 103.4°; seems very ill; serum, 10 c.cm., injected at 10 P.M.

May 10th. Condition unchanged, very restless night, seems very weak; pulse still rapid, 125, and slightly irregular; temperature again reached 103.4°. Serum, 10 c.cm., was injected at 10 A.M.

May 12th. Patient for two days showed slight improvement but general condition was very grave. To day patient is quite rational, feels better, but is very weak since shortly after last injection. Temperature has been steadily falling.

The early part of convalescence was rapid and satisfactory, but later was complicated by a typically mild attack of scarlet fever, of which several cases had arisen in convalescent patients not treated with serum; the ultimate recovery was complete. Of the 2 other cases that recovered, the clinical course of the disease so much resembled that in the cases reported as not to require detailed description.

Chart of Case II.

CASE III.—W. B. was a healthy man of 35, who suffered from a very severe confluent attack. The patient early became delirious, pulse 120 to 130, weak; respirations 24 to 30, shallow. On eighth day of disease condition was still very bad; during preceding three days the temperature varied from 100° to 102°.

April 5th. Muttering delirium, at times semicomatose, pulse 128, weak

and irregular; respirations 25, temperature 102° ; serum, 10 c.cm., injected 10 P.M.

April 6th. Next day, twelve hours later, the temperature fell to 101.4° , but afterwards rose to 102.8° ; still very delirious, general condition rather worse; at 2 P.M. serum injected.

April 7th. The temperature gradually fell to 100.8° at 10 A.M., very delirious; pulse 140, feeble, slightly irregular; respirations 38, shallow, slight tremor present; temperature, 6 P.M., 103° ; serum, 10 c.cm., injected.

April 8th. Less delirium, feels better, highest temperature 102° ; subsequent progress was slow, but the temperature did not rise again above 102° , reaching 99° three days later; convalescence was delayed by furuncles, but no abscesses formed.

CASE IV.—J. P., a previously healthy man of 21, admitted with initial temperature of 101° . A very severe confluent attack, accompanied by much active delirium and some bronchitis.

April 21st. Pulse 118, weak; respirations 28, irregular, tremulous at times, tries to get out of bed. Patient very weak. Temperature 101.8° ; serum, 10 c.cm., injected at 6 P.M.

April 22nd. Temperature rose to 103° at 6 A.M., but gradually declined to 101° ; general condition rather better.

April 23rd. Very delirious, had to be tied in bed. In very exhausted condition; pustules drying up. At 10 A.M. the temperature rose to 104° ; serum, 10 c.cm., injected. Temperature, 10 P.M., 102° .

April 24th. Very weak; pulse 130, irregular; respirations 34, shallow. In the afternoon the temperature rose to 104° ; sponged, fell to 101° , later rose to 103° ; serum, 10 c.cm., injected at 6.30 P.M.

April 25th. The temperature has steadily fallen to 100.2° ; seems better, much less delirium; pulse 110, regular; respirations 25, very weak; temperature at 2 P.M., 103° ; sponged; temperature 101° ; subsequently temperature only exceeded 100° on one occasion. The prostrated condition lasted for several days, but convalescence was later very rapid, though accompanied by almost complete alopecia.

CASE V.—A woman, aged 44. Admitted on third day of disease (temperature 105°) with severe confluent attack. For five days after admission she was very restless. At times delirious; pulse 110 to 120, temperature fluctuating below 101° . On the fifth day the temperature reached normal; on the sixth day secondary fever began. During this period she was very restless and at times maniacal; the vesicles remained of small size, and early ran together in large irregular confluent areas. At this period the condition was very grave; pustulation was in progress, there was much vomiting and delirium; the pulse was 120, weak; respiration 25, cardiac sounds feeble; the temperature steadily rose, till at 10 P.M. on the ninth day it reached 103.2° . Serum, 10 c.cm., was then injected. Four hours later the temperature reached 104° , after which it steadily fell to 100° at 2 P.M. on the tenth day. At 10 P.M. the temperature again reached 103° . Serum, 10 c.cm., was injected during this period. The woman's condition was very critical; she was getting weaker, and evidently was losing ground. Eight hours after the serum the temperature fell to 101.4° , but later rose to 104.2° , when more serum was injected, followed by a fall of temperature to 103° ; death, however, occurred twelve hours later on the twelfth day.

CASE VI.—W. H., labourer, admitted on the fifth day, with very copious eruption on face, neck, hands, and arms, less copious on trunk; secondary pyrexia began on the eighth day, and pustulation occurred about the same time. The general condition and progress closely resembled that described in other cases. On the twelfth day the temperature reached 104° , when serum, 10 c.cm., was injected. In twelve hours the temperature fell to 101.8° . The pulse was 125, very weak and irregular. Respirations 25 twenty-four hours later. On the thirteenth day the temperature again reached 103.8° . More serum was then injected, with the result that the temperature fell in sixteen hours to 101° on the fourteenth day. From this date till the eighteenth day the temperature only once reached 102° , the patient seemed stronger and took food well, though pustules cleared up slowly. On the eighteenth day the condition became worse, very restless; irregular pulse, 130, and some tremor; temperature 103° ; more serum injected, resulting in a fall of temperature to 100.2° in twelve hours. Temperature again began to rise, and patient, who showed signs of cardiac failure, died four days later, the temperature for eight hours before death being 107° . The long interval preceding the third injection was due to accidental causes.

In earlier cases serum injections were repeated only as the temperature rose after a previous injection, and the interval between the injections therefore varied. Later, however, better results were obtained after injections at regular intervals of twenty-four hours, beginning, as a rule, with the onset of pustulation. In no case were more than three injections given. Probably in some cases the injections should be continued for longer periods, or the amount of the earlier injections increased.

In cases treated with serum, the critical period following the absorption of the pustules was shortened, the intensity of the toxæmia seemed to be less, there was much less tendency to cardiac failure or collapse. The pulse throughout, though fast, was regular, and the strength of the beat was well maintained. Convalescence set in earlier and progressed rapidly. These patients were out of danger, and out of bed, before others with attacks of similar intensity. Most important of all, not one of the four patients treated with serum who recovered developed a single abscess throughout the convalescence. In cases of similar severity it was often necessary to make from 30 to 60 incisions in the same patient for abscesses or cellulitis, and many patients died in this condition of semi-pyæmia. That the severity of attack in these reported cases was as great as in those often fatal, where complications above referred to arose, is shown by the fact that

they, too, suffered from acute furunculosis, almost complete alopecia, and flesh was lost as rapidly as in other cases. No local pain, local abscess, arthritis, or hyperpyrexia ensued in any of these cases.

BACTERIOLOGY.

To establish bacteriologically the value, whether great or little, of this treatment, would require a prolonged and careful bacteriological examination of the vesicles, pustules, and resulting abscesses in all their stages, such as the writer was unable to carry out. More or less roughly, by cultivations on ordinary media, organisms were obtained from pustules and from the pus of abscesses giving reactions similar to streptococci, and in a few cases to staphylococci. The contents of a vesicle, which appeared to have undergone purulent change in its interior, rather than on the surface, was examined by the Clinical Research Association, they reported staphylococcus aureus and staphylococcus albus present.

The bacteriology of variola is very incompletely known. Cohn¹ found cocci in variolous lymph, and this observation was confirmed later by Burdon Sanderson. Weigert² found the lymphatics in the region of a pustule, both in human small-pox and in sheep-pox, filled with cocci which were later named streptococcus variolæ et vaccinæ. Hlava³ established the presence of streptococcus pyogenes in small-pox pustules. Garre⁴ found streptococci in the internal organs in a case of variola hæmorrhagica, and in a fatal case complicated by pemphigus he found the streptococcus pyogenes aureus in the pemphigus vesicles. Steinberg⁵ says that the microbes found in variolous pus are the usual pyogenic organisms.

Finally, the object of this paper is to set forth clinical facts that occurred under the writer's observation; not to propose a theory, but to leave others, whose opportunities for treating variola patients and for making the necessarily extended bacteriological inquiry are more prolonged than was the writer's, to test by a fair trial the value of this treatment.

REFERENCES.

- ¹Supplementary Local Government Board Report, 1892-93. ²Journal of Bacteriology and Pathology, vol. ii. ³Virchow's Archiv, Bd. lv, 1878. ⁴Anatom. Beitr., 2, L. v. d. Pocken. ⁵Quoted by Cruickshank, Manual of Bacteriology of Infectious Diseases. ⁶Fortschritte d. Med., 165, 1885. ⁷Manual of Bacteriology.

THE TREATMENT OF INGUINAL HERNIA IN CHILDREN.*

By W. McADAM ECCLES, M.S.LOND., F.R.C.S.ENG.,
Assistant-Surgeon to the West London Hospital, and to the City of London Truss Society, and Demonstrator of Operative Surgery at St. Bartholomew's Hospital, London.

INGUINAL hernia in children is a curable affection. Nature herself tends to bring about a cure, and it should be the aim of the surgeon to aid her in this. Yet it is a fact that there are very divergent opinions as to the best method in which art may thus assist the natural disappearance of the condition.

It is only my intention to discuss the treatment of the protrusion; yet to understand this thoroughly a few points in the etiology and the anatomy of the affection may not be out of place.

ANATOMICAL CONSIDERATIONS.

In both sexes a finger-like process of parietal peritoneum passes down through the inguinal canal, in the male accompanying the testis, and in the female the round ligament of the uterus. It is usual for this prolongation of the serous membrane to become obliterated and converted into a fibrous cord, wholly in the female, and except the lowest part in the male, this latter portion remaining as the tunica vaginalis. This closure should be complete very soon after, if not before, birth. Any failure in the obliteration of the process predisposes to protrusion of some of the abdominal viscera into the ready-formed sac. But I firmly believe that a very considerable proportion of infants who have a patent processus vaginalis never become the subjects of inguinal herniæ, and therefore it is necessary to look further for an exciting cause other than this predisposing condition.

* Read before the West Surrey District of the South-Eastern Branch of the British Medical Association.