flexion of the foot on the leg, of the leg on the thigh, of the thigh on the pelvis, a flexion of the toes on the metatarsus. In some persons, on exciting the sole the toes remain to all appearance motionless, but they do not ever (in health) execute a movement of extension. This applies especially to the big toe. In certain pathological conditions, however, excitation of the sole provokes extension of the toes, especially of the big toe. This modification of the reflex he calls the "toe phenomenon." Without proceeding to the consideration of various details, I may say that in all the cases in which Babinski has observed this phenomenon its relation to disorder of the pyramidal system has been incontestably established, and he has not once noted its existence in a case in which the pyramidal system was certainly in a state of integrity. According to his experience it is never seen in hysterical paraplegia, of flaccid or spastic character. During the last five or six months I have had the opportunity of testing the observation in a variety of cases, and especially in many examples of insular sclerosis, and am able so far to confirm the exactness of Babinski's description. It appears, therefore, that by means of this very interesting test our power of distinguishing between organic and functional paraplegia is likely to be greatly increased. If the foot-sole, on being tickled, responds by a movement of extension of the big toe (care being taken to be sure that it is a movement of extension of the metatarso-phalangeal joint) we may take it as tolerably certain that there is an affection of the pyramidal tract, and that we have not to do with a case of functional paraplegia. In insular sclerosis it must be a rare event for the pyramidal tract in some part, at least, of its extent, not to be invaded by islets of sclerosis, and the value of the test is therefore strikingly exhibited. Time will not permit me to dwell further on this interesting discovery. The subject has been ably discussed by Dr. James Collier, and I cannot do better than refer you to a contribution from him which appears in the spring number of *Brain*.⁴

NOTES AND REFERENCES.

¹ *Simulation of Hysteria by Organic Diseases of the Nervous System*, 1891; Atrophy of the Optic Nerve as a Symptom of Chronic Disease of the Central Nervous System, *BRITISH MEDICAL JOURNAL*, October 7th, 1893; Insular Sclerosis and Hysteria, *Lancet*, January 2nd, 1897; Retro-ocular Neuritis, *Optic. Soc. Trans.*, 1897. ² Absence of reflex from the sole of the foot is a very constant symptom in hysterical paraplegia..... Persistence in the application of a stimulus to the sole will, in hysterical cases, frequently but not always succeed in provoking a response after a few minutes.—Clinical Lecture on Diseases of the Nervous System, 1882. ³ See especially Le Phénomène des Orteils, *La Presse Médicale*, July 27th, 1898. ⁴ An Investigation upon the Plantar Reflex, with reference to the significance of its variations under pathological conditions, including an inquiry into the etiology of acquired pes cavus.

THE LOCALISATION AND SYMPTOMS OF DISEASE OF THE CEREBELLUM CONSIDERED IN RELATION TO ITS ANATOMICAL CONNECTIONS.*

By ALEXANDER BRUCE, M.A., M.D., F.R.C.P.E.,
Assistant Physician, Royal Infirmary, Edinburgh.

IN bringing this communication before the Medico-Chirurgical Society to-night, my main object is to draw attention to the importance of certain nerve tracts, which have been proved by the recent investigations of neurologists to bring the cerebellum into intimate relation with the spinal cord and higher parts of the central nervous axis.

I do not propose to approach the subject from the point of view of the anatomist, but rather with the intention of impressing upon the clinician that a knowledge of their position and relations, when considered in connection with the results of experiment on the cerebellum and its peduncles, may afford valuable data for the explanation of some of the symptoms of cerebellar disease, and may assist him in the difficult task of its localisation.

The symptoms of cerebellar disease may for our purpose to-night be divided into two classes: first, those which are without special significance beyond the fact that they indicate the

existence of an intracranial lesion, but which have no further localising values; and, secondly, those which are regarded as being more definitely characteristic of affections of the cerebellum. Among the former may be placed headache, optic neuritis, hebetude, and vomiting; among the latter the disturbances of equilibrium and co-ordination indicated by the peculiar reeling gait, the ataxia, the severe vertigo, and the deviations and oscillations of the eyeballs.

One important and puzzling feature of cerebellar disease is that these characteristic symptoms, as is well known, are not invariably present in all cerebellar lesions, for some such lesions may run a course which is more or less latent, or may be indicated by symptoms of so vague a nature as, in the present state of our knowledge, to be practically useless for localising purposes; and on the other hand similar symptoms may result from lesions which do not affect the cerebellum directly. One of the first steps towards the explanation of these peculiar variations in the phenomena of cerebellar disease was the recognition of the fact that these differences in the symptoms are not due to differences in the nature of the lesion (occurring as they do equally in tumour, hæmorrhage, softening, and sclerosis), but depend, as Nothnagel was the first to show, on differences in its situation. This author states in his *Topische Diagnostik der Gehirnkrankheiten* that "the conclusion to be drawn from all these observations is that the situation of the lesion is the factor that determines the presence or absence of symptoms; only when the lesion is limited to one hemisphere can it run the course without symptoms" (p. 47). "There are certainly many cases which show that lesions in the hemispheres may also produce symptoms....., but the analysis of these cases shows, in our opinion, without doubt, that the lesion in such cases involved adjacent parts in one way or other." It may be concluded, therefore, that the loss of the substance of one hemisphere does not of necessity produce any symptoms, or at least any symptoms as yet known to us. "Lesions actually limited to one hemisphere cannot be diagnosed" (p. 48). "Disturbances of co-ordination only occur when the lesion indirectly or directly involves the vermis—that is, the middle lobe of the cerebellum" (p. 50).

"Another question (p. 51) is whether disturbances of co-ordination must invariably appear when the vermis is involved. In my earlier publications I answered this in the affirmative on the strength of the observations then at my disposal."

"The later observations show undoubtedly that tumours may be situated in the vermis without causing any symptom of cerebellar ataxia, and therefore lesions of the vermis need not be invariably associated with disturbances of co-ordination."

"In this connection I may mention that these exceptions are always cases of tumour—that is, of slowly-growing lesions, and that in all cases a greater or less part of the vermis was left intact."

Nothnagel is frequently credited with the unqualified statement that it is only tumours in the middle lobe that cause disturbances of equilibrium, and that tumours of the lateral lobes always run a latent course. From what has just been quoted, however, it will be seen that this is not a correct interpretation of his opinion.

I shall now endeavour to explain from the anatomical standpoint why it is that disturbances of equilibrium occur specially in connection with lesions of the middle lobe, and why in certain instances these disturbances may also be present in lesions of the lateral lobe. The anatomical data have been obtained partly by the collation of observations of various investigators on the human fœtus, partly by the study of degenerations following experimental and pathological lesions of the brain and cord.

The cortex of the middle or vermiform lobe of the cerebellum forms the central termination for at least six important tracts derived from the spinal cord, two arising from the cord directly and the remainder from the nuclei of the postero-medial and postero-external columns, namely, the clavate and cuneate nuclei (two arising from these nuclei on the same side and two from those of the opposite side, each represented by a single line in Fig. 1). The uninterrupted fibres consist of the direct cerebellar tract and the ascending antero-lateral tract of Gowers. All these tracts, except the last mentioned,

*Read before the Medico-Chirurgical Society, Edinburgh.

are contained in the restiform body or inferior peduncle of the cerebellum.

Fig. 1.—The constituents of the inferior peduncle. P. C. N., Each oval area represents the combined nuclei of the two divisions of the posterior columns, and gives a double tract represented by a single line to the restiform body of its own side and another to the opposite side. These represent the interrupted paths from the posterior column. R. N., roof nuclei. C. V. T., cerebro-vestibular tract from roof nuclei to nucleus of Deiters.

The tract of Gowers passes further up the pons and hooks round the fifth nerve before bending into the vermis, as has been shown by Loewenthal, Mott, and Tooth in animals and by myself in man. The majority of the fibres terminate in the upper part of the middle lobe, some on the same side and some after crossing the middle line.

While it must be admitted that the functions of these tracts are still somewhat doubtful, it is certain that they are not concerned in the conduction of impulses which give rise to impressions of touch, temperature, or pain, since they may be destroyed without interference with these forms of sensibility, and it must be presumed that some, if not all, of them, are concerned in the conduction of muscular sense impressions. It is important to note that none of these tracts pass to the lateral lobes, nor does any other tract from the cord, as far as is known, enter the lateral hemispheres or become connected with them, except indirectly, and even then to a comparatively slight extent. These tracts are all undoubtedly afferent to the cerebellum.

Another tract which was at one time regarded as a purely afferent sensory tract, and was therefore termed by Edinger "the direct sensory cerebellar tract," connects the nucleus of Deiters (one of the end nuclei of the vestibular nerve, lying at the lateral angle of the fourth ventricle), with the roof nuclei of the cerebellum.

Until quite recently the relations of Deiters's nucleus remained very obscure. Its most characteristic feature, and one that has impressed itself on most observers, is the large size of its cells, which in appearance are practically identical with the motor cells of the spinal cord. So close is this

resemblance that, from the first time my attention was directed to these cells; I felt convinced that they must belong to a motor nucleus, although at that time it was not possible to trace their connections. Subsequent research has, however, determined to a very great extent the destination of the fibres derived from this nucleus. Although our knowledge of its relations is still far from being complete, enough has been learned to indicate its great importance. Monakow, in 1883, showed that the cells of this nucleus became atrophied after hemisection of the cord in the cervical region. The significance of this fact was, perhaps, not at first fully apparent, but the discovery of the existence and nature of the neuron showed that its cells must give origin to fibres which passed down the cord, and which had been divided in the above-mentioned operation. This view had been actually advanced by Deiters himself long before this. In 1888-89-92 I described two tracts arising from this nucleus; of these one passed in a downward direction, and could be followed as far as the outer side of the inferior olive; but, owing to the method of investigation then employed, it was not possible to trace it further.¹ Held subsequently observed that it was continued downwards into the antero-lateral columns of the cord, and more recent investigations have confirmed this view. The other tract passed inwards under the floor of the fourth ventricle, and, after giving a certain proportion of its fibres to the sixth nucleus, it crossed the middle line and apparently passed upwards in the posterior longitudinal fasciculus as high as the third nucleus. More recently our knowledge of the connection of these two tracts has been further extended by the study of sections stained by Marchi's method after recent injury to the nucleus of Deiters.² It is

Fig. 2.—Scheme of the connections of the nucleus of Deiters. now known that the first tract—which may be called the vestibulo-spinal tract—passes downwards and enters the

same side of the cord, where it divides into two parts, one division taking up its position at the side of the anterior median fissure and the other in the lateral column. Risien Russell has traced some of these fibres as far as the lower end of the thoracic region of the cord. They terminate in the anterior cornu of the cord on their own side at various levels.

The second set of fibres, that which passes inwards under the floor of the fourth ventricle, divides into several groups. One of these terminates in the sixth nucleus of the same side; the remaining groups enter the posterior longitudinal fasciculi (P. L. F.) on both sides, some turning upwards and some downwards. The two groups which bend upwards terminate in relation to the two third nuclei; those which have a descending direction pass down the anterior columns of the cord, where they occupy a position at the margin of the anterior median fissure. They appear to terminate by entering the anterior cornua of the cord.

It will be seen that, by means of these various tracts, the nucleus of Deiters is placed in a position to influence the nuclei of the ocular nerves and the motor cells of the cord throughout the greater part of its extent.

The nucleus of Deiters has another very important connec-

Fig. 3.—Diagrammatic transverse section of cerebellum to show connections of middle lobe: C. D., nucleus dentatus; S. C. P., superior cerebellar peduncle; C. R., inferior peduncle.

tion that can be traced backwards from it along the side of the fourth ventricle, then across the mesial plane of the cerebellum to terminate in a nucleus which is called, from its position in relation to the fourth ventricle, the roof nucleus. This tract was formerly regarded as the continuation into the cerebellum of the auditory nerve, and even of the lower cranial sensory nerves, and, as already said, was termed the "direct sensory cerebellar tract" by Edinger, but Ferrier and Turner, and more recently Risien Russell and André Thomas, have shown that the tract has its origin in the roof nucleus, and is thus not an afferent but an efferent tract, arising in the middle lobe of the cerebellum, and terminating in Deiters's nucleus.

When the middle lobe of the cerebellum is further examined (Figs. 2 and 3), it can be seen that its cortex is connected by a series of sagittal fibres with the roof nuclei. This important link completes a chain of afferent, internuncial, and efferent fibres between certain sensory tracts in the cord, the middle lobe of the cerebellum, the nucleus of Deiters, the various oculo-motor nuclei, and certain motor centres in the cord. The integrity of this chain appears to be essential to the perfect maintenance of equilibrium. If we follow the course of the fibres in this system (as indicated in Fig. 2), we see that an impulse would travel upwards through the restiform body (C. R.) and Gowers's tract (not drawn) to the cortex of the middle lobe of the cerebellum (superior vermis). It would then pass by means of the sagittal fibres (S. F.) to the roof nuclei (R. N.). Starting again from one of them (the connection with the other roof nucleus being ignored for the sake of simplicity), it would pass by the cerebello-vestibular tract to the nucleus of Deiters. Here there might be a reinforcement by other impulses travelling from the semi-circular canals (S. C.) by way of the vestibular nerve (VIII.). At Deiters's nucleus fibres to the nuclei of the sixth and third, and perhaps the fourth nerve, as well as to the anterior cornua of the cord, mainly on the same side, would be simu-

lated, and in this way there would be put into operation a very complex mechanism which is well adapted to affect the equilibrium of the body. We can thus understand how irregular or defective stimulation of Deiters's nucleus may lead to lateral oscillations of the eyeballs, and to irregular or impaired movements of the same side of the body, resulting in vertigo and disturbances of equilibrium.

It will be seen from what has just been said that the cerebellar termini of the constituents of this system are in the cortex and roof nuclei of the middle lobe; no such connection of the spinal cord or its nuclei with the lateral lobes of the cerebellum can be made out. These lobes seem to receive afferent fibres from the nuclei in the pons, which, in their turn, are in relation with the cortex of the frontal and temporal lobes of the cerebrum.

If we now consider the system of the superior cerebellar peduncle, which is also undoubtedly concerned in the maintenance of equilibrium, we see that this peduncle arises mainly from the nucleus dentatus, and terminates above in the red nucleus and optic thalamus of the opposite side. The nucleus dentatus is situated at the junction of the middle and lateral lobes, and is connected extensively with both of these portions of the cerebellum, as well as, through the inferior peduncles, with the inferior olives of the medulla (see Fig. 1).

It will be observed from Fig. 3 that this nucleus is in such a position as to be readily affected by lesions involving either the middle or the lateral lobes. It will be seen, for instance, that a section dividing the middle from the lateral lobes will cut through the nucleus dentatus, thus severing the origin of the superior peduncle, and may at the same time cut across both the afferent and efferent tracts of the middle lobe—namely, the inferior peduncle and the cerebello-vestibular tract. From these facts we can readily understand why it is that removal of one-half the cerebellum or of one lateral lobe causes practically the same symptoms as removal of one-half of the middle lobe, since in both cases the afferent and efferent fibres of one side of the middle lobe are divided, and the superior cerebellar peduncle arising from the nucleus dentatus is also involved to a greater or less extent. We are also in a position to understand why a section in the mesial plane of the middle lobe may cause few symptoms of disturbances of equilibration, and why these disturbances may be readily recovered from. Such a section will destroy the afferent and efferent fibres from both sides to an equal extent, so that the resulting interference with equilibration will be merely a symmetrical weakening of that function on both sides. This can be compensated for to a great extent by that portion of the mechanism for equilibration which has remained unaffected by the lesion. For the same reason a tumour of the vermis occupying a symmetrical position with reference to the middle line, or at least which destroys, symmetrically, afferent and efferent fibres of the vermis, would be marked by few or no symptoms, especially if its growth were so slow as to allow of compensation. The pronounced symptoms in the case of unilateral lesions are due undoubtedly to the fact that the interference with the mechanism is limited to one side, and the results produced are therefore asymmetrical. If one cerebello-vestibular tract or one nucleus of Deiters is alone affected, the usual stimuli from the latter to the nuclei of the ocular nerves and to the anterior cornua of the corresponding side of the spinal cord will be wanting; thus we can understand (1) the yielding of the limbs of the same side and the tendency to fall to that side, and (2) the oscillatory movements of the eyeballs.

These symptoms may, however, be absent after lesions of the lateral lobe external to these tracts, which would leave these systems entirely, or practically entirely, intact. Turner,³ in reviewing Lucian's work, states that "in our experiments rotation round the longitudinal axis certainly occurs more constantly when the peduncles are divided than when the lateral lobes are removed. But this is not an absolute rule, and one of the most marked instances of rotation round the vertical axis which we observed was an animal in which only the cortex of the lateral lobe had been cauterised and under conditions likely to excite irritation of neighbouring parts." This statement may be regarded as being so far in harmony with the view just enunciated.

Thomas⁴ expresses himself even more unequivocally:

"Lesions involving a considerable extent of the cortex, but not extending deeply and not implicating the central grey nuclei, produce disturbances which are of relatively short duration and of moderate intensity. Lesions limited to the cortex of one hemisphere, and not extending deeply, may be free from any disturbance."

It is well known that section of a superior or inferior cerebellar peduncle produces rotatory movements, and other motor disturbances, and also that similar disturbances follow removal of one half of the vermis or of one lateral lobe. This second fact may be readily understood when the arrangement of the parts is considered. An examination of Fig. 3 shows that removal of one half the vermis or of one lateral lobe involves a section of the intracerebellar portion of these peduncles. It is also stated that section of the middle peduncles produces somewhat similar symptoms, and since the cerebellar terminations of this peduncle are limited to the lateral lobe, it will naturally be asked why removal of the lateral lobe, which must involve an intracerebellar section of the fibres of this peduncle, should not in like manner produce similar symptoms. In answer to this, I would suggest that many of the results ascribed to section of the middle peduncle are really due to involvement of the upper or lower peduncle or of the cerebello-vestibular tract, or of all of these, during the operation employed for division of the middle peduncle, and that the results of a section limited to the middle peduncle have yet to be determined.

If the views here adduced be well founded, we may expect disturbances of equilibrium to be produced by lesions situated within an area bounded by the intracerebellar path of the two inferior peduncles, of the two superior peduncles, and the dentate nuclei in which the latter arise. This area contains the middle lobe (superior and inferior vermis, the roof nuclei, and the sagittal fibres connecting the latter with the cortex), and the cerebello-vestibular tracts from the roof nuclei to the nucleus of Deiters (Fig. 3). Lesions within this area may produce no such disturbances, provided they are symmetrically situated with reference to the mesial plane, and especially if their growth is so slow that compensation is established *pari passu* with the disturbances they may tend to cause.

On the other hand, lesions situated in the lateral lobes may produce no disturbance of equilibrium provided they are situated entirely external to the intracerebellar paths of the upper and lower peduncles, and of the nucleus dentatus (area of possible latency Fig. 3). If, however, these structures are interfered with, either by pressure or by direct involvement, then the characteristic symptoms of cerebellar disease will be produced, and will depend in their character and amount on the nature and extent of this interference.

Fig. 4. Relation of Deiters's nucleus to the mechanism for conjugate deviation.

If the cerebello-vestibular tract, or Deiters's nucleus, be in-

jured, then the usual stimuli will not pass either to the anterior cornua of the cord or to the sixth (fourth) or third nuclei. Hence may result the weakness of the same side, the tendency to fall to that side, the impairment of the conjugate deviation to that side, the tendency of both eyes to be directed to the opposite side, and the lateral nystagmus which occurs especially when the eyes are directed towards the same side. A glance at Fig. 2 will show how the spinal symptoms may be produced, since the nucleus of Deiters will not act through the vestibulo-spinal tract and posterior longitudinal fasciculi upon the anterior cornua, and more especially upon that of the same side. The effect on the position and movements of the eyes will be easily understood from Fig. 4, if it be remembered that the sixth nucleus is the nucleus for the conjugate deviation of the eye to the same side, supplying, as it does, not only the external rectus, but also, through the opposite third nucleus and nerve, the internal rectus of the opposite side. A stimulus of Deiters's nucleus, acting on the sixth nucleus, will turn the eyes to the same side; absence of this stimulus will allow the other sixth nucleus to pull the eyes towards the opposite side and will thus weaken the voluntary attempt to look to the same side. On the other hand, lesions which merely irritate but do not destroy the nucleus of Deiters may cause rigidity or spasms of the same side of the body, or such an excess of movement as may cause the body to fall to the other side, and by acting on the sixth nucleus may turn the eyes to the same side.

Luciani, Ballance, and Thomas have also drawn attention to the possible effect which the superior peduncle may exert through the red nucleus and the optic thalamus on the ocular nuclei and the opposite motor nucleus, and it is not my intention to question this influence in producing the symptoms of the cerebellar disease. At the same time, the importance of the nucleus of Deiters as a co-ordinating centre seems too evident to be ignored.

It is by no means claimed that the above described mechanism is as yet so fully understood as to be capable of explaining all the symptoms of cerebellar disease, but it, at least, affords a working hypothesis on which the most characteristic and frequent among them may be interpreted.

REFERENCES.

- ¹ *Proceedings of the Royal Society of Edinburgh*, xvii, 1883-1884, p. 26.
- ² Ferrier and Turner, Risien Russell, and André Thomas. ³ *Brain*, 1884, p. 7. ⁴ *Le Cervelet*, p. 317.

ON THE PHYSIOLOGICAL ACTION OF CHOLINE AND NEURINE.*

BY
F. W. MOTT, M.D., F.R.S., Pathologist to the London County Asylums.
and
W. D. HALLIBURTON, M.D., F.R.S., Professor of Physiology, King's College, London.

THE cerebro-spinal fluid removed from cases of brain atrophy, particularly from cases of general paralysis of the insane, produces when injected into the circulation of anæsthetised animals (dogs, cats, rabbits), a fall of arterial blood pressure, with little or no effect on respiration. This pathological fluid is richer in proteid matter than the normal fluid, and among the proteids, nucleo-proteid is present. The fall of blood pressure is, however, due not to proteid, nor to inorganic constituents, but to an organic substance which is soluble in alcohol. This substance is precipitable by phospho-tungstic acid, and by chemical methods was identified as choline. The crystals of the platinum double salt, which, when crystallised from 15 per cent. alcohol, are characteristic octahedra, form the most convenient test for the separation and identification of this base.

The nucleo-proteid and choline doubtless originate from the disintegration of the brain tissue, and their presence indicates that possibly some of the symptoms of general paralysis may be due to auto-intoxication; these substances pass into the blood, for the cerebro-spinal fluid functions as the lymph of the central nervous system. We have identified choline in the blood removed by venesection from these patients