majority of cases as it can be more rapidly performed, it presents perfect security against sepsis by escape of the contents of the stomach, provided that the opening of the stomach is delayed for a day or two, and it is possible to dispense with the use of buried sutures, which in Witzel's operation cannot be avoided.

SUCCESSFUL REMOVAL OF AN ENLARGED AND DISPLACED SPLEEN.

By D'ARCY POWER, F.R.C.S.ENG.,

Assistant Surgeon to, and Teacher of Surgery at, St. Bartholomew's Hospital; Surgeon to the Victoria Hospital for Children, Chelsea.

A WIDOW lady, aged 43, mother of two children, the youngest being 25, consulted my friend, Dr. W. S. A. Griffith, for an abdominal tumour which was thought to be ovarian. Examination showed that the uterus and its appendages were normal, and the patient then came under my care.

History.—Three and a half years previously she struck her left side just over the lower ribs whilst she was moving a piece of furniture, and she dates the beginning of her abdominal discomfort very definitely from the time of the blow, though it was not until two years later that she noticed the lower part of her abdomen increasing in size. She has suffered much pain and inconvenience from the swelling during the last twelve months, and she has been losing flesh lately and has had attacks of indigestion and vomiting.

Anamnesis.—The patient was a healthy-looking woman with a florid complexion, who complained less of the inconvenience of the tumour than of the remarks to which it gave rise, for her abdomen was as large and prominent as it had been during the sixth and seventh months of her former pregnancies. Examination of the abdomen showed a large tumour filling the whole of the left side and extending to the right of the umbilicus. The surface was smooth and there was a definite edge, notched at its lower part. It was freely movable from side to side, but any attempt to push it upwards caused pain, and it was tender at the upper part just beneath the ribs. The tumour was dull on percussion in the flank, but resonant in front where the intestines lay above it.

Operation.—I opened the abdomen at 3.30 P.M. on Wednesday, September 6th, 1899, in the presence of Mr. S. T. Huggins and his son, Dr. S. P. Huggins, making a 5-inch incision in the left linea semilunaris. The spleen was easily delivered from the abdominal cavity, and measured 8 inches in length. It was attached by so long a pedicle that it hung free by the side of the patient, and almost lay upon the operating table. The patient showed signs of collapse when the pedicle was stretched, but recovered as soon as the spleen was supported. The pedicle measured 4 inches across, and consisted of fatty connective tissue with many large veins; the tortuous splenic artery was easily seen lying in the centre. The connective tissue overlying the artery was first dissected away for about an inch, and a double silk ligature was passed beneath the vessel, which was tied in two places about half an inch apart and divided. The pedicle was afterwards ligatured in half-inch segments from below upwards and was divided. The cut end of the pedicle was carefully examined to ascertain whether any vessel had escaped ligature, and the operation was delayed for nearly ten minutes until I had satisfied myself that it was absolutely free from oozing. When everything appeared secure the stump of the pedicle was returned to the abdomen. The suspensory ligament was only represented by a little loose connective tissue, and gave no trouble, as it was secured by a single ligature of silk. The wound was closed by a continuous suture of fine silk through the peritoneum, three interrupted sutures of sterile catgut through the abdominal muscles and aponeurosis of the external oblique, and a continuous suture of silkworm gut through the skin. There was very little hæmorrhage during the operation, but the spleen contained a good deal of blood, which ran out and clotted after the organ had been removed. The operation lasted an hour and a-half from the beginning of the anæsthetic—gas and ether being administered during the first three-quarters of an hour, chloroform afterwards—to the application of the dressing, some part of the time being occupied in making the skin aseptic.

After-History.—The patient bore the operation excellently, her pulse being 96 and of good volume after she had been put to bed. She passed a fairly good night, and on the following day was only complaining of a little flatulence, which soon passed off. From the time of the operation until the fifth day the patient presented evidence of sympathetic irritation in a curiously agonised expression, of which she was quite unaware, reminding me of those who suffer from abdominal cancer or commencing tetanus. It seemed to be due to a slight tonic contraction of the platysma and its connections, but in no other way was there any indication of sympathetic disturbance, and the patient was quite cheerful. She remained comfortable for many days with a normal temperature and a pulse of 64. The stitches were removed on the seventh day, when the wound was healed except for a slight stitch abscess at the lower part, which had disappeared on the ninth day. I then went away for my summer holiday, and on my return found that she had progressed satisfactorily until September 28th—the twenty-second day after the operation—when her temperature began to rise. Two days later she was suddenly seized with intense pain in her left side, whilst the action of the heart and diaphragm was impeded, and she was thought to be dying. The symptoms, which were evidently very alarming for a short time, passed off spontaneously, and were succeeded by a similar attack two days later. Mr. Walsham then kindly saw the patient in my absence, and opened a localised subdiaphragmatic abscess by an incision carried parallel to the scar of the former wound, and about 2 inches to the outer side. There was free bleeding for several days, apparently from a small artery at the bottom of the abscess cavity, pus was discharged by the bowel, and a small fecal fistula formed. The fistula soon closed, leaving only a small sinus through which

a silk ligature came away from time to time. The general condition of the patient remained good; she increased in weight; her temperature again became normal; and she went home on October 18th, the forty-second day after the operation. I have since seen her on two or three occasions, the last time being in September, 1900, and except for the sinus and some occasional colicky pains on the left side, she appears to be well, and is meditating a return to her former profession. Physiologically, she has sustained no inconvenience from the loss of her spleen.

Description of Spleen Removed.—The spleen was estimated to weigh about 2½ lbs. immediately after the operation, and whilst it was full of blood. It weighed between 17 and 18 ozs. after it had been kept in spirit for six months. The enlargement was due to simple hypertrophy.

Blood Examination.—Dr. J. H. Drysdale, who kindly examined the blood of the patient on September 14th, 1899, the eighth day after the operation, reports: "Hæmoglobin, 69 per cent.; red cells, 4,230,000; number of leucocytes, 17,000; lymphocytes, 18.75 per cent.; large mononuclear, 12.5 per cent.; polymorphonuclear, 65.5 per cent.; eosinophile, 3.5 per cent. There were no myelocytes or atypical cells. The blood was taken before the midday meal, the patient having had a boiled egg, toast, and coffee for breakfast, with lamb, vegetables, and a fried sole on the previous day. *Remarks.*—Leucocytosis, the large lymphocytes (mononuclear), relatively as well as positively increased."

The case bears out the conclusion that the splenectomy is a successful operation in cases of enlarged and displaced spleens when the enlargement is not leucocythæmic. Some credit may be given to the patient's statement that the whole mischief started from an injury over the region of the spleen, as she was a healthy woman who had not been pregnant for many years, but had not approached the menopause, and consequently there had been no disturbance of the relations of her abdominal contents. The operation showed that some absorption of the suspensory ligament had enabled the spleen to weigh anchor and sail away into the depths of the peritoneal cavity, whilst it also proved that the shock attending manipulation of the enlarged organ is due chiefly to impulses communicated through the pedicle, and not, as some have suggested, to interference with the diaphragm through the suspensory ligament. The gradual elongation of the pedicle led to alterations in the vascular supply causing chronic congestion of the spleen ending in hypertrophy. The etiology of the enlargement, therefore, is somewhat clearer than usual. I was somewhat surprised during the operation to find how little the spleen shrank when it was handled, due apparently to the fact that the muscular tissue had not increased proportionately to the other elements in its structure. The formation of an abscess near the pedicle was a most unfortunate occurrence, and marred what would have otherwise been a very successful result. It was due, apparently, to a contamination of one of the silk ligatures used in securing the pedicle. The organism, however, could not have been very virulent, as the patient seemed to be perfectly well for twenty-two days after its introduction. The sudden and alarming effects which the abscess produced when it reached the under surface of the diaphragm are well worthy of remembrance.

The interesting statistics collected by Dr. Vanverts¹ show that between 1876 and 1897 16 cases of movable, displaced, and enlarged spleen had been removed with a single death. All the cases occurred in women.

I owe my best thanks to Mr. S. T. Huggins for his kind and skilful attendance upon the patient during my absence, and to his son, Dr. S. P. Huggins, formerly my House-Surgeon at St. Bartholomew's Hospital, for taking careful notes of the case.

REFERENCE.

¹ *De la Splenotomie*, Paris: Steinheil, 1897.

ABDOMINAL TUMOUR (INCLUDED FŒTUS)

OCCURRING IN A CHILD AGED 3 MONTHS: LAPAROTOMY: DEATH.

BY

G. A. WRIGHT,

and

D. S. WYLIE,

M.B., F.R.C.S.,

M.B., Ch.B.,

Surgeon, Manchester Royal Infirmary and Manchester Children's Hospital.

Late Senior Resident Medical Officer, Manchester Children's Hospital.

[With a report on the Specimen by J. W. BALLANTYNE, M.D., F.R.C.P.E.; and drawings by W. E. FOTHERGILL, M.D.]

HISTORY (BY DR. WYLIE).

THE child was a female, aged 2 months when admitted to the Manchester Hospital for Sick Children on March 10th, 1899. The mother gave the following history: The child's abdomen had been swollen since birth, labour having been long and