teacher than to aid the comprehension of the student. Our pathological museums are chaotic, and therefore repellent to the student. Medicine is a science without a method, and until our leaders adopt one we shall not take the place that belongs to us.

THE RELATION BETWEEN CAUSE AND EFFECT IN DISEASE.*

By CHARLES POWELL WHITE, M.D., F.R.C.S.,
Demonstrator in Pathology, Yorkshire College, Leeds.

In considering the pathology of diseases we often have occasion to use the term "specific." It is well, then, to ask ourselves what this term means. A "specific disease" is literally a disease which has certain definite characters which distinguish it from all other diseases. There are two ways in which a disease may be specific.

1. It may be due to a specific causal agent.
2. It may be characterised by specific lesions.

It is obvious that our conception of any disease will differ according to which of these criteria we adopt as our basis of specificity.

For example, let us consider tuberculosis. If we adopt the anatomical basis—that is, if we consider tuberculosis to be a disease characterised by the presence of certain definite anatomical lesions called tubercles, we must include in this term, not only those cases in which the tubercle bacillus is the causal agent, but also other cases due to other organisms such as aspergillus, and we must exclude all diseases not characterised by these lesions. If, on the other hand, we adopt the etiological basis, we must exclude all cases which are due to organisms other than the tubercle bacillus, and we must include cases in which the anatomical tubercles are not present; for the tubercle bacillus occasionally gives rise to a diffuse instead of a nodular inflammation.

Thus the term "tuberculosis," which originally had a purely anatomical significance, has now come to have a meaning which is purely etiological, and in consequence those diseases which are characterised by anatomical tubercles, but which are not due to the tubercle bacillus, have been designated by the unsatisfactory term "pseudo-tuberculosis."

The same change is now, unfortunately, taking place in the meaning of the term "pneumonia." This originally meant a disease characterised by the anatomical condition of inflammatory consolidation of the lung. Now, however, there is a growing tendency to include pneumonia among the specific infective diseases, that is, among the diseases due to a specific micro-organism. If this is done we shall have to describe pneumonia, not as a disease characterised by an inflammation of the lung, but as a disease which is the result of infection by the diplococcus pneumoniae, and we shall have to include in it certain cases of broncho-pneumonia, pericarditis, meningitis, otitis, empyema, etc., any of which conditions may be caused by the diplococcus with or without the characteristic inflammation of the lung. Judging from the analogy of tuberculosis we may expect in the future to read about "pneumonia" of the meninges, ear, pericardium, etc., and we may also find the term "pseudo-pneumonia" introduced to describe those diseases characterised by inflammation of the lung, but not due to the diplococcus pneumoniae.

It is thus a great mistake to give an etiological significance to a term the significance of which is purely anatomical. It is sure to lead to great confusion in medical nomenclature. With regard to this question of specificity I should like to lay down the following two complementary laws.

1. No specific causal agent always gives rise to characteristic lesions.
2. No specific lesion is always produced by a single definite causal agent.

Besides the examples given above, I will give the following:

1. The diphtheria bacillus usually gives rise to a fibrinous inflammation of the fauces or larynx with the production of membrane. Sometimes, however, the membrane is wanting, and instead of a fibrinous there is a gangrenous or oedematous inflammation. On the other hand, fibrinous inflammation of

the fauces may be due to other organisms than the diphtheria bacillus.

2. Congenital syphilis, if it affects the liver, usually gives rise to a pericellular cirrhosis. It may, however, give rise to other forms of cirrhosis, or to gummata. On the other hand, pericellular cirrhosis of the liver is occasionally found in patients presenting no other evidences whatever of congenital syphilis.

3. The streptococcus pyogenes in the subcutaneous tissue may give rise either to a suppurative or a non-suppurative inflammation, and suppuration may be produced by organisms other than the streptococcus.

4. Chronic alcohol poisoning, if it affect the liver, usually gives rise to a multilobular cirrhosis. It may, however, give rise to other forms of cirrhosis, or to fatty degeneration without cirrhosis. On the other hand, multilobular cirrhosis may be due to other causes than alcohol poisoning.

These examples might be multiplied indefinitely, but I have given a sufficient number to demonstrate the general truth of the two laws given above.

So far I have considered only anatomical lesions. It remains to consider those lesions which may be termed functional, such as alterations in secretion or motor activity. At first sight it appears doubtful whether we can apply the laws to specific alterations in functional activity, but, on closer investigation, it will be seen that, although there are apparent exceptions, yet there are numerous cases in which the laws hold. At any rate, the second law would appear to hold good in all cases.

5. Morphine, while usually producing narcosis, sometimes produces excitement. On the other hand, narcosis is produced by many agents besides morphine.

6. Lead poisoning, if it affect the central nervous system, may give rise to either excitement or dulness.

As an apparent exception to the first law may be mentioned the action of tetanus toxin on the central nervous system. It is quite possible, however, that the tetanus toxin may sometimes reach the central nervous system without producing the characteristic convulsions, since an examination for the tetanus bacillus is only made when the characteristic convulsions are present.

We have now to consider the following question, Why does the effect produced by a specific causal agent vary in different cases? The answer is to be found in the fact that no disease is dependent on one cause alone. The factors concerned in the production of a disease can be divided into (1) the essential factor, and (2) the adjuvant factors; the essential factor being that without which the disease could not occur, and the adjuvant factors being those which assist in production of the disease, but which in themselves are not individually essential.

In those diseases which are due to a specific causal agent, this agent itself is the essential factor, the adjuvant factors being those usually known as the predisposing causes. It will be found that, granting the access of the causal agent to the body, the adjuvant factors can be reduced to two in number: (1) The locality to which the causal agent is applied; (2) the relative virulence (or dose) of the causal agent. By "relative virulence" I mean the virulence of the causal agent considered in relation to the susceptibility of the patient.

1.—The Locality to which the Causal Agent is applied.

- (a) An irritant applied to an epithelial surface usually gives rise to a catarrhal inflammation—for example, inflammations of the lung due to inhalation, or to the spreading of an inflammation along the air passages, usually have the characters of broncho-pneumonia.

- (b) An irritant applied to a surface covered by endothelium usually sets up a fibrinous inflammation, such as is seen in pericarditis, etc.

- (c) An irritant acting on connective tissue tends to produce a suppurative or an indurative inflammation, etc.

2.—The Relative Virulence (or Dose) of the Causal Agent.

The intensity of any disease varies with the virulence of the causal agent and with the susceptibility of the patient. Thus an irritant of low relative virulence acting on connective tissue gives rise to a productive inflammation, if the virulence is greater the result is suppuration, and if the relative viru-

* Communicated to the Leeds and West Riding Medico-Chirurgical Society, November 16th, 1900.

lence is very great the effect is a serous inflammation leading to spreading oedema.

Conclusions.

From these facts I draw the following conclusions:

1. If a disease is due to a specific causal agent it is not characterised by specific anatomical or functional lesions.
2. If a disease is characterised by specific lesions we must not expect to find a specific causal agent. From this it follows that from the characters, histological or macroscopical, of the lesions we cannot positively affirm that the disease is due to a specific causal agent. The only means by which we can affirm this is to find and isolate the causal agent itself.
3. Variations in the characters of the lesions in diseases due to specific causal agents depend not on the nature of the agent, but on the locality to which it is applied and on its relative virulence.

MYOKYMIA, OR PERSISTENT MUSCULAR QUIVERING.

By R. T. WILLIAMSON, M.D.LOND., F.R.C.P.,

Physician to the Ancoats Hospital, Manchester; and Assistant Lecturer on Medicine, Owens College.

A SHORT time ago a patient came under my care at the Ancoats Hospital, suffering from a peculiar persistent quivering of the muscles of the limbs and trunk. The symptoms corresponded to those met with in the very rare affection to which Professor Schultze has given the name of "myokymie" ($\mu\upsilon\varsigma$, a muscle, and $\kappa\upsilon\mu\alpha$, a wave), and Dr. Kny (who had previously reported two similar cases) that of "myoclonus fibrillaris multiplex." I have only been able to find records of eight cases in medical literature, but probably others have been reported.

The following are the notes of my case:

J. C., aged 21, clerk, was admitted as an in-patient at the Ancoats Hospital, August 29th, 1900, suffering from a peculiar persistent quivering of the muscles of the limbs, trunk, and face.

Previous History.

The left leg had been suddenly paralysed in infancy (probably owing to acute anterior poliomyelitis), and had remained useless. At the age of 16 years the left (paralysed) leg had been seriously injured, through an accident, and amputation had been necessary at the upper part of the left thigh.

Two years ago (at the age of 19) he had first noticed quivering of the muscles of the right leg. The onset was gradual, and the quivering spread in course of time to the muscles of the arms, trunk, face, and tongue. There had been no cessation or diminution of the symptom since its onset. After the amputation of the left leg he had been able to get about with the aid of a crutch, but recently he had found great difficulty in doing so, owing to the right leg frequently "giving way" through the muscular quivering.

He had always been a teetotaler, but had taken strong tea with each meal for many years. Three times a day for many years he had taken two cups (about one pint) of strong tea. He had only smoked about an ounce and a-half of tobacco weekly.

Condition on Examination.

The patient was fairly well nourished, and there was no definite wasting of any of the muscles of the arms, right leg, or trunk. There was a well-marked lateral curvature of the spine, which had gradually developed since the left leg had been amputated. The most striking symptom was a persistent rapid quivering of the muscles of the right leg, the arms, trunk, face, and tongue. Sometimes one small bundle of muscle fibres contracted rapidly, giving the appearance of fibrillary contraction; sometimes several bundles of muscle fibres contracted; at other times the whole muscle would contract rapidly. The symptom was best seen in the leg, and the quivering affected chiefly the thigh and calf muscles. At one moment a bundle or several bundles of muscle fibres would contract rapidly; directly afterwards a wave of contraction would pass over the whole muscle. This muscular quivering produced a very slight shaking or trembling of the whole leg, even when the patient was in bed. When the quivering was very violent, rapid and repeated slight flexion movements of the foot occurred at the ankle, owing to the quivering of the calf muscles. The arms were affected in a similar manner but not quite so markedly as the leg, the muscles of the shoulder, upper arm, and forearm being most affected, those of the hand very slightly. In the arms the quivering or contraction did not affect the whole of a muscle so frequent as in the leg. The rapid and repeated quivering of the muscles of the arms and leg continued more or less all day long; the muscles were affected in a most irregular manner, and the frequency of the quivering varied considerably.

The muscles of the chest and abdomen presented occasional fibrillary contractions. The muscles of the face and tongue, the masseters and neck muscles presented frequent fibrillary contractions. The eyes were not affected; there was no nystagmus. The muscular quivering and contractions in the limbs and trunk continued when the patient was at rest in bed, as well as when he was going about. When the arms were held out in front of the chest, the muscular quivering produced a slight tremulous movement of the whole limb resembling that so frequently

seen in Graves's disease; but there was not the flexion and extension movements of the fingers or wrist which are seen in paralysis agitans. The muscular quivering and slight tremor of the limbs were very little affected by voluntary movement; they were not diminished thereby; if any change occurred at all, they were slightly increased. The muscular quivering was increased by excitement, and when the patient was undergoing examination. The patient was able to write fairly well, and though the writing was slightly jerky, it was much better than in the various other forms of tremor, with the exception of paralysis agitans. He was also able to draw a straight line on paper fairly well.

The patient could perform all the movements of the arms and legs, and there was no localised wasting or paresis of muscles. There was no rigidity of the limbs. The knee-jerk was a little increased. There was no ankle clonus. On attempting to obtain ankle clonus occasionally a few jerks were obtained, but at other times this symptom was entirely absent. The plantar reflex was normal (flexor type). There was no loss of sensation to touch, pain, or temperature. The sensory cranial nerves were unaffected. There were no mental symptoms. The patient was not emotional or nervous. There was no affection of speech. The patient often complained of palpitation of the heart, and the pulse was often quick, 84 to 108 per minute. Examination of the heart, lungs, and abdomen revealed no evidence of organic disease. The patient complained of frequent sweating. There was no exophthalmos, no Graefe's symptom, and no thyroid enlargement. There was a little nausea after food; the bowels were regular.

The urine was often pale, at other times of normal colour. The quantity was increased; it was measured for sixteen days, and found to vary in amount from 64 to 124 ounces daily. The specific gravity varied considerably; when the urine was pale it was often low—1005 to 1008; at other times it was 1012 to 1015. Albumen and sugar were absent. The reaction was sometimes alkaline, sometimes acid. The temperature was normal. Insomnia was a frequent symptom.

As the patient had taken strong tea at every meal for many years I allowed him no fluids except milk and water for two weeks, but without any result as regards his symptoms. During the four weeks he was in the hospital there was no definite change; some days the muscular quivering was a little less, but the diminution was never permanent. The muscular quivering continued more or less all day long, just as it had done for two years.

The symptoms in the case just recorded corresponded to those in the affection to which the name of "myokymia" has been given. Similar cases have been recorded by Kny, Schultze, Hoffmann, Biancone, Buber, Karcher, and Mayer. In the 8 cases of which I have been able to find records in medical literature, the ages were from 21 to 71. All the patients were males. Five of the eight were peasants or country labourers. One patient was a painter, another a lead solderer; both of these patients suffered also from symptoms of lead poisoning. One patient suffered from chronic double sciatica. In all of the 8 cases the legs were most affected. In 2 cases the legs only were affected; in 1 the legs and muscles of the back; in 2 the arms and legs; in 1 the legs and arms but not the hands; in 1 almost all the muscles of the body; in 1 almost all the muscles except those of the face and neck. [In the case which I have just reported, the muscles of the limbs and trunk, as well as those of the neck and face, were affected.] In 4 cases recovery occurred in the course of a few months; in 1 case there was distinct improvement. The treatment, in the cases in which recovery occurred, was by warm baths and galvanism (Kny), by bipolar faradic baths (Kny), by rest in bed and warm baths (Schultze). Buber's case improved distinctly under treatment with the galvanic current and warm baths. Biancone recommends complete rest in bed, warm baths, and sodium salicylate.

The cause and pathology of this peculiar affection are at present unknown.

REFERENCES.

- Kny, *Archiv f. Psych. u. Nervenkrankheiten*, Bd. xix. Schultze, *Deutsche Zeitschrift f. Nervenheilkunde*, 1894, vi (abstract), *Neurologisches Centralblatt*, No. 6, 1895. Hoffmann, *Neurologisches Centralblatt*, No. 6, 1895. Mayer, *ibid.*, No. 14, 1897 (abstract). Buber, *ibid.*, No. 15, 1897. Biancone, *ibid.*, No. 4, 1899 (abstract). Karcher, *ibid.*, No. 7, 1899 (abstract).

A BRIEF ACCOUNT OF THE RECENT EPIDEMIC OF CHOLERA IN KASHMIR.

By ERNEST F. NEVE, M.D., F.R.C.S.E.,
Kashmir.

ONCE more Kashmir has been visited by a sharp epidemic of cholera. Thus, in 13 years there have been three invasions. In 1888 there were probably 18,000 attacks, and a total mortality of nearly 10,000; in 1892 the official returns showed 16,845 cases, with 11,712 deaths; owing to the concealment then extensively practised by the people, it is probable that