severely, and she was then in one which had lasted over an hour. Morphine gr. $\frac{1}{2}$, with atrop. sulph. gr. $\frac{1}{10}$, was injected, soon after which the fits ceased. As the apærent had not acted, enemata were administered without effect. She remained free from fits till the evening, when they again returned with great severity; the os was then slowly dilating, and the pains coming on regularly, but with little force. Another injection of morphine gr. $\frac{1}{2}$ caused a cessation of the fits, which did not return. On the succeeding morning morphine gr. $\frac{1}{2}$ was injected, and the labour was found to be progressing slowly. During these two days she remained in a semi-conscious condition, not recognising anyone, though capable of feeling pain. On the evening of the second day, as the os was well dilated, and there was uterine inertia, chloroform was administered, and a dead child was delivered with forceps. The mother was in a restless and sleepless condition for four days afterwards; this was relieved by injections of morphine gr. $\frac{1}{2}$. The urine, which in the early stage of the illness contained one-eighth albumen, was free from it on the fifth day. The bowels were at last moved by repeated doses of mist. sennæ co.

These cases suggest the conclusion that to induce labour, however severe the fits, is a great mistake, the two cases so treated terminating fatally, whereas those in which no operative procedures were carried out with a view to remove a cause of the fits recovered perfectly. In both cases the fits were almost at once checked by the administration of morphine, and did not return so long as the patient was under its influence. Another advantage is that morphine is free from the depressing effect upon the heart exerted by chloroform or chloral. The old idea that it exercised a bad effect in cases of renal trouble would appear to have no foundation in fact.

SYPHILIS IN THE ROYAL NAVY.

By J. P. H. GREENHALGH, M.B., CH.B.VICT.

Surgeon, R.N.

COMPARATIVELY recently it was pointed out in the BRITISH MEDICAL JOURNAL that the decrease in the late venereal returns of the Indian Army could be accounted for on two grounds, these being (1) the inclusion of syphilis under the Cantonments Act for infectious diseases; (2) more efficient treatment. With reference to this question of treatment the following significant passage occurs:

We had occasion some 18 months ago to call attention to the fact that in very many instances soldiers were only treated for syphilis as long as there was some outward sign of the disease, and that no regular and long-continued treatment was carried out. After the secondary symptoms had disappeared temporarily, and the man was discharged from hospital, only too often no further medicine was taken until the disease broke out again, probably in a more virulent form.

This is a serious indictment, touching as it does our professional responsibility to those under our charge, which, whether in the public service or out of it, I take to be our first obligation. It is no less a matter of concern to have to admit that in the Royal Navy at any rate, this allegation contains a present-day element of truth. Two examples will suffice to show its applicability:

CASE I.—A. B., aged 23, stoker, contracted syphilis in March, 1899. He was sent to hospital early in July of the same year, where he remained for a period of seven weeks. In the following January he developed late secondaries; mucous patches and small ulcers were present on the lips, palate, and tonsils, and staining of the skin indicated the antecedent occurrence of a rash. He was again sent into hospital in February.

CASE II.—C. D., aged 20, stoker, contracted syphilis in the middle of January, 1900. He was returned from hospital about the end of March; three weeks later he reappeared for treatment. He had then a large indurated sore occupying the same position as the original chancre, and a roseolar rash over the trunk, characteristically syphilitic.

Reading between the lines, it seems that the first case was treated until secondary symptoms had disappeared, the second until the primary lesion had subsided; both men were then discharged to duty.

The fault does not lie at the doors of the hospital staff. It is obviously impracticable for men to be retained in hospital for periods of from twelve to eighteen months in order to undergo a continuous course of treatment. And it would be ridiculous to suggest that medical men in the navy do not recognise the imperative necessity for prolonged mercurialisation in syphilis equally with their professional brethren in other spheres of practice. The evil is rather the outcome of a faulty nomenclature.

Any case of venereal disease is, in accordance with the nosological requirements at the Admiralty, returned as primary syphilis. Whether the lesion be clearly one of Hunterian chancreoid infection or one of local contagious and non-infecting ulceration, the designation is the same, and here we have the crux of the whole question. A man is sent into hospital with venereal disease, and is discharged after a

few weeks apparently well, with "primary syphilis" entered on his medical history sheet. Should treatment be carried on?

It is not to be denied that many difficulties at present bar the way to a satisfactory answer. It may be argued that where a differential diagnosis has been established before the patient is forwarded to hospital, it is incumbent on the medical officer primarily responsible to continue or discontinue treatment when the man returns under his care, as the case requires. Where, however, the doctor or the patient is shortly transferred elsewhere, other arrangements are necessary. In such an event a supplementary note on the history sheet would clear up the ambiguity of the official entry, and the purpose would still be served. Such an innovation, however, the effectiveness of which depends on its universal application, could only be introduced through the proper channels, and even if this were not so, is it satisfactory that individual expedients should be a substitute for official action? To the medical mind the term primary syphilis is specifically limited to the true infective process, and that it should include other conditions which *in ipso* are pathologically distinct is as confusing as it is unscientific. Once the wide-reaching difference between Hunterian syphilis and non-infecting venereal ulceration is recognised, and its importance made obvious on the medical history sheet, the way for thorough treatment, which means diminished overcrowding of hospitals by cases of secondary syphilis, improved statistics, and a corresponding increase in the effective strength of the navy is made easy. In every depôt and on every ship a time list of men under treatment would be kept; in the case of men newly joined a brief inspection of their history sheets, coupled with the certain knowledge that where specific disease existed treatment had been regularly carried out up to date, would give all the information necessary to the completion of a prophylactic circle embracing the entire navy, and to the carrying on of an adequate campaign against this formidable and insidious disease.

NOTE ON A CASE OF PROTRACTED STUPOR OR TRANCE,

TREATED IN THE ROYAL INFIRMARY, NEWCASTLE-UPON-TYNE.

By DAVID DRUMMOND, M.D.,
Senior Physician to the Hospital.

D. B., a clerk, aged 26, was admitted into the Royal Infirmary, Newcastle-upon-Tyne, on March 31st, 1900, in a state usually described as lethargy, trance, or stupor.

History.

He was always more or less melancholic and reserved, and religious to the verge of fanaticism. During the previous two years he became less and less communicative and increasingly taciturn, and for twelve months prior to his admission he seldom spoke unless addressed, lost his appetite, and became thin and pale, but attended regularly to his employment until seven weeks before his admission, when he voluntarily retired into private life, and spent his time reading the Bible and taking solitary walks. A little later he declined to leave his bed, and refused food, and towards the end of March his illness culminated in a "fit" (?), when he passed into the state to be presently described.

Family History.

A neurotic family history was elicited. His grandfather died in an asylum and his brother committed suicide, whilst one brother died of phthisis during the patient's stay in the infirmary.

Condition on Admission.

He was a tall, spare, sickly, and anæmic young man. There was no history of sexual irregularities. At the time of admission to hospital it was stated by friends who brought him that he had not moved or spoken for a week. He lay motionless and took no notice of his surroundings. His eyes were open and appeared to stare in front of him with a far-away look that suggested subconsciousness. His limbs were cold, flaccid, and unresisting. The tongue was coated and the breath foul, and nutrition much lowered. The reflexes, superficial and deep, were just perceptible, but the corneal reflex