INTRAPERITONEAL RUPTURE OF THE BLADDER.

OPERATION FOUR DAYS AFTER INJURY: RECOVERY.

By W. PERCY BLUMER, F.R.C.S. EDIN.,
Surgeon to the Sunderland Infirmary.

J. P. B., labourer, single, aged 35, was admitted into the Sunderland Infirmary at 1.30 P.M. on August 2nd, 1900, complaining of pain in the abdomen and inability to pass urine.

History.

On the evening of July 29th, while walking along a street, he was seized with a feeling of giddiness and fell, striking his abdomen on the kerbstone. On recovery, he felt little the worse for his accident, and walked home. On reaching home he discovered that all attempts at micturition were fruitless; there was, however, a slight escape of what he took to be blood. On July 30th he stayed in bed and sent for a doctor, who prescribed a pill and saw him only once. He remained in bed till August 2nd, and feeling no better, walked to the infirmary, a distance of over two miles. He had been drinking heavily just before his accident, and said his bladder was full at the time.

Condition on Examination.

The abdomen was greatly distended all over, somewhat tender on palpation, and absolutely dull in both flanks and across the abdomen to a little above the umbilicus. There was no extravasation into the perineum. Hiccough was constant and distressing; this had been the case for the three days prior to admission. He appeared very ill; the face was pale and drawn, with sunken eyes. No food had been taken, except milk and water, since the accident. Pulse 60 per minute, regular and fairly full. Temperature 96° F.

A soft rubber catheter passed easily, and immediately bloody urine flowed, 196 ounces in all being drawn out (quantity carefully measured). The abdomen thereafter resumed a more normal appearance. There was no doubt as to the nature of the injury.

Operation.

Laparotomy was performed at 4 P.M., with the assistance of Mr. T. F. Hopgood. The patient having been put under chloroform an incision was made in the middle line just above the pubis, and the abdominal cavity entered in the usual way. A quantity of fluid similar in character to that previously drawn off flowed out. The bladder was lightly contracted, and showed a vertical tear on the posterior wall, which admitted two fingers easily, the peritoneum covering its wall being torn an inch lower down. There was no peritonitis.

The rent was closed with continuous sutures, the bladder tested, the abdominal cavity flushed out with hot boracic solution and the abdominal wound stitched up, a Keith's drainage tube being inserted, and a soft catheter tied into the bladder.

After-History.

The tube was exhausted one hour afterwards (fluid $\frac{1}{2}$ ounce), then again in two hours (fluid 1 ounce). Thereafter exhaustion was made every four hours. The tube was withdrawn at the end of forty-eight hours, and the opening closed with a flying suture.

On the evening of August 5th the patient became very noisy and restless, attempting to get out of bed, pulling out the catheter and threatening the nurses; he suffered also from slight hallucinations. A hypodermic injection of morphine, $\frac{1}{4}$ grain, followed by $\frac{1}{2}$ grain one hour later, had the effect of quieting him, and that night he slept better. On August 6th he was quiet, and slept seven hours.

On August 7th, when dressing the wound, the lower stitches appeared very tense, and on cutting two a drachm or so of pus, and what was evidently urine, escaped. The opening was syringed with 1 in 5,000 corrosive solution and drained with iodoform gauze. That evening a consultation was held as to the advisability of opening up the abdominal wound; it was, however, deemed inexpedient on account of the sloughing state of the tissues and the almost entire absence of symptoms of peritonitis. A fine rubber drainage tube was in-

serted instead of the iodoform gauze, and hot boracic fomentations applied every two hours. A seidlitz powder was administered.

There was evidently a little leakage from the bladder, and small quantities of urine continued to flow through the tube in the abdominal wall, the channel being cut off from the general peritoneal cavity.

The remaining stitches were taken out on the tenth day, when the whole length of the wound burst open, and a large slough was taken out. The urine was now very alkaline. On August 9th, 99 ounces of urine were secreted in the twenty-four hours. The bladder was syringed out with boracic lotion, and boracic acid, 10 grains, was given thrice daily. This treatment was continued for ten days, after which salol was administered in 5-grain doses every two hours, in place of boracic acid, up to 60 grains in the day. Under this treatment the cystitis rapidly yielded.

The temperature was generally subnormal for the first ten days, but on the third and tenth days it reached 99° F. On the sixteenth and twenty-first days it reached 101° F. It then fell to normal, and remained so until the end.

The fistula was rather slow in healing, but the patient was discharged quite well on October 29th. During the last week of his convalescence he had three epileptic fits of a transitory nature.

Remarks.

I have not been able to find on record such a case. The points calling for special mention are:

1. The length of time between the injury and operation.
2. The large amount of fluid in the abdomen causing comparatively little trouble.
3. That four days after the injury the patient walked over two miles.
4. The absence of symptoms of peritonitis.

I am indebted to Dr. C. G. Wotherspoon, house-surgeon, for the notes of this case. His care and attention materially contributed to the patient's recovery.

NOTES ON A CASE OF PROFUSE HYDROPERITONEUM COMPLICATING UTERINE FIBROIDS.

By A. C. BUTLER-SMYTHE, F.R.C.S.E.,

Surgeon to the Samaritan Free Hospital for Women and Children, and to the Grosvenor Hospital for Women and Children, Westminster.

In abdominal surgery one frequently meets with cases in which uterine fibroids are accompanied by varying quantities of hydroperitoneum, the amount seeming to depend more on the mobility of the growth than on its size. Instances of profuse hydroperitoneum occurring in patients free from thoracic or other abdominal disease are, however, uncommon, and therefore I have ventured to publish this case.

Mrs. T., aged 40, consulted me in December, 1895, on account of an abdominal swelling.

History.—She had been married seventeen years and had had three children, the youngest being then 14 years of age. Menstruation began at 15, and her periods had always been regular and painless, the loss being free and the flow lasting usually seven days. Her last period had occurred in November, 1895. In her early life she had had the usual minor ailments, and after marriage pneumonia. She had also two attacks of influenza. Six months previous to her visit to me she experienced some abdominal pain, and later on a swelling was noticed in her right side. This swelling seemed to become bigger before each period, and in the intervals it decreased in size. Of late she suffered from constipation, and there were symptoms of bladder pressure. There had been no loss of flesh within the last six months, her weight being now 8 st. 11 lbs. Up to December she had enjoyed good health and was well able to attend to her business and household duties.

Condition when First Seen.—Medium height, pallid features, skin moist, tongue clean, pulse regular and of fair volume, temperature normal, respirations natural, slight hæmic murmur at base, but otherwise the heart seems healthy, bowels constipated, urine free from albumen and sugar, liver normal. Bimanual examination revealed two movable tumours in the abdomen, one situated in the right iliac and lumbar regions, the other occupying the hypogastric and left iliac regions. Pressure on one or both of the tumours was communicated to the uterus, showing them to be closely connected with that organ. The cervix uteri was hard and granular, the os patulous, and the uterine cavity measured $3\frac{1}{2}$ inches. The diagnosis arrived at was 'subperitoneal uterine fibroids,' and it was agreed that operative interference was uncalled for in the absence of severe symptoms. The patient was therefore put upon ergot and iron, and advised to live frugally, avoiding wines and stimulating diet.

Progress.—The patient remained in fair health until August, 1896, when she had an attack of peritonitis, and in my absence from town consulted Sir Spencer Wells, and later on Mr. Alban Doran, both of whom advised against operative measures. During the following ten months she had little to complain of, and continued at her business, being able to take long walks and look after her house. On June 28th, however, she met with an accident, being knocked down by a cart when crossing the road. She was much bruised on the abdomen, thighs, legs, and arms, and narrowly escaped being killed. The accident was followed by abdominal pain, and from that date she rapidly lost ground.

Repeated Aspirations.—The tumours became tender, emaciation was marked, and on August 24th free fluid was detected in the abdomen. By September 13th she was much distended, and 18 pints of clear amber-coloured fluid were removed by aspiration. On September 24th she was again relieved, 20 pints being withdrawn. At the end of two weeks, on October 6th, the aspirator was again necessary, and about 20 pints were evacuated. On October 25th the patient was admitted into the Grosvenor Hospital for Women, Westminster, and two days later she was for the fourth time aspirated, 21 pints being removed. The reaccumulation of fluid, however, was so rapid that on November 2nd about 14 pints were taken away.

Operation.—The patient now decided to have the tumours removed. Accordingly on November 6th I operated. On opening the abdomen about a gallon and a half of fluid escaped. The parietal peritoneum was unusually thick and injected. The omentum was adherent to the parietes, and also to the anterior surfaces of both tumours. There was much intestinal adhesion, the caecum and appendix caeci having to be dissected off the lower part of the right tumour before it could be brought outside the abdomen. The *serre-noud* was then placed in position, and the operation completed. Chiefly owing to the extent of intestinal adhesions the operation occupied two hours, and the patient when put to bed was in a very collapsed condition.

Recovery.—She rallied steadily, and made a good recovery. There has been no return of the hydroperitoneum, and at the present time the patient enjoys the best of health.

To my mind it is clear that the hydroperitoneum was directly caused by the injury to the peritoneum covering the fibroids, inasmuch as up to the date of the accident the patient was in very fair health, whereas all the bad symptoms rapidly developed after the injury, and were at once relieved, and the patient cured by the removal of the uterus and tumours. The following points seem to me to be worthy of consideration, and may be of interest to my hearers:

1. The ovaries and tubes were healthy.
2. The tumours removed were simple fibromyomata, and showed no signs of malignant or other degeneration, but the peritoneum covering them was thick and injected. The tumours were not oedematous.
3. Up to the date of the accident no free fluid had been detected, and there was no loss of weight.
4. Emaciation was first noticed, and free fluid in the abdomen first detected, about six weeks after the accident.
5. So far as could be ascertained the patient was free from thoracic or abdominal disease.
6. Though aspiration had to be performed five times in six weeks, and about 20 pints of fluid were evacuated each time, yet there was no reaccumulation of fluid after the supravaginal hysterectomy, and the patient remains in perfect health (July, 1900).

OTOMYCOSIS IN THE TROPICS.

By H. CAMPBELL HIGGETT, C.M., M.D.,

Physician to the Royal Palace, Bangkok.

OTOMYCOSIS, or the growth of fungus in the external auditory meatus, is apparently a rare disease in temperate climates if one may judge from the scanty literature on the subject. During eight years' work in Singapore and Bangkok, however, my experience has been that it is quite a common—if not the most common—disease affecting the external auditory meatus which one meets with in the tropics. The reason for this appears to be that all the conditions favourable to the growth of fungi are present—namely, heat, moisture, and the presence practically everywhere in abundance of the spores of the fungi. All my cases were in adult Europeans. Children are said to be exempt from the disease.

Symptoms.—These vary in intensity according to the severity of the case. There may be simply a sensation of blocking of the ear with slight impairment of hearing. Itching, pain, and often a considerable amount of watery discharge, is complained of, and the patient notices that his pillow has been stained during the night by a yellowish fluid which he finds exuding from one or both ears. Both ears are usually affected, but often in different degree. When an attack of acute diffuse inflammation of the meatus supervenes, as is unfortunately a

frequent complication, great pain and sleeplessness follow, and often a considerable rise of temperature is noted for a few days.

Physical Appearances.—In typical first attacks the meatus is seen to be filled up with a soft wool-like substance which varies in colour according to the type of fungus present. As I have seen it it is usually of a pale lemon yellow or a very pale yellowish green. On passing a probe into the canal the obstruction is found to be soft and moist and is readily removed. The walls of the canal may be found to present quite a normal appearance, but as a rule they are somewhat reddened, and the irritation may have gone as far as to have led to some desquamation of the epithelium. In cases in which the attention of the patient has been suddenly drawn to his ears by an acute attack of pain and deafness, it will be found that there are signs of extension of the catarrh to the middle ear. In chronic cases, besides the soft masses of recent fungus, impacted masses of a material like sodden newspaper are found in the meatus. These come away in large scales, or even casts of the canal, and the walls are found to have undergone thickening, so much so as in some cases to hide the tympanic membrane. Microscopical examination of the fungus reveals the presence of such common fungi as the *penicillium glaucum*, *aspergillus*, and more commonly in my experience the *mucor mucedo*.

The complications noted in my cases were diffuse inflammation of the external auditory meatus, acute serous catarrh of the middle ear, chronic catarrh of the same, perforation of the membrana tympani, swelling and even abscess of the lymphatic glands at the angle of the jaw, and eczema of the meatus and pinna. The acute adenitis, the result of septic infection from the ears, was so severe in one case as to require the administration of chloroform on two separate occasions in order to abscesses on either side of the neck.

Diagnosis is usually easy, and is at once settled by the microscope.

Prognosis.—This is a readily curable disease, but is only so if the case be taken charge of by the physician himself. To hand over the details of treatment to the patient is a plan which leads to much trouble and disappointment. I have known of cases going on for weeks and even months when carelessly attended to.

Treatment.—As much of the fungus as possible should be removed with a cotton and probe, and then the canals should be well syringed with a warm solution of bichloride of mercury (1 in 5,000). The canal is then thoroughly dried with cotton wool, so as to get rid of all water, which favours the growth of the fungi, and finally it is sponged with a solution of bichloride of mercury in absolute alcohol (1 in 1,000). This gives rise to considerable pain for a moment, but it soon passes off. The alcohol is allowed to evaporate, and then the meatus is plugged with sterilised cotton-wool, which is not removed until the following day. It will then be found that many of the spores that had resisted the action of the germicide applied the previous day have germinated in the interval, and there is apparently as much fungus as before. It is not so dense, however, and is more easily removed by the same means. This process is repeated daily until no more fungus forms, then the eczema or other complication is treated in the usual way. After the fungus has ceased to grow I often insufflate a powder of boric acid 2 parts, bismuth salicylate 1 part, and oxide of zinc 3 parts. Such a combination completes the cure not only of the otomycosis, but also of the eczema of the meatus.

Prophylaxis consists in keeping the ear canals very clean and dry, and especially in avoiding the entrance of sea water, which by reason of its deliquescent salts apparently conduces more than fresh water to the growth of the fungi.

AMBULANCE WORK.—At Wallsend, recently, 29 certificates of the St. John Ambulance Association were presented to members of the ambulance corps in connection with Messrs. Swan and Hunter's shipyard on the Tyne, and subsequently Dr. Robert Alfred Wilson, honorary surgeon to the corps, was presented with a silver mounted umbrella.

At a recent meeting of the Stockport Infirmary a resolution was passed expressing the regret with which the resignation of Dr. Keay—who had been resident medical officer for four years, and since 1887 honorary surgeon—was received.