Suppression In Vivo of Human Papillomavirus Type 18 E6-E7 Gene Expression in Nontumorigenic HeLa × Fibroblast Hybrid Cells

FRANZ X. BOSCH,^{1†*} ELISABETH SCHWARZ,² PETRA BOUKAMP,³ NORBERT E. FUSENIG,³
DUSAN BARTSCH,² AND HARALD ZUR HAUSEN²

*Institut für Zell- und Tumorbologie,¹ Institut für Virusforschung, Projektgruppe Angewandte Tumorstudiologie,²
and Institut für Biochemie,³ Deutsches Krebsforschungszentrum, Im Neuenheimerfeld 280,
D-6900 Heidelberg, Federal Republic of Germany*

Received 14 March 1990/Accepted 21 June 1990

The E6 and E7 genes of the cancer-associated human papillomavirus (HPV) types 16 (HPV16) and 18 (HPV18) can induce cell immortalization in vitro in normal human keratinocytes. This, however, is not associated with tumorigenicity in vivo. On the other hand, tumorigenicity of HPV18-positive HeLa cervical carcinoma cells can be suppressed by fusion of HeLa cells with normal human keratinocytes or fibroblasts. We have addressed the question of whether suppression of tumorigenicity in HeLa × fibroblast hybrid cells might be due to a reduced ability of these cells to express the HPV18 E6-E7 genes in vivo. Nontumorigenic hybrid cells and tumorigenic hybrid segregants were transplanted as organotypical cultures or injected subcutaneously into immunocompromised mice and were analyzed for HPV18 E6-E7 gene expression by RNA-RNA in situ hybridization. The tumorigenic hybrid cells showed a continuous and invasive growth that was associated with high levels of HPV18 E6-E7 mRNAs at all time points examined. In contrast, the nontumorigenic hybrid cells stopped cell proliferation approximately 3 days after transplantation. At this time they expressed the E6-E7 genes at low levels, whereas at day 2 high expression levels were observed. However, the mRNA levels of the cytoskeletal genes β -actin and vimentin remained high for at least 14 days, demonstrating that inhibition of growth and of HPV18 E6-E7 gene expression was not due to cell death. These results suggest that growth inhibition of the nontumorigenic HeLa × fibroblast hybrid cells in vivo might be caused by suppression of HPV18 E6-E7 gene expression and are compatible with the idea of an intracellular surveillance mechanism for HPV gene expression existing in nontumorigenic cells.

Cancer of the uterine cervix is regarded as a late consequence of persistent infections of the cervical epithelium with human papillomaviruses (HPVs) (54, 62). Of the more than 60 genotypes of HPVs identified so far, of which about one-third infect the anogenital tract of women and men (15, 42), only certain HPV types, in particular HPV types 16 (HPV16) and 18 (HPV18), are strongly associated with malignant genital tumors. Altogether, about 90% of cervical carcinomas are positive for HPV DNA; HPV16 represents the most prevalent HPV type, followed by HPV18. Additional experimental data strongly indicate that, besides the presence of HPV DNA in cervical epithelium, other events are required for full malignant conversion of cells infected by "oncogenic" HPVs. A change in the state of the viral DNA may represent one factor in progression toward malignancy. In premalignant lesions of the cervix (cervical intraepithelial neoplasias) the viral DNA is present as an episome, with a tendency toward integration in higher-grade cervical intraepithelial neoplasias (29). In cervical carcinomas either integrated or episomal HPV DNA or a mixture of both forms was detected (17, 31), whereas cell lines established from cervical carcinomas have been shown to harbor exclusively integrated HPV16 or HPV18 DNA. The integrated viral DNA has usually suffered a disruption in the E1 and E2 open reading frames, accompanied in many cases by deletion of parts of the open reading frame E2-to-L2 region (1, 9, 18, 44, 55). In all cell lines and the majority of carcinoma biopsies

analyzed, the viral open reading frames E6 and E7 are preserved and expressed as mRNAs and proteins (43, 46, 48). These data have indicated that expression of the HPV E6 and E7 genes may be crucial for maintenance of the transformed phenotype in vitro as well as of the malignant phenotype of cervical cancer cells and that DNA integration may represent a means for the E6-E7 genes to escape from regulation by viral control factors, such as the E2 *trans* repressor (5).

Several studies have provided evidence that the HPV16 and HPV18 E6 and E7 genes can profoundly influence proliferation and differentiation capacities of rodent and human cells in tissue culture (in vitro). It has been shown for the HPV18-positive cervical carcinoma cell line C4-I that proliferation of these cells in vitro depends on a certain level of E6-E7 gene expression (53). Transfection studies with rodent cells as targets have revealed that the E7 gene is able to induce immortalization of primary cells (3, 24), malignant transformation of established cells (4, 23, 28), and, in cooperation with an activated *ras* oncogene, malignant transformation of primary cells (12, 30, 35). After transfection of the DNA of the cancer-associated HPV types, normal human epithelial cells gain an indefinite life span in vitro (i.e., cell immortalization) and show altered differentiation in vitro (16, 25, 32, 36, 41, 58). Cell immortalization in vitro is brought about by the cooperative activities of the E6 and E7 genes (2, 21, 22, 33).

A common feature of all HPV-immortalized keratinocyte lines tested so far is that they do not form tumors in nude mice. Thus, the ability to express the HPV16 or HPV18 E6-E7 genes in vitro, which results in cell immortalization, is

* Corresponding author.

† Present address: Hals-Nasen-Ohren Klinik, Universität Heidelberg, Im Neuenheimerfeld 400, D-6900 Heidelberg, Federal Republic of Germany.

obviously not sufficient to confer tumorigenic growth potential to these cells *in vivo*.

The lack of tumorigenicity of these HPV-immortalized cell lines may reflect some early stages in HPV-linked carcinogenesis in which HPV infection is only one of several necessary events. This is also evident from the long latency period between primary HPV infection and cancer appearance (which is of low incidence) and the detection of HPV16 infections also in women with normal cervical cytology (62). In an attempt to explain these features at the cellular level, a model has been proposed that views the development of virus-linked human tumors as the result of a failing intracellular surveillance of persisting viral genes by putative cellular genes named cellular interfering factors (CIF genes) (60, 61). In the case of HPV-associated genital tumors, the CIF genes are postulated to be present as intact genes only in nontumorigenic cells and to act *in vivo* by suppression of HPV E6-E7 gene expression (61).

To examine the prediction that the HPV E6-E7 genes are negatively regulated in nontumorigenic cells *in vivo*, we used somatic cell hybrids that were established by fusion of tumorigenic HeLa cells with normal human fibroblasts (49, 51). These hybrid cells show the phenomenon of tumor suppression when inoculated into nude mice, i.e., they are nontumorigenic *in vivo*, although they exhibit all characteristics of transformed cells in tissue culture (*in vitro*). Rare segregants have been isolated that express the tumorigenic phenotype, a property that has been correlated with the loss of one copy of fibroblast chromosome 11 (52). We have used these hybrid cells, since HeLa cells are derived from an adenocarcinoma of the cervix and contain integrated and transcriptionally active HPV18 DNA (44). The HeLa-specific HPV18 DNA integration pattern is retained in the hybrid cells, and E6-E7 transcription is observed in the nontumorigenic hybrid cells as well as in tumorigenic segregants grown in tissue culture (38, 45). The nontumorigenic hybrid cells and their tumorigenic segregants are thus genetically closely related; they show the same transformed phenotype *in vitro* but differ drastically in their proliferation potential *in vivo* (nontumorigenic versus tumorigenic phenotype). We analyzed HPV18 E6-E7 gene expression by determination of mRNA levels in the nontumorigenic and tumorigenic hybrid cells *in vivo* after transplantation or subcutaneous inoculation into nude mice. In particular we wanted to examine whether the host-induced growth inhibition of the nontumorigenic hybrid cells is associated with a reduction in HPV18 E6-E7 mRNA levels. For this purpose we used RNA-RNA *in situ* hybridization on tissue sections, which allowed the determination of HPV18 mRNA levels in single cells in combination with a histological examination of the hybrid cell transplants.

MATERIALS AND METHODS

Somatic cell hybrids. The tumorigenic and nontumorigenic hybrid cells used were kindly provided by E. J. Stanbridge. After the cervical carcinoma line HeLa was fused with normal human fibroblasts, the nontumorigenic hybrid clone, designated 444, was obtained. From these cells a rare tumorigenic segregant clone, designated CGL3 (14, 49), originated. These cell clones are referred to here as nontumorigenic (clone 444) and tumorigenic (clone CGL3) cells, respectively. The cells were propagated in Dulbecco modified Eagle medium supplemented with 10% fetal calf serum. To generate organotypical cultures, 2×10^5 cells were seeded on glutaraldehyde-fixed type I collagen gels that had

been cast in silicon culture chambers as described in detail elsewhere (6; P. Boukamp, D. Breitkreutz, H.-J. Stark, and N. E. Fusenig, *Differentiation*, in press).

Transplantation of organotypical cultures. Approximately 17 h after seeding of the hybrid cells, the organotypical cultures were transplanted. At that time the cells had formed a complete monolayer. The medium was drained, and the organotypical cultures were covered with hatlike transplantation chambers. The whole unit was then placed onto the muscle fascia of pathogen-free nude mice (obtained from Wiga; for details see reference 59 and Boukamp et al., in press). At different times thereafter, the transplants were dissected *in toto*. They were snap frozen in isopentane precooled close to its freezing point in liquid nitrogen for cryostat sectioning and *in situ* localization analyses, or they were fixed in neutral buffered Formalin and embedded in paraffin for histological examination. In some experiments, the hybrid cells were injected subcutaneously. For this purpose, cells were harvested with trypsin-EDTA and suspended in 200 μ l of growth medium, 5×10^6 cells were injected subcutaneously.

RNA analysis. Total RNA to be analyzed by Northern (RNA) blot hybridization was extracted from nodules grown for different lengths of time after subcutaneous injection. The RNA was prepared by the guanidinium isothiocyanate procedure (8), separated by denaturing agarose gel electrophoresis, and transferred with capillary to Gene Screen Plus membranes. The probes used for the Northern blots were cDNA fragments specific for HPV18 E6-E7 or human β -actin and were labeled with 32 P by random priming (20). For *in situ* hybridization analyses, cRNA probes were used prepared by incorporation of either α - 32 P- or 35 S-labeled CTP or UTP into transcripts synthesized *in vitro* from cDNAs cloned into the Bluescribe vector (Promega Biotec Co., Madison, Wis.). In addition to E6-E7 and human β -actin, a cRNA probe specific for the human vimentin gene, which was kindly provided by H. Herrmann (Division of Membrane Biology and Biochemistry), was used. As a negative control, cRNAs transcribed in the orientation yielding sequences identical to the mRNAs in question were used (sense cRNA instead of antisense cRNA).

***In situ* hybridization.** A hybridization procedure based on the studies of Cox et al., (11) was carried out as described previously (7), except that the proteinase K treatment of the sections was shortened to 10 to 15 min.

Indirect immunofluorescence. The following murine monoclonal antibodies were used to identify the transplanted cells and to discriminate between the tumorigenic and nontumorigenic cells: antibody M 20, reacting with human cytokeratin no. 8 and not cross-reacting with other species, kindly provided by van Muijen, Department of Pathology, University of Leiden, The Netherlands; antibody Ks.18.174, specific for cytokeratin no. 18, obtained from Progen, Heidelberg, Federal Republic of Germany; and the broad-range cytokeratin antibody lu-5, from Böhringer GmbH, Mannheim, Federal Republic of Germany. An antibody against a 75-kilodalton protein expressed on the surface of tumorigenic but not nontumorigenic cells, thus specifically decorating tumorigenic segregant cells (27), was kindly provided by E. J. Stanbridge.

RESULTS

Continued expression *in vivo* of the HPV18 E6-E7 genes in tumorigenic cells. In initial experiments, total RNA from tumorigenic cells grown in tissue culture was extracted and

FIG. 1. Northern blot hybridization analysis of HPV18 E6-E7 (A) and β -actin (B) mRNAs. Samples of 10 μ g of total RNA extracted from cultured hybrid cells of line 444 (nontumorigenic, lane 1), line CGL3 (tumorigenic, lane 5) and the parental HeLa cells (lane 2) as well as from two different tumors induced by HeLa cells (lanes 3 and 4) and CGL3-cells (lanes 6 and 7) were fractionated, blotted, and hybridized to E6*-E7 DNA (A) and hybridized again to β -actin DNA (B). Both DNA probes were 32 P labeled by random priming (20). The tumors were excised 3 weeks after subcutaneous injection into nude mice.

compared with RNA from tumors induced by the same cells in nude mice. The RNA preparations were subjected to Northern blot hybridizations with 32 P-labeled HPV18-specific cDNA probes. The results confirmed the continued expression of the HPV18 E6-E7 genes in transplanted tumorigenic cells in vivo. A comparison of the hybridization signals with an internal standard, which was provided by hybridization of the filters with a β -actin gene probe, re-

vealed that in the tumors the steady-state levels of the E6-E7 mRNAs were substantially (about three- to fivefold) higher than in the cultured cells (Fig. 1; compare lanes 3 and 4 with lane 2 and lanes 6 and 7 with lane 5, respectively).

HPV18 E6-E7 mRNA levels in nontumorigenic hybrid cells grown in vitro. The nontumorigenic cells, when grown in culture, also expressed the E6-E7 genes (Fig. 1; compare lane 1 with lanes 2 and 5) (38, 45). We have ascertained that there was no subpopulation of cells among the nontumorigenic cells that did not express the E6-E7 genes: in situ hybridization to cultured nontumorigenic and tumorigenic cells, with an HPV18 E6*-E7 antisense cRNA probe, showed in both cell types a homogeneous or nearly homogeneous expression of the E6-E7 genes, as exemplified in Fig. 2 for the nontumorigenic cells. This was also the case when the hybridization was performed after overnight growth of the cells on the collagen gel used for transplantation. Thus, the nontumorigenic cells used for the transplantation experiments were not heterogeneous in their expression of the early HPV genes.

Cell growth and HPV18 E6-E7 gene expression in vivo in nontumorigenic and tumorigenic hybrid cells. Concerning the expression of E6-E7 genes in vivo, Northern blot hybridization data of the nontumorigenic cells were variable and inconclusive, yielding weak or only background hybridization signals.

Therefore, we analyzed the status of the E6-E7 expression of the hybrid cells under in vivo conditions by in situ hybridization-histochemistry to obtain information on the histological state of the nontumorigenic and tumorigenic cells as well as on the cellular levels of expression of the E6-E7 genes of HPV18, in comparison with the expression of cellular genes.

Tumorigenic and nontumorigenic hybrid cells were trans-

FIG. 2. HPV18 E6-E7 mRNA in cultured nontumorigenic cells analyzed by in situ hybridization. The cells were grown on microscope slides to semiconfluency, washed with phosphate-buffered saline, and fixed in 4% paraformaldehyde for 15 min. Hybridization to a E6*-E7 cRNA was for 3 h at 50°C. Acid, heat, and proteinase K treatment were omitted. A, bright field; B, dark field illumination. Magnifications, $\times 380$.

planted onto nude mice (see Materials and Methods). At different times thereafter transplants were removed and subjected to histological examination and to *in situ* hybridization with the following strand-specific cRNA probes: HPV18 E6*-E7 antisense cRNA, for the detection of the E6-E7 mRNAs; HPV18 E6*-E7 sense cRNA (negative control) and antisense cRNA probes specific for the cytoskeletal mRNAs of human β -actin and vimentin. The latter were used to control for the metabolic and transcriptional activity of the transplanted cells.

Since only a limited number of transplantation chambers and animals could be manipulated at one time, three consecutive experiments were performed in which transplants of nontumorigenic and tumorigenic hybrid cells were harvested at the following times after transplantation (p.t.): days 2, 6, and 12 (experiment 1); days 5, 7, and 11 (experiment 2); days 3, 6, 14, and 21 (experiment 3). In all experiments, each time point involved at least two animals, each of which provided a sufficiently large number of tissue sections for *in situ* analysis; an additional animal was used for histological examination only. All cells used for transplantation were derived from the same stock of frozen cells. The tumorigenicity status of cells was assayed by subcutaneous inoculation and long-term observation: none of the animals inoculated with nontumorigenic cells developed a tumor during a 6-month period, whereas tumorous growth was evident in all animals that had been inoculated with tumorigenic cells. Both the histology and the patterns of HPV18 E6-E7 gene expression were consistent among the transplants harvested at the same time points. The data from these three experiments are described and discussed together.

Histological examination of the transplant sections revealed that the nontumorigenic cells initially continued to proliferate. Under the *in vivo* conditions employed here, this period lasted for about 3 days. Figure 3 shows the histology of sections of transplanted cells at low magnification; cells were harvested at day 2, 3, 6, and 14 p.t., respectively. At day 2 p.t., both tumorigenic and nontumorigenic cells had formed a thin epitheliumlike tissue (Fig. 3a and b). At day 3, the nontumorigenic cells were present in a yet larger number, but at no time point thereafter did they show further growth (Fig. 3b, d, f, and h). The tumorigenic cells, on the other hand, continued to grow fast and invaded the mesenchyme of the host animal (Fig. 3a, c, e, and g).

Dark-field photomicrographs showing the hybridization signals obtained from these sections after hybridization with ³²P-labeled HPV18 E6*-E7 antisense RNA probe are presented in Fig. 4. The tumorigenic cells showed a strong, homogeneous expression of the E6-E7 genes at all times analyzed (Fig. 4a, d, g, and j). At day 2, the mRNA level in the nontumorigenic cells was quite similar to that in the tumorigenic cells (Fig. 4b). Already at day 3, however, the nontumorigenic cells showed a sharp reduction in the level of the E6-E7 mRNAs when compared with the tumorigenic cells harvested at the same time and in the nontumorigenic cells on day 2 p.t. (compare Fig. 4e with 4b and d). These reduced levels persisted further (Fig. 4h and k). Thus, there was a high level of HPV18 E6-E7 gene expression in both

tumorigenic and nontumorigenic cells early after transplantation (2 days p.t.), followed by a decline of HPV18 mRNAs specifically in the nontumorigenic cells. A crude estimation of quantitative differences in E6-E7 gene expression between tumorigenic and nontumorigenic cells and at different times was obtained by grain counting. A comparison of the hybridization signals generated by the E6-E7-specific probe in the nontumorigenic cells at day 2 p.t. with those generated at days 3 and later revealed differences by a factor of approximately 4 and higher. The suppression of E6-E7 gene expression was equally evident when the hybridization signals displayed by the nontumorigenic cells at day 3 or 6 were compared with those displayed by the tumorigenic cells at the same time points.

Cellular gene expression *in vivo* in nontumorigenic hybrid cells. The transplanted cells were also examined for the expression of cellular genes. For that purpose the cytoskeletal genes β -actin and vimentin were chosen, since their level of expression should be a direct indicator of the metabolic and transcriptional activity of the cells. Both genes showed constantly high mRNA levels in the tumorigenic cells at all times analyzed (data not shown) and in the nontumorigenic cells until day 14 p.t. (for β -actin, see Fig. 4c, f, i, and l; data for vimentin are not shown). Only at day 21 p.t. were the mRNA levels of β -actin and vimentin reduced, indicating that cell death had now commenced (data not shown). A crude estimation of grain densities generated by the β -actin and vimentin probes in both cell types confirmed the interpretation derived from visual inspection that the mRNAs of both genes stayed at high levels at least until day 14 in the nontumorigenic cells. These results demonstrated that the nontumorigenic cells were still vital 11 days after suppression of HPV18 E6-E7 gene expression.

Heterogeneity of HPV18 E6-E7 mRNA levels in nontumorigenic hybrid cells *in vivo*. At day 14 p.t., the collagen was largely dissolved and/or repopulated by mouse mesenchymal cells, and so the transplanted cells were now dispersed among the mesenchymal cells of the host animal. Indirect immunofluorescence staining with monoclonal antibodies specific for human cytokeratins no. 8 (data not shown) and 18 (Fig. 5b) was therefore employed to identify the transplanted human hybrid cells. The great majority of the immunofluorescence-positive cells revealed no detectable E6-E7 mRNAs (Fig. 4k and 5a), but there were isolated foci of cells displaying significant E6-E7 mRNA levels (Fig. 5a). However, at day 21 p.t. such foci could hardly be observed anymore (data not shown). To determine whether these focally increased levels of E6-E7 mRNAs (denoted by arrows in Fig. 5a) were due to contamination by tumorigenic cells or even to the appearance of tumorigenic segregants, we reexamined all sections by using an antibody against a 75-kilodalton protein recognizing a surface antigen expressed in the tumorigenic hybrid cells (and in HeLa cells) but not in the parent nontumorigenic hybrid cells (27). Whereas the epithelia formed by the tumorigenic hybrid cells reacted positively at all time points tested (Fig. 5c), the nontumorigenic hybrid cells never showed any staining,

FIG. 3. Low-magnification micrographs of organotypical cultures of tumorigenic cells (a, c, e, g) and nontumorigenic cells (b, d, f, h) grown *in vivo* for different periods of time. The cells were transplanted onto different animals as described in Materials and Methods, and the transplants removed *in toto* after 2 days (a, b), 3 days (c, d), 6 days (e, f), and 14 days (g, h). The cryostat sections were stained with hematoxylin and eosin. During sectioning, the nontumorigenic cells grown for 6 days have folded back. Note the continuous growth of the tumorigenic cells and the stop in growth of the nontumorigenic cells between days 3 and 6. Magnifications: a and b, ca. $\times 116$; c through h, ca. $\times 174$.

FIG. 4. HPV18 E6-E7 gene expression in organotypical cultures of tumorigenic cells (a, d, g, j) and nontumorigenic cells (b, e, h, k) and of β -actin gene expression in nontumorigenic cells (c, f, i, l): low-magnification, dark-field micrographs. The same sections or sections parallel to those presented in Fig. 3 are shown. Note that at day 2 p.t. HPV18 E6-E7 gene expression in the nontumorigenic cells was very similar to that seen in the tumorigenic cells (a, b, c), but at day 3 (d, e, f) and later (day 6: g, h, i; day 14: j, k, l) was drastically reduced. β -Actin gene expression (c, f, i, l), on the other hand, remained at high levels. Magnifications are as in Fig. 3.

including the foci with increased E6-E7 mRNA levels in 14-day-old transplants (Fig. 5d, showing region corresponding to Fig. 5a and b). This clearly excluded cross-contamination by tumorigenic cells.

The nontumorigenic cells showed heterogeneous behavior regarding the extent of E6-E7 mRNA suppression also at the earlier times p.t., as shown at higher magnification in Fig. 6. Starting with day 3 p.t., in a large subset of the nontumorigenic cells HPV18 E6-E7 expression was completely shut off (Fig. 6c and d) when compared with the hybridization signals obtained in the nontumorigenic cells at day 2 p.t. and also in the tumorigenic cells at day 3 and later. However, a further subset of the nontumorigenic cells showed a markedly reduced but still clearly detectable level of E6-E7 mRNAs (Fig. 6d); yet, none of these cells showed any reduction in β -actin or vimentin mRNA levels, confirming their viability until at least day 14 (Fig. 4c, f, i, and l).

HPV18 E6-E7 gene expression after subcutaneous inoculation. HPV18 E6-E7 gene expression of the tumorigenic and nontumorigenic cells was also analyzed in nude mice after injection by the subcutaneous route (Fig. 7). In this experiment tissue samples were removed from the sites of previous injection at days 5 and 21 (tumorigenic cells) or days 5 and 28 (nontumorigenic cells), respectively. The nontumorigenic cells were markedly down-regulated in their E6-E7 mRNA levels at both day 5 and 28 (Fig. 7). It has to be stressed, however, that even at day 28 there were some small regions still revealing hybridization levels above background, although always substantially lower than the corresponding signals seen over the tumorigenic cells.

DISCUSSION

This study has revealed quantitative differences in HPV18 E6-E7 gene expression in nontumorigenic versus tumori-

FIG. 5. Identification of and discrimination between the nontumorigenic and tumorigenic cells grown in vivo for 14 days by indirect immunofluorescence. (a) A few cells positive for HPV18 E6-E7 mRNA can be seen in this section after in situ hybridization (arrows). (b) Monoclonal antibody Ks 18.174, specific for cytokeratin no. 18, identified the transplanted human cells in adjacent sections (a). A monoclonal antibody against a 75-kilodalton protein, specifically recognizing tumorigenic cells (c), did not react with any transplanted nontumorigenic cells, including those displaying increased E6-E7 mRNA levels (d, region corresponding to a and b). Magnification, $\times 400$.

FIG. 6. Heterogeneity of HPV18 E6-E7 gene expression in nontumorigenic cells at early time points p.t. analyzed by in situ hybridization. (a) Tumorigenic cells, grown for 6 days (positive control); (b, c, and d) nontumorigenic cells grown for 2, 3, and 6 days, respectively. Arrows demarcate significantly E6-E7-positive cells, arrowheads indicate E6-E7 negative cells. For further details see the text. Magnifications: a, $\times 500$; b through d, $\times 780$.

genic HeLa $\times$ fibroblast hybrid cells when comparing growth conditions in vitro (tissue culture) and in vivo (nude mice). We have chosen the hybrid cell lines 444 (nontumorigenic) and a tumorigenic segregant thereof (CGL3) as model system, since the extensive characterization of these two cell lines has shown their very close biochemical and genetic relationship (50). For the tumorigenic HeLa $\times$ fibroblast segregant line, Northern blot and in situ hybridization experiments have established that the continuous and invasive growth in vivo of these cells was correlated with constantly high levels of HPV18 mRNAs at all times examined after transplantation or subcutaneous inoculation (Fig. 1, 4, 5, 6, and 7).

The uncontrolled growth of the tumorigenic segregants impressively contrasted to the host-induced inhibition of cell growth of the nontumorigenic hybrids (Fig. 3 and 4). Growth inhibition in vivo was apparently taking effect between day 3 and day 5 p.t., since from day 3 on the number of cell layers on the collagen gel did not increase further

(compare Fig. 3d and f). These data are in good agreement with the observations of Stanbridge and co-workers, who reported that proliferation was seen only for 3 to 4 days after subcutaneous inoculation of the nontumorigenic HeLa $\times$ fibroblast hybrid cells (50, 51).

In vitro (i.e., either in petri dishes or as organotypical cultures on collagen) as well as in the initial period of growth in vivo (i.e., 2 days after transplantation of the organotypical cultures), the level of HPV18 E6-E7 mRNA in the nontumorigenic cells was as high as that in the tumorigenic hybrid cells (Fig. 1, 2, 3, and 4). In all specimens taken at day 3 p.t., the HPV18 E6-E7 mRNA levels were already markedly reduced (Fig. 4), and they were similarly reduced in all specimens taken at later times (Fig. 3 and 4). We noted, however, that the reduction of the HPV18 E6-E7 mRNA levels was not homogeneous in these cells. Whereas in the majority of cells HPV18 E6-E7 gene expression dropped below the level of detection of the in situ hybridization method, in a subset of cells expression was still obvious but

FIG. 7. In situ hybridization analysis of HPV18 E6-E7 gene expression after subcutaneous injection of tumorigenic and nontumorigenic cells. Low-magnification dark-field (a, c, e) and bright-field (b, d, f) photomicrographs of tumorigenic cells (a, b) and nontumorigenic cells (c, d), each grown in vivo for 5 days, and nontumorigenic cells grown in vivo for 28 days (e, f). Note that at both time points the nontumorigenic cells show a substantial reduction but not a complete homogeneous shutoff of E6-E7 gene expression with small areas still displaying weak hybridization signals. Magnification, ca. $\times 172$.

clearly reduced. How much this differential pattern is due to distinct cytogenetic subpopulations or epigenetic phenomena remains to be seen.

In contrast to the reduced levels of HPV18 E6-E7 mRNAs at day 3 p.t. and onward, high levels of β -actin (Fig. 4) and vimentin (data not shown) transcripts were fully maintained in the nontumorigenic hybrid cells until (at least) day 14 p.t., with expression levels very similar to those in the tumorigenic cells (data not shown). This clearly shows that the nontumorigenic cells remained viable for at least 2 weeks after transplantation and consequently excluded cell death as responsible for the decrease of HPV18 E6-E7 gene expression. Only at day 21 p.t. were the β -actin mRNA levels in the nontumorigenic cells reduced, indicating that cell death had now commenced. Again these results obtained for the transplanted nontumorigenic cells are in agreement with data reported earlier (50, 51) indicating that the nontumorigenic HeLa $\times$ fibroblast hybrid cells, when injected subcutaneously, remain as a viable but nondividing tissue for up to 15 days and that the early cessation of cell division is not accompanied by cell death.

Thus, the histology and *in situ* hybridization data of the transplantation experiments strongly indicate that growth inhibition *in vivo* of the nontumorigenic cells is correlated with a down-regulation of HPV18 E6-E7 gene expression, whereas the continuous growth of the tumorigenic cells is correlated with consistently high levels of HPV18 E6-E7 mRNAs.

Reduced levels of HPV18 E6-E7 mRNAs were also observed in the nontumorigenic cells after subcutaneous inoculation (Fig. 7). In comparison with the transplantation assay, subcutaneous injection has several disadvantages for the examination of the *in vivo* expression of genes. First, repeating the identification and isolation of the nontumorigenic cells requires very large numbers of cells to be injected (this also applies to the tumorigenic cells for isolation at early times after subcutaneous inoculation when the cells have not yet grown to tumors of easily detectable size). This leads to pronounced necrotic reactions (51), making an exact evaluation of the metabolic and transcriptional status of individual cells very difficult. Furthermore, the large number of injected cells may lead to differential influences of the host mesenchymal environment upon the peripheral injected cells as compared with influences upon the more centrally located cells. Irrespective of these limitations, however, the results obtained after subcutaneous injection are in agreement with the findings obtained in the transplantation assay.

The E6 and E7 genes of the cancer-associated HPV16 and HPV18 have been implicated as essential factors in HPV-linked carcinogenesis because of their presence and expression in cervical carcinomas and cell lines derived from cervical carcinomas (1, 19, 43, 44, 46, 48), their ability to interfere with normal cell growth control *in vitro* (immortalization of normal human cells) (2, 21, 22, 33), and their influence on proliferation *in vitro* of malignant cells as demonstrated for the HPV18-positive cervical carcinoma cell line C4-I (53). All of these data implicate that the HPV18 E6-E7 genes might also be involved in growth regulation of the HeLa $\times$ fibroblast hybrid cells. This notion has received strong support by our observation of a synchrony between HPV18 E6-E7 gene expression and growth behavior *in vivo*. Does this synchrony reflect a causative relation in the sense that inhibition of cell growth might be a consequence of HPV18 E6-E7 down-regulation? It is tempting to speculate about such a relation (see below), but at present every answer to this question must remain conjectural. However,

the results of previous *in vitro* studies and of the *in vivo* experiments reported here identify the HPV18 E6-E7 genes as candidate regulatory genes for transformation and oncogenic growth of the HeLa-hybrid cells and as possible targets of host-mediated suppression operating only in the nontumorigenic cells.

It is, however, completely unknown which cellular regulatory mechanisms are responsible (i) for down-regulation of the HPV18 E6-E7 genes and (ii) for growth inhibition of the nontumorigenic hybrid cells *in vivo*, and whether these are (partly or completely) different or identical. In the HeLa $\times$ fibroblast hybrids, the nontumorigenic phenotype is linked to the presence of two copies of fibroblast chromosome 11 (52). In addition, a normal chromosome 11 is able to suppress tumorigenicity of HPV18-positive HeLa and HPV16-positive SiHa cervical carcinoma cells (26, 40). Furthermore, it has been shown that human embryonic fibroblasts carrying a deletion in the short arm of one chromosome 11, in contrast to diploid human embryonic fibroblasts (47), were susceptible to transformation by HPV16 early region DNA and efficiently expressed the viral genes. All of these data indicate that chromosome 11 probably harbors tumor suppressor genes whose loss of function might be important for the development of HPV-associated cervical carcinomas. Deletions in certain regions of chromosome 11 are also associated with Wilm's embryonic kidney tumor and with other malignancies and are implicated in the loss of tumor suppressor genes (56; for reviews, see references 37 and 39).

Experimental evidence for interactions between the products of cellular tumor suppressor genes and HPV transforming genes has recently been obtained by demonstrating complex formation between the retinoblastoma tumor suppressor gene product RB and the HPV16 and HPV18 E7 proteins (18, 34), and between the cellular p53 protein and the HPV16 and HPV18 E6 proteins (57). These data suggested that the HPV E7 and E6 proteins might interfere with the normal functions of the RB and p53 proteins, which seem to play crucial roles for normal cell growth control (10, 13).

In the intracellular surveillance hypothesis, a class of cellular genes called CIF genes exists in normal cells; these genes can negatively regulate incoming viral oncogenes, such as the HPV18 and HPV16 E6-E7 genes. Furthermore, the irreversible inactivation or loss of these CIF genes is postulated to be necessary to release the HPV E6-E7 genes from cellular suppression *in vivo* and to represent an essential event in the multistep development of cervical carcinoma (60, 62). In the context of this hypothesis, the shutdown of HPV18 E6-E7 gene expression *in vivo* in the nontumorigenic cells results from host-mediated activation of such cellular suppressor genes. With regard to the specific chromosome losses of tumorigenic hybrid segregants (52), intact CIF genes should be contributed by the normal parental cell to the nontumorigenic HeLa hybrid cells and be again lost in the tumorigenic segregants. This scenario would also provide a possible explanation for the nontumorigenic phenotype of HPV-immortalized human cells: cell immortalization *in vitro* due to expression of the HPV E6-E7 genes (2, 21-23) but nontumorigenicity *in vivo* at least in part due to suppression of the HPV E6-E7 genes. *In vitro* studies with 5-azacytidine have suggested the existence of host cell mechanisms that negatively regulate HPV E6-E7 transcription specifically in nontumorigenic HPV-containing cells but are not active under normal tissue culture conditions (38). It remains to be clarified, however, whether the negative regulatory mechanisms induced in the nontumorigenic hybrid cells by

5-azacytidine treatment *in vitro* are identical to those induced *in vivo*.

Thus, the *in situ* hybridization data presented in this report point to a different regulation of HPV18 E6-E7 gene expression *in vivo* in nontumorigenic versus tumorigenic cells. Future studies are necessary to analyze whether the correlated down-regulation of HPV18 E6-E7 gene expression and growth inhibition *in vivo* of the nontumorigenic cells are indeed causally related. This will require the identification of cellular genes that interfere with the expression of the HPV transforming genes. This hopefully will contribute to a better understanding of the cellular growth-regulatory networks affected in HPV-associated carcinogenesis.

ACKNOWLEDGMENTS

We thank M. Mappes and H. Steinbauer for expert technical assistance and specifically acknowledge the kind gift of antibody 75K as well as valuable discussion by E. J. Stanbridge, Department of Microbiology and Molecular Genetics, University of California at Irvine.

This work was supported by the Deutsche Forschungsgemeinschaft.

LITERATURE CITED

- Baker, C. C., W. C. Phelps, V. Lindgren, M. J. Braun, M. A. Gonda, and P. M. Howley. 1987. Structural and transcriptional analysis of human papillomavirus type 16 sequences in cervical carcinoma cell lines. *J. Virol.* **61**:962-971.
- Barbosa, M. S., and R. Schlegel. 1989. The E6 and E7 genes of HPV-18 are sufficient for inducing two-stage *in vitro* transformation of human keratinocytes. *Oncogene* **4**:1529-1532.
- Bedell, M. A., K. H. Jones, S. R. Grossman, and L. A. Laimins. 1989. Identification of human papillomavirus type 18 transforming genes in immortalized and primary cells. *J. Virol.* **63**:1247-1255.
- Bedell, M. A., K. H. Jones, and L. A. Laimins. 1987. The E6-E7 region of human papillomavirus type 18 is sufficient for transformation of NIH 3T3 and Rat-1 cells. *J. Virol.* **61**:3635-3640.
- Bernard, B. A., C. Bailly, M.-C. Lenoir, M. Darmon, F. Thierry, and M. Yaniv. 1989. The human papillomavirus type 18 (HPV18) E2 gene product is a repressor of the HPV18 regulatory region in human keratinocytes. *J. Virol.* **63**:4317-4324.
- Bohnert, A., J. Hornung, I. C. Mackenzie, and N. E. Fusenig. 1986. Epithelial-mesenchymal interactions control basement membrane production and differentiation in culture and transplanted mouse keratinocytes. *Cell Tissue Res.* **244**:413-429.
- Bosch, F. X., J.-P. Ouhayoun, B. L. Bader, C. Collin, C. Grund, I. Lee, and W. W. Franke. 1989. Extensive changes in cytokeratin expression patterns pathologically affected human gingiva. *Virchows Arch. B Cell Pathol.* **58**:59-77.
- Chirgwin, I. M., A. A. Przybyla, R. MacDonald, and W. J. Rutter. 1979. Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry* **18**:5295-5299.
- Choo, K.-B., C.-C. Pan, and S.-H. Han. 1987. Integration of human papillomavirus type 16 into cellular DNA of cervical carcinoma: preferential deletion of the E2 gene and invariable retention of the long control region and the E6/E7 open reading frames. *Virology* **161**:259-261.
- Cooper, J. A., and P. Whyte. 1989. RB and the cell cycle: entrance or exit? *Cell* **58**:1009-1011.
- Cox, K. H., D. V. DeLeon, L. M. Angerer, and R. C. Angerer. 1984. Detection of mRNAs in sea urchin embryos by *in situ* hybridization using asymmetric RNA probes. *Dev. Biol.* **101**:485-502.
- Crook, T., J. P. Morgenstern, L. Crawford, and L. Banks. 1989. Continued expression of HPV-16 E7 protein is required for maintenance of the transformed phenotype of cells cotransformed by HPV-16 plus EJ-ras. *EMBO J.* **8**:513-519.
- De Caprio, J. A., J. W. Ludlow, D. Lynch, Y. Furukawa, J. Griffin, H. Pwnica-Worms, C.-M. Huang, and D. M. Livingston. 1989. The product of the retinoblastoma susceptibility gene has properties of a cell cycle regulatory element. *Cell* **58**:1085-1095.
- Der, C. J., and E. J. Stanbridge. 1978. Lack of correlation between the decreased expression of cell surface LETS protein and tumorigenicity in human cell hybrids. *Cell* **15**:1241-1251.
- De Villiers, E.-M. 1989. Heterogeneity of the human papillomavirus group. *J. Virol.* **63**:4898-4903.
- Dürst, M., R. T. Dzarlieva-Petrusevska, P. Boukamp, N. E. Fusenig, and L. Gissmann. 1987. Molecular and cytogenetic analysis of immortalized human primary keratinocytes obtained after transfection with human papillomavirus type 16 DNA. *Oncogene* **1**:251-256.
- Dürst, M., A. Kleinheinz, M. Hotz, and L. Gissmann. 1985. The physical state of human papillomavirus type 16 DNA in benign and malignant genital tumors. *J. Gen. Virol.* **66**:1515-1522.
- Dyson, N., P. M. Howley, K. Münger, and E. Harlow. 1989. The human papilloma virus-16 E7 oncoprotein is able to bind to the retinoblastoma gene product. *Science* **243**:934-937.
- El Awady, M. K., J. B. Kaplan, S. J. O'Brien, and R. D. Burk. 1987. Molecular analysis of integrated human papillomavirus 16 sequences in the cervical cancer cell line SiHa. *Virology* **159**:389-398.
- Feinberg, A. P., and B. Vogelstein. 1984. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **137**:266-267.
- Hawley-Nelson, P., K. H. Vousden, N. L. Hubbert, D. R. Lowy, and J. T. Schiller. 1989. HPV16 E6 and E7 proteins cooperate to immortalize human foreskin keratinocytes. *EMBO J.* **8**:3905-3910.
- Hudson, J. B., M. A. Bedell, D. J. McCance, and L. A. Laimins. 1990. Immortalization and altered differentiation of human keratinocytes *in vitro* by the E6 and E7 open reading frames of human papillomavirus type 18. *J. Virol.* **64**:519-526.
- Kanda, T., A. Furuno, and K. Yoshiike. 1988. Human papillomavirus type 16 open reading frame E7 encodes a transforming gene for rat 3Y1 cells. *J. Virol.* **62**:610-613.
- Kanda, T., S. Watanabe, and K. Yoshiike. 1988. Immortalization of primary rat cells by human papillomavirus type 16 subgenomic DNA fragments controlled by the SV40 promoter. *Virology* **165**:321-325.
- Kaur, P., and J. K. McDougall. 1988. Characterization of primary human keratinocytes transformed by human papillomavirus type 18. *J. Virol.* **62**:1917-1924.
- Koi, M., H. Morita, H. Yamada, H. Satoh, J. C. Barrett, and M. Oshimura. 1989. Normal human chromosome 11 suppresses tumorigenicity of human cervical tumor cell line SiHa. *Mol. Carcinogen.* **2**:12-21.
- Latham, K. M., and E. J. Stanbridge. 1990. Identification of the HeLa tumor-associated antigen, p75/150, as intestinal alkaline phosphatase and evidence for its transcriptional regulation. *Proc. Natl. Acad. Sci. USA* **87**:1263-1267.
- Le, J.-Y., and V. Defendi. 1988. A viral-cellular junction fragment from a human papillomavirus type 16-positive tumor is competent in transformation of NIH 3T3 cells. *J. Virol.* **62**:4420-4426.
- Lehn, H., L. L. Villa, F. Marziona, M. Hilgarth, H.-G. Hillemanns, and G. Sauer. 1988. Physical state and biological activity of human papillomavirus genomes in precancerous lesions of the female genital tract. *J. Gen. Virol.* **69**:187-196.
- Matlashewski, G., J. Schneider, L. Banks, N. Jones, A. Murray, and L. Crawford. 1987. Human papillomavirus type 16 DNA cooperates with activated *ras* in transforming primary cells. *EMBO J.* **6**:1741-1746.
- Matsukura, T., S. Koi, and M. Sugase. 1989. Both episomal and integrated forms of human papillomavirus type 16 are involved in invasive cervical cancers. *Virology* **172**:63-72.
- McCance, D. J., R. Kopan, E. Fuchs, and L. A. Laimins. 1988. Human papillomavirus type 16 alters human epithelial cell differentiation *in vitro*. *Proc. Natl. Acad. Sci. USA* **85**:7169-7173.
- Münger, K., W. C. Phelps, V. Bubbs, P. M. Howley, and R. Schlegel. 1989. The E6 and E7 genes of the human papillomavirus type 16 together are necessary and sufficient for transforma-

- tion of primary human keratinocytes. *J. Virol.* **63**:4417-4421.
34. Münger, K., B. A. Werness, N. Dyson, W. C. Phelps, E. Harlow, and P. M. Howley. 1989. Complex formation of human papillomavirus E7 proteins with the retinoblastoma tumor suppressor gene product. *EMBO J.* **8**:4009-4105.
 35. Phelps, W. C., C. L. Yee, K. Münger, and P. M. Howley. 1988. The human papillomavirus type 16 E7 gene encodes transactivation and transformation functions similar to those of adenovirus E1A. *Cell* **53**:539-547.
 36. Pirisi, L., K. E. Creek, J. Doniger, and J. A. DiPaolo. 1988. Continuous cell lines with altered growth and differentiation properties originate after transfection human keratinocytes with human papillomavirus type 16 DNA. *Carcinogenesis* **9**:1573-1579.
 37. Ponder, B. 1988. Gene losses in human tumours. *Nature (London)* **335**:400-402.
 38. Rösl, F., M. Dürst, and H. zur Hausen. 1988. Selective suppression of human papillomavirus transcription in non-tumorigenic cells by 5-azacytidine. *EMBO J.* **7**:1321-1328.
 39. Sager, R. 1989. Tumor suppressor genes: the puzzle and the promise. *Science* **246**:1406-1412.
 40. Saxon, P. J., E. S. Srivatsan, and E. J. Stanbridge. 1986. Introduction of human chromosome 11 via microcell transfer controls tumorigenic expression of HeLa cells. *EMBO J.* **5**:3461-3466.
 41. Schlegel, R., W. C. Phelps, Y.-L. Zhang, and M. Barbosa. 1988. Quantitative keratinocyte assay detects two biological activities of human papillomavirus DNA and identifies viral types associated with cervical carcinoma. *EMBO J.* **7**:3181-3187.
 42. Schneider, A. 1990. Latent and subclinical genital HPV infections, p. 2-5. *In* Papillomavirus report; international literature on papillomavirus infection, vol. 1.
 43. Schneider-Gädicke, A., and E. Schwarz. 1986. Different human cervical carcinoma cell lines show similar transcription patterns of human papillomavirus type 18 early genes. *EMBO J.* **5**:2285-2292.
 44. Schwarz, E., U. K. Freese, L. Gissmann, W. Mayer, B. Roggenbuck, A. Stremlau, and H. zur Hausen. 1985. Structure and transcription of human papillomavirus sequences in cervical carcinoma cells. *Nature (London)* **314**:111-114.
 45. Schwarz, E., A. Schneider-Gädicke, and H. zur Hausen. 1987. Human papillomavirus type-18 transcription in cervical carcinoma cell lines and in human cell hybrids, p. 47-53. *In* B. M. Steinberg, J. L. Brandsma, and L. B. Taichman (ed.), *Cancer cells* vol. 5. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
 46. Seedorf, K., T. Oltersdorf, G. Krämmmer, and W. Röwekamp. 1987. Identification of early proteins of human papillomaviruses type 16 (HPV 16) and type 18 (HPV 18) in cervical carcinoma cells. *EMBO J.* **6**:139-144.
 47. Smits, H. L., E. Raadsheer, I. Rood, S. Mehendale, R. M. Slater, J. van der Noordaa, and J. ter Schegget. 1988. Induction of anchorage-independent growth of human embryonic fibroblasts with deletion in the short arm of chromosome 11 by human papillomavirus type 16 DNA. *J. Virol.* **62**:4538-4543.
 48. Smotkin, D., and F. O. Wettstein. 1986. Transcription of human papillomavirus type 16 early genes in a cervical cancer and a cancer-derived cell line and identification of the E7 protein. *Proc. Natl. Acad. Sci. USA* **83**:4680-4684.
 49. Stanbridge, E. J. 1976. Suppression of malignancy in human cells. *Nature (London)* **260**:17-20.
 50. Stanbridge, E. J., and R. Ceredig. 1981. Growth-regulatory control of human cell hybrids in nude mice. *Cancer Res.* **41**:573-580.
 51. Stanbridge, E. J., C. J. Der, C.-J. Doersen, R. Y. Nishimi, D. M. Peehl, B. E. Weissman, and J. E. Wilkinson. 1982. Human cell hybrids: analysis of transformation and tumorigenicity. *Science* **215**:252-259.
 52. Srivatsan, E. S., W. F. Benedict, and E. J. Stanbridge. 1986. Implication of chromosome 11 in the suppression of neoplastic expression in human cell hybrids. *Cancer Res.* **46**:6174-6179.
 53. Von Knebel Doeberitz, M., T. Oltersdorf, E. Schwarz, and L. Gissmann. 1988. Correlation of modified human papilloma virus early gene expression with altered growth properties in C4-1 cervical carcinoma cells. *Cancer Res.* **48**:3780-3786.
 54. Vousden, K. H. 1989. Human papillomaviruses and cervical carcinoma. *Cancer Cells* **1**:43-50.
 55. Wagatsuma, M., K. Hashimoto, and T. Matsukura. 1990. Analysis of integrated human papillomavirus type 16 DNA in cervical cancers: amplification of viral sequences together with cellular flanking sequences. *J. Virol.* **64**:813-821.
 56. Weissmann, B. E., P. Saxon, S. R. Pasquale, G. R. Jones, A. G. Geiser, and E. J. Stanbridge. 1987. Introduction of a normal human chromosome 11 into a Wilms' tumor cell line controls its tumorigenic expression. *Science* **236**:175-180.
 57. Werness, B. A., A. J. Levine, and P. M. Howley. 1990. Association of human papillomavirus types 16 and 18 E6 proteins with p53. *Science* **248**:76-79.
 58. Woodworth, C. D., P. E. Bowden, J. Doniger, L. Pirisi, W. Barnes, W. D. Lancaster, and J. A. DiPaolo. 1988. Characterization of normal human exocervical epithelial cells immortalized *in vitro* by papillomavirus types 16 and 18 DNA. *Cancer Res.* **48**:4620-4628.
 59. Worst, P. K. M., I. C. Mackenzie, and N. E. Fusenig. 1982. Reformation of organized epidermal structure by transplantation of suspensions and cultures of epidermal and dermal cells. *Cell Tissue Res.* **225**:65-77.
 60. zur Hausen, H. 1977. Cell-virus gene balance hypothesis of carcinogenesis. *Behring Inst. Mitt.* **61**:23-30.
 61. zur Hausen, H. 1986. Intracellular surveillance of persisting viral infections. *Lancet* **ii**:489-491.
 62. zur Hausen, H. 1989. Papillomaviruses as carcinomaviruses. *Adv. Viral Oncol.* **8**:1-26.