Sexuality and older people

Doctors should ask patients, regardless of age, about sex

PHOTOSEARCH

RESEARCH, p 151

Peggy J Kleinplatz associate professor and clinical professor, Department of Family Medicine, Faculty of Medicine and School of Psychology, University of Ottawa, Ottawa, ON, Canada K1N 6N5 kleinpla@uottawa.ca

Competing interests: None declared.

Provenance and peer review: Commissioned; not externally peer reviewed.

Cite this as: *BMJ* 2008;337:a239
doi: 10.1136/bmj.a239

Much of the literature on sexuality in elderly people focuses on sexual problems, leaving clinicians with the impression that older adults have either dismal or non-existent sex lives. Few data are available on “normal” sexuality in elderly people, let alone the entire spectrum of sexual expression including optimal sexuality.¹ Beckman and colleagues’ linked study makes a welcome contribution to the limited literature on sexuality in older people.²

A major contribution of Beckman and colleagues’ study is that it focuses on sexual attitudes and behaviour in a sample of people—not patients—who are not seeking treatment for sexual dysfunction or attending a general medical clinic. The methodology is strong, using consistent interviewing techniques over a 30 year period to produce four comparable sets of cross sectional data from 1971 to 2001.

Current knowledge suggests that sexual functioning and frequency decline with age and that sex decreases in importance over time.³ The existing literature emphasises the widespread prevalence of sexual difficulties in men and women.⁴⁻⁵ In contrast, Beckman and colleagues provide good news—sex is an important and positive part of the lives of their 70 year old participants, and more so for the current cohort of men and women than for their predecessors in 1971. Although these data are invaluable, the study does have limitations. Sexual activity was defined as sexual intercourse, and the researchers’ questions about same sex activities and self stimulation were discontinued after 1971 for fear of offending participants.

The study reports that subjective sexual satisfaction is increasing, especially in women, even if sexual dysfunctions are present. Some dysfunctions such as female anorgasmia and erectile dysfunction are decreasing, whereas others such as ejaculatory dysfunction in men have increased over the past 30 years. The authors speculate that the decrease in erectile dysfunction in 70 year old men may result from the availability of phosphodiesterase type 5 inhibitors. Male sexual dissatisfaction and ejaculatory dysfunction increased in the latest cohort. One interpretation is that older men are “performing” better sexually thanks to erectogenic drugs, but enjoying themselves less, thus the difficulty in male orgasm. The meaning of these findings is worthy of further investigation.

Attitudes to sexuality seem to be converging in men and women even though some behaviours remain strikingly constant. Beckman and colleagues seemingly link the increasingly early sexual debut seen over the past

30 years with increasingly positive attitudes to sexuality over time. The implication is that a generation’s sexual change—perhaps linked to the sexual revolution of 1965-75—is evident in this latest cohort of 70 year olds in 2001. Yet, interestingly, both men and women continue to blame men when sexual intercourse ceases between them. This finding has been consistent for 40 years.⁶⁻⁷

Perhaps some aspects of heterosexual relationships are so deeply ingrained that they are more resistant to change. Even if women seem to be coming into their own sexually—and more satisfied than ever in the latest cohort—years of men being in charge of “making the first move” in adolescent sexual encounters in the 1940s and during marriage in early adulthood in the 1950s and 1960s has led to the expectation that men remain responsible for making sex happen. Thus, attributing the responsibility for the frequency or lack of sex to men continues. Perhaps the findings are a manifestation of the time lag between a change in attitude and the ultimate shift in sexual behaviour patterns in heterosexual couples. Clinicians should be sensitive to this mindset when probing into patients’ concerns over sexual frequency, desire, initiation, satisfaction, and their meanings to all parties.

What are the implications of these findings for clinical practice? Doctors in general are known to be uncomfortable about asking patients questions about their sex lives. This is particularly so when the patients’ personal characteristics differ from their own (for example, their sex, age, sexual orientation).⁸⁻⁹ This may be especially disadvantageous when dealing with elderly patients who are already assumed to be invisible and post-sexual by society. Such people may be even less likely than most to approach their doctors with sexual problems and concerns, although research shows that most people hope that their doctors will approach them.¹⁰ Given that sex plays an increasingly valuable role in the lives of older men and women, Beckman and colleagues’ study reinforces the dictum that doctors should ask—and be trained to ask—every patient, regardless of age, “Any sexual concerns?”⁹ Doctors are well placed to normalise and affirm the value of fulfilling sexual relations for the wellbeing of older patients.

- 1 Kleinplatz PJ, Ménard AD. Building blocks towards optimal sexuality: constructing a conceptual model. *Fam J Couns Ther Couples Fam* 2007;15:72-8.
- 2 Beckman N, Waern M, Gustafson D, Skoog I. Secular trends in self reported sexual activity and satisfaction in Swedish 70 year olds: cross sectional survey of four populations, 1971-2001. *BMJ* 2008; doi: 10.1136/bmj.a279.
- 3 Lindau ST, Schumm LP, Laumann EO, Levinson W, O’Muircheartaigh

- CA, Waite LJ. A study of sexuality and health among older adults in the United States. *N Engl J Med* 2007;357:762-74.
- 4 Laumann EO, Nicolosi A, Glasser DB, Paik A, Gingell C, Moreira E, et al. Sexual problems among women and men aged 40-80 y: prevalence and correlates identified in the global study of sexual attitudes and behaviors. *Int J Impot Res* 2005;17:39-57.
 - 5 Nicolosi A, Buvat J, Glasser DB, Hartmann U, Laumann EO, Gingell C. Sexual behaviour, sexual dysfunctions and related help seeking patterns in middle-aged and elderly Europeans: the global study of sexual attitudes and behaviors. *World J Urol* 2006;24:423-8.
 - 6 Pfeiffer E, Verwoerd A, Wang HS. Sexual behavior in aged men and women. *Arch Gen Psychiatry* 1968;19:753-58.
 - 7 Verwoerd A, Pfeiffer E, Wang HS. Sexual behavior in senescence. Changes in sexual activity and interest of aging men and women. *J Geriatr Psychiatry* 1969;2:163-80.
 - 8 Maurice WL. *Sexual medicine in primary care*. St Louis: Mosby, 1999.
 - 9 Moser C. How to ask sex questions during a medical interview. *San Francisco Med* 2005;78:22-4.
 - 10 Metz M, Seifert MH. Differences in men's and women's sexual health needs and expectations of physicians. *Can J Hum Sex* 1993;2:53-9.

Improving the safety of peripheral intravenous catheters

Specialised teams could bring greater benefit than routine replacement

GARY PARKER/SPL

RESEARCH, p 157

Dennis G Maki professor of medicine and hospital epidemiologist, Section of Infectious Diseases, Department of Medicine, University of Wisconsin School of Medicine and Public Health, Madison, WI 53792, USA
dgmaki@medicine.wisc.edu
Competing interests: None declared.

Provenance and peer review: Commissioned; not externally peer reviewed.

Cite this as: *BMJ* 2008;337:a630
doi: 10.1136/bmj.a630

Current guidelines from the United Kingdom¹ and Australia² recommend routine replacement of peripheral intravenous catheters every 48-72 hours to prevent painful infusion phlebitis and rare but life threatening peripheral catheter related bacteraemia. In the United States, Centers for Disease Control and Prevention guidelines recommend replacement every 72-96 hours.³ However, numerous large prospective cohort studies⁴⁻⁸ provide convincing evidence that the risk of bacteraemia associated with the small Teflon or polyetherurethane catheters now widely used in hospitals is only about 0.1-0.3 per 100 catheters.^{9 10} Because many hospitals do not have a team of nurses responsible for the insertion and care of peripheral intravenous catheters, and the average duration of catheterisation rarely exceeds three to four days, many hospitals no longer routinely replace catheters at defined intervals.

In the linked study, Webster and colleagues report a large randomised controlled trial of different methods of managing peripheral intravenous catheters,¹¹ following an earlier pilot study,¹² seeking scientific validation that peripheral venous catheters no longer need to be replaced at least every 72 hours. In total, 755 medical and surgical patients were randomised to have their peripheral intravenous catheter routinely replaced every three days (control group) or only when clinically indicated, for phlebitis, infiltration, or unexplained fever (clinically indicated group). The study found no significant difference between the groups in premature removal of catheters for phlebitis or infiltration (relative risk 1.15, 95% confidence interval 0.95 to 1.40). The authors estimate that peripheral infusion related costs could be reduced by about 25% if hospitals replaced catheters only when clinically indicated, rather than at 48-72 hour intervals. However, they conclude that larger trials are needed to support this policy if phlebitis is used as the primary endpoint.

Considering that nearly 200 million peripheral intravenous catheters are used each year in US hospitals alone,⁹ Webster and colleagues' trial is important. A limitation of the trial, however, was that the nurses who provided clinical care assessed the insertion sites when the catheters were removed, rather than researchers. Moreover, the study was done in a hospital with a dedicated nurse intravenous therapy team. The incidence of all complications, especially phlebitis, is greatly reduced when

highly experienced nurses insert the catheters and provide follow-up monitoring and care,⁷ which may explain why the incidence of phlebitis in the trial was low.

As a consequence, it is unclear how well the results can be generalised to the majority of hospitals, which do *not* have intravenous teams. The study is also underpowered to reliably conclude that abandoning periodic replacement of peripheral intravenous catheters is unlikely to increase the incidence of infiltration or phlebitis. In addition, the cost analyses did not include the estimated costs of treating severe phlebitis and infiltration or the rare cases of peripheral intravenous catheter related bacteraemia (about 1-3/1000 catheters^{7 9 10}) that will certainly occur, mostly after 48 hours of catheterisation.^{4 7 10} Finally, large cohort studies show that the risk of intravenous phlebitis rises significantly after 48 hours not 72 hours,^{4 6 7} and a large randomised trial comparing routine peripheral intravenous catheter replacements at 48 hours with replacing catheters only when clinically indicated might well show a significant reduction in phlebitis and costs with routine replacement.

Large randomised controlled trials have shown that using specialised teams to insert and care for all peripheral intravenous catheters,⁷ or adopting simple and relatively inexpensive technological advances—such as using in-line filters to remove microparticulates within the infusate,¹³ using catheters made of polyetherurethane rather than Teflon,^{5 6} and securing catheters with a new tapeless device¹⁴—each substantially reduced the incidence of infusion phlebitis and was cost effective. Specialised teams also prevented peripheral intravenous catheter related bacteraemia.⁷ Such approaches could potentially obviate the need to replace peripheral intravenous catheters at periodic intervals.

In summary, Webster and colleagues' trial did not satisfactorily prove that not replacing peripheral intravenous catheters at 72 hour intervals is safe and cost effective, especially in hospitals that do not have specialised intravenous teams to insert and care for catheters, and the value of periodic catheter replacement remains unresolved. Although abandoning scheduled replacements may not greatly increase the incidence of infusion phlebitis and infiltration in the average hospital that currently replaces peripheral catheters at 72 hour intervals, it would probably increase the risk of catheter related bacteraemia with *Staphylococcus*

aureus.^{4 7 10} A large well designed randomised trial comparing replacements at 48 hour intervals with replacement only as clinically indicated would be likely to show significantly fewer local complications and, if the study was adequately powered, a reduced incidence of catheter related bacteraemia as well. But having a specialised team insert and care for catheters clearly reduces both phlebitis and bacteraemia. Moreover, adopting one of the technologies described can also reduce infusion phlebitis and be cost effective, whether or not the hospital replaces peripheral catheters at scheduled intervals.

- 1 British Medical Association: *Healthcare associated infections: a guide for healthcare professionals*. 2006. www.bma.org.uk/ap.nsf/Content/HealthcareAssocInfect.
- 2 Australian Department of Health and Ageing. *Infection control guidelines for the prevention of transmission of infectious diseases in the health care setting*. 2004. [www.safetyandquality.gov.au/internet/safety/publishing.nsf/Content/966A5A0D8A1E5C46CA2571D80021E034/\\$File/intravascdevicejun05.pdf](http://www.safetyandquality.gov.au/internet/safety/publishing.nsf/Content/966A5A0D8A1E5C46CA2571D80021E034/$File/intravascdevicejun05.pdf).
- 3 O'Grady NP, Alexander M, Dellinger EP, Gerberding JL, Heard SO, Maki DG, et al. Guidelines for the prevention of intravascular catheter-related infections. *Infect Control Hosp Epidemiol* 2002;23:759-69.
- 4 Tager IB, Ginsberg MB, Ellis SE, Walsh NE, Dupont I, Simchen E, et al. An epidemiologic study of the risks associated with peripheral intravenous catheters. *Am J Epidemiol* 1983;118:839-51.
- 5 Gaukroger PB, Roberts JG, Mannes TA. Infusion thrombophlebitis:

a prospective comparison of 645 Vialon and Teflon cannulae in anaesthetic and postoperative use. *Anaesth Intensive Care* 1988;16:265-71.

- 6 Maki DG, Ringer M. Risk factors for infusion-related phlebitis with small peripheral venous catheters. A randomized controlled trial. *Ann Intern Med* 1991;114:845-54.
- 7 Soifer NE, Borzak S, Edlin BR, Weinstein RA. Prevention of peripheral venous catheter complications with an intravenous therapy team: a randomized controlled trial. *Arch Intern Med* 1998;158:473-7.
- 8 Bregenzer T, Conen D, Sakmann P, Widmer AF. Is routine replacement of peripheral intravenous catheters necessary? *Arch Intern Med* 1998;158:151-6.
- 9 Maki DG, Kluger DM, Crnich CJ. The risk of bloodstream infection in adults with different intravascular devices: a systematic review of 200 published prospective studies. *Mayo Clin Proc* 2006;81:1159-71.
- 10 Collignon PJ. Intravascular catheter associated sepsis: a common problem. The Australian study on intravascular catheter associated sepsis. *Med J Aust* 1994;161:374-8.
- 11 Webster J, Clarke S, Paterson D, Hutton A, van Dyk S, Gale C, et al. Routine care of peripheral intravenous catheters versus clinically indicated replacement: randomised controlled trial. *BMJ* 2008;337:a630.
- 12 Webster J, Lloyd S, Hopkins T, Osborne S, Yaxley M. Developing a research base for intravenous peripheral cannula re-sites (DRIP trial). A randomised controlled trial of hospital in-patients. *Int J Nurs Stud* 2007;44:664-71.
- 13 Falchuk KH, Peterson L, McNeil BJ. Microparticulate-induced phlebitis: its prevention by in-line filtration. *N Engl J Med* 1985;312:78-82.
- 14 Schears G. Summary of product trials for 10,164 patients: comparing an intravenous stabilizing device to tape. *J Infusion Nurs* 2006;29:225-31.

Bullous pemphigoid and pemphigus vulgaris

Increasing incidence in elderly people in the UK

RESEARCH, p 160

Pascal Joly professor,
Department of Dermatology,
National Reference Centre for
Autoimmune Bullous Dermatoses,
Rouen University Hospital, 76031
Rouen, France

Pascal-Joly@chu-rouen.fr

Competing interests: None
declared.

Provenance and peer review:
Commissioned; not externally
peer reviewed.

Cite this as: *BMJ* 2008;337:a209
doi: 10.1136/bmj.a209

In the linked study, Langan and colleagues determine the incidence of and mortality from bullous pemphigoid and pemphigus vulgaris in the United Kingdom.¹ Autoimmune bullous diseases of the skin are potentially life threatening disorders. Autoantibodies are directed against adhesion proteins of the keratinocyte membrane (pemphigus) or proteins of the dermal epidermal junction (bullous pemphigoid). Binding of these antibodies to their target antigens disrupts intraepidermal or dermal epidermal junctions—this often results in the formation of cutaneous or mucosal blisters, which evolve into erosions. Extensive erosion can lead to failure of the skin's main functions, particularly defence against infection. Treatments aim to block production of pathogenic antibodies by plasma cells using corticosteroids, immunosuppressants, and, more recently, biological agents such as anti-CD20 or intravenous immunoglobulins.²

Bullous pemphigoid is the most common autoimmune bullous skin disease. The incidence has previously been reported as 7-30 cases per million people each year.³ By using the health improvement network, a computerised longitudinal general practice database, Langan and colleagues report an incidence of this disease in the UK of 43 cases per million people each year.¹ Interestingly, the incidence of bullous pemphigoid increased between 1996 and 2001—about 17% for each calendar year, or a 4.8-fold increase over the 11 years. The reasons for this increase are unclear. Possible explanations include the increase in age of the general population; an increase in debilitating

neurological disorders, which are major risk factors for this disease⁴; and the use of drugs such as neuroleptics and spironolactone.

Langan and colleagues confirmed previous studies indicating that the incidence of bullous pemphigoid is high in elderly people—the adjusted incidence was more than 400 cases per million people each year in those over 80. In line with previous European series, the patients in Langan and colleagues' study had a median age of 80, and this has major prognostic implications for the management of these patients.^{5 6} Indeed, a recent study showed that the main predictors of poor prognosis in patients with bullous pemphigoid were demographic factors (older age and female sex), associated medical conditions (cardiac insufficiency, history of stroke, and dementia), and poor general health. No factors directly related to the disease, not even the extent of the cutaneous lesions, were related to patients' prognosis.⁷

Langan and colleagues also confirmed the association between poor prognosis and elderly age. One year mortality in their patients was 12% and 19%, respectively—twice and three times higher than in the general population. Moreover, mortality in patients with bullous pemphigoid was probably underestimated because the information came exclusively from general practitioners. Also, patients admitted to hospital for bullous pemphigoid and those living in nursing homes—patients with the worst prognosis—were excluded.

Although this study did not record the causes of

deaths, two retrospective studies performed in France and Germany indicate that high doses of systemic corticosteroids are responsible for the poor prognosis of patients with bullous pemphigoid.^{6,7} This hypothesis was confirmed by a randomised controlled trial.⁸

Langan and colleagues showed that bullous pemphigoid and pemphigus are more common than many clinicians realise and should no longer be thought of as rare diseases in elderly patients. They also confirm that these diseases have a poor prognosis and that safer treatments are needed to avoid the severe adverse effects of high doses of corticosteroids, especially in elderly patients.

- 1 Langan SM, Smeeth L, Hubbard R, Fleming KM, Smith CJP, West J. Bullous pemphigoid and pemphigus vulgaris—incidence and mortality in the UK: population based cohort study. *BMJ* 2008; doi: 10.1136/bmj.a180.
- 2 Joly P, Mouquet H, Roujeau JC, D'Incan M, Gilbert D, Jacquot S, et al.

A single cycle of rituximab for the treatment of severe pemphigus. *N Engl J Med* 2007;357:545-52.

- 3 Gudi VS, White MI, Cruickshank N, Herriot R, Edwards SL, Nimmo F, et al. Annual incidence and mortality of bullous pemphigoid in the Grampian region of north-east Scotland. *Br J Dermatol* 2005;153:424-7.
- 4 Cordel N, Chosidow O, Hellot MF, Delaporte E, Lok C, Vaillant L, et al. Neurological disorders in patients with bullous pemphigoid. *Dermatology* 2007;215:187-91.
- 5 Joly P, Roujeau JC, Benichou J, Picard C, Dreno B, Delaporte E, et al. A comparison of oral and topical corticosteroids in patients with bullous pemphigoid. *N Engl J Med* 2002;346:321-7.
- 6 Rzany B, Partscht K, Jung M, Kippes W, Mecking D, Baima B, et al. Risk factors for lethal outcome in patients with bullous pemphigoid: low serum albumin level, high dosage of glucocorticosteroids, and old age. *Arch Dermatol* 2002;138:903-8.
- 7 Joly P, Benichou J, Lok C, Hellot MF, Saiag P, Tancrede-Bohin E, et al. Prediction of survival for patients with bullous pemphigoid: a prospective study. *Arch Dermatol* 2005;141:691-8.
- 8 Roujeau JC, Lok C, Bastuji-Garin S, Mhalla S, Enginger V, Bernard P. High risk of death in elderly patients with extensive bullous pemphigoid. *Arch Dermatol* 1998;134:465-9.

Cross border health care in Europe

European Commission provides legal clarity and more information

Martin McKee professor of European public health, London School of Hygiene and Tropical Medicine, London WC1E 7HT
martin.mckee@lshtm.ac.uk

Paul Belcher EU government affairs adviser, Royal College of Physicians, London NW1 4LE

Competing interests: MM participated in the impact assessment of the EU's proposals and coauthored a working paper on quality of health care in Europe. This editorial draws on research conducted in the Europe for patients project, supported by the European Commission's sixth framework programme.

Provenance and peer review: Commissioned based on an idea from the author; not externally peer reviewed.

Cite this as: *BMJ* 2008;337:a610
doi: 10.1136/bmj.39398.456493.80

This week the European Commission publishes its long delayed proposals for how citizens of the European Union should obtain health care in other member states. These proposals will be considered by the European Parliament and governments of the member states.¹ The proposals build on a process involving health ministries and other stakeholders that began in 2003.

Current arrangements for cross border health care were established in the 1970s, when it became clear that free movement within Europe would require that people could receive health care when abroad. It was also recognised that people might need to be sent abroad for treatment, but that this should be controlled by the organisations paying for the care.²

The number of people crossing European borders has increased exponentially.³ A new generation of Europeans sees national frontiers as increasingly irrelevant. Some of them have challenged what they see as unjustifiable restrictions on their right to obtain health care in another country and, in many cases, their arguments have been upheld by the European Court of Justice.⁴ This has resulted in a legislative framework regulating cross border care that is full of holes created by legal precedents, but with little clarity about what those precedents mean in practice. It was never going to be easy to resolve this confused situation, and a recent attempt to treat health services like any other service foundered when the many specificities of health care became clear.

The commission's proposals are incremental. They maintain provisions already in place and focus on areas where clarity is needed. They have three main strands.

Firstly, the values (universality, access to good quality care, equity, and solidarity) and principles (quality, safety, care based on evidence and ethics,

patient involvement, redress, and privacy and confidentiality) underpinning European health systems are clearly stated. These were previously agreed by Europe's health ministries.⁵ Responsibility for ensuring that services comply with these values and principles should lie with the member state on whose territory the care is provided, although subject to future agreement on mechanisms to ensure that core values and principles are adhered to everywhere.

Secondly, a specific framework will be introduced for the aspects of cross border care not already covered by existing legislation, such as that covering people who fall ill while temporarily abroad. Key elements relate to people who choose to go abroad to obtain care. If this is non-hospital care, they simply arrange it themselves. If it is hospital or other specialised care, which the commission will define, countries may introduce systems that require people to seek prior authorisation before obtaining care abroad, but only if they can show that this is necessary to prevent outflow of patients from making their hospitals non-viable. Refusals must be limited to those necessary to avoid such adverse effects on their existing hospital system. Where the home country has a system of primary care gate keeping, this will be respected. In both cases, patients will be entitled to reimbursement only up to the amount that would be paid at home and will not be allowed to make a profit. This framework also proposes a network of national contact points for patients seeking information and tidies up several unresolved matters. For example, it will make it clear that pharmacies should honour prescriptions issued by doctors in other member states, subject to necessary checks and with certain drugs excluded.

Thirdly, mechanisms will be established to foster European collaboration on health services, such as

shared facilities in border areas, common methods of technology assessment, and centres of excellence for rare conditions.

The proposals provide much needed legal clarity in many areas and should provide patients with more information than they have at present. Yet the proposals also raise new problems that may need to be tested in the courts. The most controversial of these may be how member states apply the agreed values and principles. At the very least, the proposals will probably lead to a wide ranging debate on the considerable cultural differences seen in European health systems.

Patients will not be able to demand procedures abroad that are not authorised at home, although their home system can use existing mechanisms to access such procedures if thought to be appropriate. The phenomenon of “postcode prescribing”—where drugs are available to patients in one primary care trust but not another—has already been tested in the English courts.⁶ In future, such cases may have a European dimension as European citizens become more aware of national differences in entitlements.

The proposed legal instrument is a framework directive. This establishes the principles underlying subsequent legislation and sets the broad parameters within which it can operate. However, it leaves flexibility to respond to specific problems and changing circumstances. This will be necessary given the extensive agenda ahead. In particular, the proposals leave responsibility for quality and

safety with individual member states, even though some do not yet have effective systems in place, but it does hint at future European action to facilitate improvements.⁷ The need to deal with the compatibility of mechanisms for maintaining professional competence is particularly pressing, given the more mobile workforce.^{8,9}

The commission's proposals should be seen not as a route map for patients' mobility in Europe, but rather as a set of general directions, which would leave governments and other stakeholders to fill in the details.

- 1 European Commission. *Proposal for a directive of the European Parliament and of the council on the application of patients' rights in cross-border healthcare*. Brussels: European Commission, 2 July 2008.
- 2 Council of the European Union. *Regulation (EEC) no 1408/71 of the council of 14 June 1971 on the application of social security schemes to employed persons and their families moving within the community*. Brussels: Council of the European Union, 1971.
- 3 Rosenmöller M, McKee M, Baeten R. *Patient mobility in the European Union: learning from experience*. London: European Observatory on Health Care Systems, 2006. www.euro.who.int/observatory/Publications/20060522_4.
- 4 Kanavos PG, McKee M. Cross-border issues in the provision of health services: Are we moving towards a European health policy? *J Health Serv Res Pol* 2000;5:231-6.
- 5 Council of the European Union. *Council conclusions on common values and principles in EU health systems*. Luxembourg: Council of the European Union, 1-2 June 2006.
- 6 White C. Final guidance issued on Herceptin after appeal rejected. *BMJ* 2006;333:409.
- 7 Legido-Quigley H, McKee M, Nolte E. *Assuring the quality of health care in the European Union*. London: European Observatory on Health Care Systems (in press).
- 8 Merkur S, Mossialos E, Long M, McKee M. Physician Revalidation in Europe. *Clin Med* (in press).
- 9 McKee M, Belcher P, Coulter A, Raje D. *Consultation regarding community action on health services*. London: Royal College of Physicians, 2007.

The future of primary and community care in England

The permanent NHS revolution reaches into general practice

Chris Ham professor of health policy and management, Policy and Management, Health Services Management Centre, University of Birmingham, Birmingham B15 2RT c.j.ham@bham.ac.uk

Competing interests: CH was director of the strategy unit in the Department of Health between 2000 and 2004.

Provenance and peer review: Commissioned, not externally peer reviewed.

Cite this as: *BMJ* 2008;337:a819
doi: 10.1136/bmj.a819

Starting with the bold statement that “primary and community care services are regarded with pride at home and admiration abroad,”¹ Lord Darzi's plans for the future extend the process of permanent NHS revolution into the heart of general practice.

The government's strategy for primary and community care in England acknowledges that many improvements have occurred in the past decade. Despite these achievements, the standard of primary care varies widely and needs to adapt to rising public expectations, changes in risk factors, and an ageing population. Accordingly, the strategy outlines proposals for reform in three areas.

Firstly, it argues that there is a need to promote healthy lives through a greater emphasis on prevention. The main initiative in this area is a vascular risk assessment programme for people aged 40 to 74 to be delivered by general practitioners, pharmacies, and other services, as foreshadowed in the prime minister's first major speech on the NHS in January 2008.²

Secondly, the strategy sets out proposals for continuously improving the quality of care. This is

to be done through refining the quality and outcomes framework to focus more on health outcomes rather than process measures, and by collecting and publishing data on service quality, including outcomes reported by patients. The Department of Health will also work with the Royal College of General Practitioners to develop an accreditation scheme for general practices.

Thirdly, and most radically, the strategy includes plans to increase choice and give patients a greater role in shaping services. These plans include the continuing development of health centres led by general practitioners and offering walk-in services and bookable appointments, the use of care plans for anyone with a chronic disease who wants them, and the piloting of individual budgets. Individual budgets presage the introduction of patient based commissioning, giving service users direct control over resources.

Absent from the government's plans is any mention of polyclinics. Although the idea of patients having access to a wider range of services in the

community is commended in the strategy, it places the emphasis on providing these services in different settings rather than in a single building. Whether this amounts to a change of heart or a change of presentation is unclear.

Underpinning the government's policy proposals is a series of ideas on how they can be implemented. As in the main report on the NHS Next Stage Review,³ the emphasis is on change being driven locally and by clinicians. To this end, the strategy makes a bold commitment to "redefine and reinvigorate practice based commissioning" in the process acknowledging, if only implicitly, that practice based commissioning has so far failed to deliver.

In seeking to reinvigorate practice based commissioning, it proposes incentives for a broader range of clinicians, including hospital based specialists, to become involved. This is intended to support the development of more integrated care for patients. In an ambitious and welcome extension of this principle, the strategy includes a commitment to test and evaluate how primary and community clinicians can improve health outcomes through a pilot programme of integrated care organisations.

The theme of integration recurs in the discussion of the relationship between health and social care, although in this context it seems to be more of an afterthought than a well considered policy. Lacking specific proposals, the document expresses a commitment to support organisations that wish to go further in integrating health and social care services, effectively kicking this idea into the long grass by announcing the setting up of a group, chaired by the health minister, to work with stakeholders to decide how this should be done.

The strategy throws down the gauntlet to community health services by signalling the intention to focus on ways of improving productivity and

quality in these services. This includes developing measures of performance that can be used by commissioners and providers to secure better value for money. Community health services could undergo substantial organisational change as primary care trusts divest themselves of their responsibilities to provide services in a rerun of Nigel Crisp's ill fated plans from 2005.⁴

In general practice, patient choice and competition between providers will be driven by a fairer system of funding in which money will follow patients' choices. This sounds the death knell for the guarantee of a minimum income for practices and the protection it has offered some practices since the introduction of the new contract. The impact of this commitment will depend on how and over what period it is implemented, and the willingness of the BMA to treat with health ministers, with whom they have not enjoyed the best of relations.

There is much common ground between the strategy and the government's previous vision for community services, *Our Health, Our Care, Our Say*, published in 2006.⁵ The need to reiterate earlier policy commitments suggests that the biggest challenge in taking forward the vision rests in the means of implementation and whether clinical leadership and money following patients in primary care will be fit for purpose.⁶

Lord Darzi believes that an appeal to professionalism is the best way to advance consumerism and implementation of the proposals in his strategy. This may be what the profession wants to hear,⁷ but sits uneasily alongside the claims of Ben Bradshaw, his fellow health minister, that GPs operate cartels that hinder patient choice. With Darzi cast as the good cop and Bradshaw the bad cop, there is a risk of mixed messages from the government.

Whether the strategy will help to consolidate domestic pride in and international admiration of primary care, or whether it will undermine the fragile flower that is British general practice, remains to be seen. The strategy is good in parts, and in many areas there is the prospect that it will strengthen a system that is already functioning well. Much now hinges on how it is taken forward and the skill with which health ministers and the medical profession navigate the treacherous waters that lie ahead.

- 1 Department of Health. *NHS next stage review: our vision for primary and community care*. July 2008. www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_085937
- 2 Ham C. Gordon Brown's agenda for the NHS. *BMJ* 2008;336:53-4
- 3 Secretary of State for Health. *High quality care for all*. June 2008. (Cm 7432.) www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_085825
- 4 Timmins N. How the mighty came to fall. *BMJ* 2006;332:628.
- 5 Secretary of State for Health. *Our health, our care, our say*. January 2006. www.dh.gov.uk/en/Healthcare/Ourhealthourcareoursay/index.htm
- 6 Bevan H, Ham C, Plsek PE. *The next leg of the journey. How do we make High Quality Care for All a reality?* London: NHS Institute for Innovation and Improvement, 2008. www.institute.nhs.uk/images//documents/About_US/nextlegofthejourney_bevan-ham-plsek.pdf
- 7 Horton R. The Darzi vision: quality, engagement and professionalism. *Lancet* 2008;372:3-4.

Lord Darzi (left) believes an appeal to professionalism will achieve his goals. This contrasts with Ben Bradshaw's claim that GPs operate cartels that hinder patient choice