and tenderness in the right hypochondrium, with pain in the right shoulder; appetite bad; foul tongue and great thirst; complains of a sense of distension after meals, with flatulence; bowels costive; urine scanty and high coloured, with œdema of the legs and ankles, and swelling and fluctuation of the abdomen.

R. Pil. Hydrarg., gr. v.; Extr. Coloc. Co., gr. vj. M. Fiat pil. ij., hora somni sumend. R. Haust. Salin. Aper., primo mane sumend.

9th. Bowels well acted upon, the evacuations being of a clay colour; other symptoms about the same.

R. Pil. Hydrarg., gr. iij.; Pulv. Scillæ, Digitalis, utr., gr. j. M. Fiat pilula, mane et vespere sumenda.

R. Potass. Iodid., scr. ij.; Tinct. Hyoscyami, dr. ij.; Infusi Cascariæ, oz. vj.; M. Capt. cochl. tria ter indies.

11th. Legs and ankles considerably œdematous; more swelling of abdomen; breathing difficult in the recumbent posture; great thirst; urine still scanty and high coloured.

To continue the pills. To omit the former mixture and to take the following:—R. Potassæ Bicarbon., dr. j.; Infusi Buchu, oz. vj. M. Capt. cochl. ampla iij., quarta quaque hora. To drink broom-tea, and to have an ointment of tartar emetic and iodide of potassium rubbed into the right side.

13th. Symptoms alleviated; has passed more urine; slept better; thirst diminished; bowels confined.

To continue the medicines; to have an aperient draught containing one scruple of Pulvis Jalapæ Comp.

17th. Greatly improved; has passed large quantities of urine; the skin from being dry and harsh, has become cool and moist; tongue clean; no œdema of the legs and ankles, and decidedly less swelling of the abdomen; is able to lie down comfortably; sleeps much better; appetite improved.

To omit the blue pill and squill, and to have—R. Pil. Plummerii, gr. iv.; Extr. Hyoscyami, gr. v. M. Fiat pil. ij., hora somni omni nocte sumend. To continue the mixture.

19th. From this time she began to amend rapidly. Has been latterly much purged, whether from the buchu or not I am unable to say; the evacuations the same as from the use of hydragogues; the kidneys have been powerfully acted upon, and she has continued to pass large quantities of urine. On the 24th there was no swelling of the abdomen whatever, and on the 26th, she was so far convalescent as to be enabled to go into the country.

CASE II.

M. R., aged 14. This was a case of anasarca, succeeding about the usual period, a mild attack of scarlatina. It resisted the treatment which I adopted in the first instance, of bark and aromatic confection, and the symptoms became aggravated, and on November 18th, there was great thirst, quick pulse, white tongue, and constipated bowels, the dropsy in this case, as Dr. Gregory states, depending upon a continuance of a highly excited state of the cutaneous capillaries, which, earlier in the disease, occasions desquamation of the cuticle. After the

exhibitions of saline purgatives, combined with small doses of digitalis, I again tried the buchu, with the same good effects as before, and in a few days the dropsy disappeared. In this case also, there was great purging, when the buchu had been taken for a short time, and also an abundant secretion of urine.

SHORT NOTES OF THE OPINIONS AND PRACTICE OF THE LATE JOHN PEARSON, ESQ., F.R.S., ON SYPHILITIC DISEASE.*

By W. S. OKE, M.D., Physician to the Royal South Hants Infirmary.

Ulcers on the Nipple, Navel, Alæ Nasi, and Upper Lip.—Ulcerations of the areola and nipple in both sexes, particularly in women, are very tender to the touch, and discharge a thin ichorous fluid. When of long continuance they occasion inflammation and abscess of the breast. They occur chiefly in pregnant women who have lues, but they occur so rarely as not to be seen perhaps in more than one case out of 500. In the umbilicus also ulcers are more frequent in women than in men, and are very offensive.

When the alæ nasi and upper lip are affected by venereal ulcer, a pimple will form, and discharge an ichor; or there may be soreness and excoriation produced, which if suffered to go on will destroy the alæ. The sore always spreads from the angle between the cheek and nose, and frequently degenerates into a painful spreading ulcer, which will sometimes extend to, and destroy, the upper lip. These cases are not common. Cancer generally affects the lower lip; but the secondary sore from syphilitic poison usually attacks the upper lip.

Local Treatment of Ulcers of the Nipple, Umbilicus, Alæ Nasi, and Upper Lip.—In the treatment of the two first, cleanliness must be insisted on. Where the nipple is sore and inflamed, a poultice, made with linseed meal and Goulard water, may be applied till the swelling is reduced; afterwards a solution of the Argenti Nitras, or of the bi-chloride of mercury, may be used. The wash, commonly called "the black wash," may be tried; but no greasy application should be made use of. Sometimes these sores will not heal till after delivery.

The same mode of treatment will be applicable to ulcers on the umbilicus. Ulcers on the alæ nasi and upper lip may be dressed with the Unguentum Hydrargyri Nitrico-Oxidi, and occasionally touched with a solution of the nitrate of silver.

[Case.—A female, of middle age, consulted me for an ulcer on the left alæ nasi. It occupied the greater part of the alæ, which was much thickened. It was foul, with high jagged edges, and had all the appearance of a carcinomatous character. It produced great deformity of the countenance, had continued nearly twelve

months, and had resisted every means that had been tried to heal it. I at first doubted whether I should do her any good, but having been consulted by her some years previously for ulceration of the throat, I thought it probable that it might have a syphilitic origin. Accordingly, I prescribed five grains of the *Pilula Hydrargyri* every night, and four grains of the iodide of potassium three times a day, desiring that the ulcer might be washed twice a day with a little diluted tincture of myrrh. Under this treatment the sore completely healed in a month, and all hardness of the ala disappeared.]

Venereal Sarcocoele.—This is a tumefaction of the testicle, and by no means a common occurrence. When it does occur, it is always connected with an affection of the skin. Both testicles are usually enlarged, and become soft and pulpy. There is also a fulness of the integuments, which are sometimes covered with tubercles or a scaly furfuraceous appearance. The swelling is pretty uniform, and there is not much pain nor any affection of the spermatic vessels. Suppuration of the scrotum rarely occurs from the syphilitic poison.

Syphilitic Affection of the Ears.—The sebaceous glands behind the ears, and upon the concha, and sometimes the whole cartilaginous substance, forming the external ear, are involved in the syphilitic action. The parts become thickened and painful, often discharging offensive matter; at other times the disease will extend to the Eustachian tubes, particularly where the tonsils and fauces are affected, producing tinnitus aurium. The ossicula auditus are not so apt to be involved as in scrofula, and therefore deafness is probably the result of thickening of the parts. From this patients generally recover.

Intertrigo Syphilitica.—By intertrigo is meant a painful excoriation of the integuments, with an offensive ichorous discharge, often extending from the anterior to the posterior perineum; and on the inside of the upper part of the thigh there is sometimes a morbid surface, having a whitish appearance, and covered with a thick sebaceous matter. The patient is disturbed by it at night, and cannot sleep. The ulceration never becomes deep, and is seldom met with in the higher ranks of life. It is more the result of a want of cleanliness than a characteristic of lues, although it occurs in those affected with this disease. Intertrigo pudendi is often the result of gonorrhœa in women; it may also be produced by the action of mercury, when it looks as if the epithelium had been separated from the organs of generation.

Treatment of Venereal Sarcocoele in Syphilitic Affection of the Ears and Intertrigo.—The swelling of the testicle will generally begin to subside when the mercury acts on the constitution; but if the disease proves obstinate, a warm solution of the muriate of ammonia, with vinegar, may be locally applied, or the camphorated spirits of wine, with a little of the Liquor

Ammoniae, made warm over a candle. Should the swelling of the testicle remain in an indolent state, a poultice of the fresh leaves of hemlock may prove useful.

Ulcerations of the ears should be well washed with a decoction of poppies and chamomile flowers, and dressed with the *Unguentum Hydrargyri Nitrico-Oxidi*. If the ulceration extend to the internal ear, fumigation of cinnabar will be useful.

Intertrigo seldom demands much attention from the surgeon, and may be removed by interposing a silk handkerchief between the scrotum and thigh, supposing mercury to be always used. Intertrigo pudendi may be treated with Goulard water and lime water, to which a little spirit is added, or with a weak solution of the bichloride of mercury. A strong infusion of tea or chamomile flowers will sometimes relieve the symptoms. Should there be much excoriation, the saturnine ointment or calamine cerate may be applied.

Intertrigo ani is by far the most painful, and is frequently so severe as not to admit of relief. Large doses of opium will be necessary to lessen irritation; and in these cases the form of powders answers better than unguents. Equal parts of powdered myrrh, white lead, and *Lapis Calaminaris*, may be tried, or the sulphate of zinc in powder; and the parts should be frequently washed with a soft sponge wrung out in warm elder flower water, or warm water only.

Warts and Excrecences.—Warts and excrecences may be considered diseases of the skin, and caused by spurious growth. When they remain after the total disappearance of the venereal symptoms, it is probable that in the majority of cases they will not again infect the constitution; there are, however, exceptions to the rule. There are cases where, after the removal of warts from the prepuce, &c, by excision, the sore will not heal, except under a course of mercury. Syphilitic warts may be divided into the *verruca vulgaris*, and *acrochordis*, and syphilitic excrecences into *condylomata*, *fici*, and *cristæ*. The common wart is a small, hard, and almost insensible tubercle, arising from the surface of the skin, covered with a dense cuticle, and of the colour of the integuments. They occur on the penis, nymphæ, perineum, &c., and especially in those who have had the venereal disease. They do not arise from the cuticle, and are always sensible. They are more luxuriant where the epidermis is kept moist by secretion, and are black in the negro. They are not to be considered as newly formed parts, but rather a redundancy of the normal parts, and hence their base does not increase, although the wart itself may acquire a considerable size.

The *acrochord* or pencil-shape kind springs from a moist surface,—viz., from within the labia of females, and is of a softer texture than the common wart; they are sometimes met with within the nose, where they are not always of the syphilitic character.

Condylomata are smooth, soft, and of a dark red colour; they are covered with cuticle, and not much elevated above the skin; their surface is as broad as their base, and they are commonly found about the anus. Sometimes they assume a shape and colour resembling figs, and hence are called figs.

Cristæ occur about the anus, perineum, and glans; they have a broad base, and a fimbriated appearance. When seedy they have been compared to a cauliflower. Sometimes they are produced in great number on the perineum of females, forming a large bulk, and discharging a most offensive ichorous matter. They sometimes occur in children, and without any venereal cause; they are not probably derived from the same sources as the common wart, but from the sebaceous glands of the part.

Treatment of Verruæ, Condylomata, &c.—The hard warts may be destroyed by inflammation, decomposed by caustic, or extirpated by the knife. It is difficult to excite a sufficient inflammation to destroy warts, without injuring the surrounding parts. It is best first to soften them by a poultice, containing soda, after which the top is to be pared off, and a mineral acid applied, till the whole is destroyed. Where a ligature can be used it is a still better remedy, as the wart is then less liable to return. It is always applicable to the pencil kind. It appears that the part of the wart below the ligature dies as well as that above it. If the prepuce be studded with warts, it is better to remove it; and if there be masses of them on the penis, a small portion only should be destroyed at a time, lest the inflammation produce gangrene. Condylomata frequently shrink and disappear during the mercurial course, without any local application; if they should remain, a solution of the bi-chloride of mercury, or of mercury in dilute nitric acid may be used.

Cristæ are to be reckoned amongst the most inveterate venereal excrescences. They will be regenerated again and again. The most useful applications are the following:—A solution of the bi-chloride of mercury, the hydrochlorate of ammonia, powder of savin, acetate of copper, sesqui-chloride of antimony, (butler of antimony,) arsenic and fumigations, with the bi-sulphuret of mercury. Those caustics which produce most inflammation, are the best. Should a crista, after having been destroyed by caustic, sprout up again, great benefit will be derived by a strong solution of alum in a decoction of oak bark. The following is recommended by Plenck:—

R. Alcohol, Aceti Destillati, utrque oz. iss.
Hydrargyri Bi-Chloridi dr. j.
Aluminis Sulphatis, Camphoræ, utrque . dr. ss.
Misce. Fiat Lotio.

Whenever a ligature can be used in cases of crista it ought to be applied. A needle with a double ligature will be found the most effectual method of applying it. If it be thought fit to excise these

excrescences, the excisions must be deep in order that the sebaceous glands, from which they probably spring, may be extirpated with them.

Rhagades.—Rhagades are fissures on the hands, about the anus, or perineum, and sometimes on the lips. It is important to distinguish the syphilitic from the common kind. The common rhagades are dry and give no disturbance but by their dryness; they sometimes appear like little nodes or corns in patients who have a tendency to lepra. Those which are derived from syphilitic causes may be distinguished by their being very painful and discharging profusely a foetid matter.

Treatment of Rhagades.—Those on the hands generally begin to heal when the system is under the influence of mercury, and will get well without any local application, if attention be paid to cleanliness; those on the anus are very painful and difficult to heal. They should be dressed so as to expose the bottom of the sore, and the best application will be the Unguentum Hydrargyri Nitratis, or a solution of the nitrate of silver.

(To be continued.)

ON THE TAXIS.

TO THE EDITOR OF THE PROVINCIAL MEDICAL AND SURGICAL JOURNAL.

SIR,

Some late numbers of your Journal contain communications on the subject of incarcerated hernia, which are calculated to cast imputations on the practice of some of the most eminent surgeons and most popular writers on surgery, from Desault to Druitt.

One of your correspondents, Mr. Hunt, seems fearless in maintaining his position in regard to the general practicability of the reduction by the taxis. He deserves our best acknowledgments, inasmuch as he has insisted on a method which, I verily believe, might have been successfully adopted in many of the cases which have been subjected to the severer operation. For the cause of suffering humanity, therefore, I am happy that I can offer my testimony to the superiority of the mode of proceeding Mr. Hunt has recommended.

If the explanation of the physiological principles on which the misfortune depends be correct,—namely, the congestive state of the veins, then we must admit, consequently, the impossibility of returning the venous blood or other contents of the hernial tumour by a hasty trial of compression. The reduction of strangulated hernia by the taxis is a slow process, but speaking from my own experience, it is always possible. He is wanting in patience or practical tact who does not succeed when called in to a case before acute inflammation has supervened. The directions for the reduction of hernia are usually absurd. To talk of kneading the tumour is mischievous,—of first “pulling out” further the contents is impossible, or if not impossible, only adding to the existing evil. Lastly, to limit the