Published in final edited form as:

Org Lett. 2006 May 11; 8(10): 2175–2178.

Solvolysis of a Tetrahydropyranyl Mesylate: Mechanistic Implications for the Prins Cyclization, 2-Oxonias-Cope Rearrangement, and Grob Fragmentation

Ramesh Jasti and Scott D. Rychnovsky*

Department of Chemistry, 1102 Natural Sciences II, University of California, Irvine, Irvine, California, 92697-2025

Abstract

A solvolysis reaction is used to demonstrate that a tetrahydropyranyl cation is a common intermediate for Prins cyclizations, 2-oxonia Cope rearrangements, and Grob fragmentations of tetrahydropyran rings.

The Prins cyclization and the 2-oxonia-Cope rearrangement are highly stereoselective carbon-carbon bond forming reactions that have received considerable attention from organic chemists.^{1,2,3} Both transformations are initiated by an oxocarbenium ion and a tethered alkene and thus are mechanistically related. Not surprisingly, the two processes are often competing pathways.⁴ Recently, we reported our investigations into the relative rates of 2-oxonia-Cope rearrangements and Prins cyclization reactions.⁵ The mechanistic model set forth in that document was inspired by calculations from Alder suggesting a common cationic intermediate for both transformations.⁶

Figure 1 outlines the proposed mechanistic pathways.⁵ In this model, tetrahydropyranyl cation **3** serves as a branch point in a 2-oxonia-Cope rearrangement (**2** to **4**) and Prins cyclization reaction (**2** to **5**).⁷ Alder suggested that tetrahydropyranyl cation **3** has increased stability due to extensive delocalization. The optimal geometry for this delocalization places the hydrogen in the pseudoaxial position which directs nucleophilic attack along an equatorial trajectory. The extensive delocalization therefore provides rationale for the 2,4,6-cis stereoselectivity commonly observed for Prins cyclization reactions.^{8,9}

The energy diagram in Figure 2 further details the calculations reported by Alder.⁶ Tetrahydropyranyl cation **6** was calculated to have a groundstate energy 13.4 kJ/mol higher in energy than oxocarbenium ion **7**. In addition, the barrier for ring-opening was calculated to be only 1.9 kJ/mol. Based on these calculations, we surmised that generation of a tetrahydropyranyl cation should rapidly lead to an oxocarbenium ion. Such a ring-opening process would represent the first known Grob fragmentation of a tetrahydropyran ring, the reverse of a Prins cyclization. In this way, the delocalized tetrahydropyranyl cation may serve not only as an intermediate for Prins cyclizations and 2-oxonia-Cope rearrangements, but also for Grob fragmentations of tetrahydropyran rings. In this communication, we report a solvolysis reaction linking all three processes to the delocalized tetrahydropyranyl cation intermediate proposed by Alder.

Figure 3 outlines a test to identify a Grob fragmentation pathway for a tetrahydropyran ring. A solvolysis reaction of appropriately substituted tetrahydropyran **8** should generate tetrahydropyranyl cation **9**. Cation **9** could then undergo nucleophilic attack to produce optically active tetrahydropyran **10**. In contrast, tetrahydropyranyl cation **9** may undergo a fast ring-opening process to generate achiral oxocarbenium **11**. Prins cyclization of achiral oxocarbenium ion **11** would then necessarily lead to racemic **10**. Therefore, racemization of tetrahydropyran **10** can be used as an indicator for a Grob fragmentation pathway.

With the racemization test in mind, we turned our attention to the synthesis of optically active tetrahydropyranyl mesylate **17** (Scheme 1). Ester **14** was prepared in 42% yield from optically active alcohol **12**¹⁰ and carboxylic acid **13**.¹¹ Reductive acetylation conditions developed in our laboratory smoothly converted ester **14** to acetoxy ether **15** as a 1:1 mixture of diastereomers.¹² Prins cyclization of **15** followed by methanolysis delivered tetrahydropyran **16** in 67% yield over 2 steps.¹³ Reduction and mesylation completed the synthesis of tetrahydropyran **17**.

Trifluoroacetolysis¹⁴ of optically active tetrahydropyran **17** followed by methanolysis provided racemic tetrahydropyran **18** as a single diastereomer in 82% yield (Scheme 2).¹⁵ Racemization of tetrahydropyran **18** indicates that ring-opening is faster than nucleophilic attack by trifluoroacetate. This result provides the first evidence of a Grob fragmentation pathway for tetrahydropyran rings. One question remaining was whether this was a concerted or a stepwise process. Figure 3 illustrates a Grob fragmentation of **8** occurring by way of discrete carbocation intermediate **9** (i.e. asynchronous Grob fragmentation). Is it possible that solvolysis of **8** in fact leads directly to achiral oxocarbenium ion **11** via a synchronous Grob fragmentation?

In his classic studies, Grob distinguished between asynchronous and synchronous fragmentation reactions by comparing rates of solvolysis with carbon analogs (Figure 4).¹⁶ Cyclohexyltosylate **19** has been shown to have a half-life of 35 min in trifluoroacetic acid at room temperature.¹⁷ In contrast, subjecting tetrahydropyran **20** to similar conditions resulted in no reaction. In a synchronous fragmentation reaction, the heteroatom substituted compound should undergo solvolysis more rapidly or at a similar rate to the carbon analog due to favorable n_o to σ^*_{C-C} interactions. In an asynchronous fragmentation, the heteroatom inductively destabilizes the C4 cationic intermediate, thus slowing the rate of solvolysis relative to the carbon analog. Our results are consistent with an asynchronous fragmentation, in which the inductive effect of the heteroatom decreases the reactivity of tetrahydropyran **20** towards acid-catalyzed solvolysis.

With the results from the solvolysis reactions in hand, a mechanistic scheme can be constructed in which the Prins cyclization, 2-oxonia-Cope rearrangement, and Grob fragmentation all proceed through one common intermediate (Figure 5). Trifluoroacetolysis of mesylate **17** leads to tetrahydropyranyl cation **21**. Tetrahydropyranyl cation **21** can then ring-open, presumably, with equal probability to either oxocarbenium ion **23** or achiral oxocarbenium ion **24**. Tetrahydropyranyl cation **21** therefore is an intermediate for a Grob fragmentation of tetrahydropyran **17**. In addition, oxocarbenium ions **23** and **24** can interconvert by undergoing a ring-closure process followed by a ring-opening. In this way, tetrahydropyranyl cation **21** also serves as an intermediate for a 2-oxonia-Cope rearrangement. Furthermore, the asynchronous nature of the Grob fragmentation implies that the reverse reaction, a Prins cyclization, is also asynchronous. Thus, oxocarbenium ions **23** and **24** lead to **22** by way of the same tetrahydropyranyl cation intermediate **21**. Interestingly, although the Prins cyclization proceeds through a cationic intermediate, only one isomer is observed at C4. This result is consistent with Alder's description of extensive delocalization for tetrahydropyranyl cations such as **21**.^{6,5}

Ring-opening of tetrahydropyranyl cation **21** to achiral oxocarbenium **24** was assumed to be the only pathway for racemization. In order to test the validity of this assumption, saturated tetrahydropyran **25** was prepared. In this case, an achiral intermediate is not available by a simple Grob fragmentation. Surprisingly, trifluoroacetylation of saturated tetrahydropyran **25** followed by methanolysis led to alcohol **26** in only 49% ee (Scheme 3). The loss in optical activity for **26** indicates the presence of a second racemization pathway that is independent of achiral intermediate **24**. The erosion in enantiomeric excess for this reaction may involve isomerization of an (*E*)-oxocarbenium ion to a (*Z*)-oxocarbenium ion followed by a 2-oxonia-Cope rearrangement.¹⁸ Efforts to unravel this unusual racemization pathway are currently underway in our laboratory and will be reported in due course.

In conclusion, we have demonstrated a novel solvolysis reaction of a tetrahydropyranyl mesylate. The outcome of this solvolysis reaction correlates well with a mechanistic model in which a delocalized tetrahydropyranyl cation serves as an intermediate for Prins cyclizations, 2-oxonia-Cope rearrangements, and Grob fragmentations of tetrahydropyran rings. This knowledge should be useful for further development and application of these transformations.

Supporting Information Available

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the National Cancer Institute (CA-081635).

References

1. For reviews on the Prins cyclization see: (a) Arundale E, Mikeska LA. *Chem Rev* 1952;52:505–555. (b) Adams DR, Bhatnagar SP. *Synthesis* 1977:661–672. Snider, BB. *The Prins Reaction and Carbonyl Ene Reactions*. Trost, BM.; Fleming, I.; Heathcock, CH., editors. 2. Pergamon Press; New York: 1991. p. 527–561. (d) Overman LE, Pennington LD. *J Org Chem* 2003;68:7143–7157. [PubMed: 12968864]
2. For recent work on the Prins cyclization see: (a) Bolla ML, Patterson B, Rychnovsky SD. *J Am Chem Soc* 2005;127:16044–16045. [PubMed: 16287289] (b) Dalgard JE, Rychnovsky SD. *J Am Chem Soc* 2004;126:15662–15663. [PubMed: 15571386] (c) Yadav KV, Kumar NV. *J Am Chem Soc* 2004;126:8652–8653. [PubMed: 15250708] (d) Hart DJ, Bennet CE. *Org Lett* 2003;5:1499–1502. [PubMed: 12713308] (e) Barry CSJ, Crosby SR, Harding JR, Huges RA, King CD, Parker GD, Willis CL. *Org Lett* 2003;5:2429–2432. [PubMed: 12841747] (f) Miranda PO, Diaz DD, Padron JI, Bermejo J, Martin VS. *Org Lett* 2003:1979–1982. [PubMed: 12762701]
3. For recent work on the 2-oxonia-Cope see: (a) Nokami J, Yoshizane K, Matsuura H, Sumida SI. *J Am Chem Soc* 1998;120:6609. (b) Sumida SI, Ohga M, Mitani J, Nokami J. *J Am Chem Soc* 2000;122:1310. (c) Loh TP, Hu QY, Ma LT. *J Am Chem Soc* 2001;123:2450. [PubMed: 11456905] (d) Loh TP, Ken Lee CL, Tan KT. *Org Lett* 2002;4:2985–2987. [PubMed: 12182605] (e) Loh TP, Hu QY, Ma LT. *Org Lett* 2002;4:2389–2391. [PubMed: 12098254] (f) Lee CLK, Lee CHA, Tan KT, Loh TP. *Org Lett* 2004;6:1281–1283. [PubMed: 15070317]
4. (a) Lolkema LDM, Semeyn C, Ashek L, Hiemstra H, Speckamp WN. *Tetrahedron* 1994;50:7129–7140. (b) Roush WR, Dille GJ. *Synlett* 2001:955–959. (c) Rychnovsky SD, Marumoto S, Jaber JJ. *Org Lett* 2001;3:3815–3818. [PubMed: 11700146] (d) Crosby SR, Harding JR, King CD, Parker GD, Willis CL. *Org Lett* 2002;4:577–580. [PubMed: 11843595] (e) Crosby SR, Harding JR, King CD, Parker GD, Willis CL. *Org Lett* 2002;4:3407–3410. [PubMed: 12323030] (f) Hart DJ, Bennet CE. *Org Lett* 2003;5:1499–1502. [PubMed: 12713308]
5. Jasti R, Anderson CD, Rychnovsky SD. *J Am Chem Soc* 2005;127:9939–9945. [PubMed: 15998101]
6. Alder RW, Harvey JN, Oakley MT. *J Am Chem Soc* 2002;124:4960–4961. [PubMed: 11982351]
7. For trapping experiments of proposed charged intermediates in Prins cyclizations and 2-oxonia Cope rearrangements see: (a) Barry CS, Bushby N, Harding JR, Hughes RA, Parker GD, Roe R, Willis CL.

- Chem Commun 2005;29:3727–3729. (b) Rychnovsky SD, Marumoto S, Jaber JJ. Org Lett 2001;3:3815–3818. [PubMed: 11700146]
8. Under standard Prins cyclization conditions, nucleophilic trapping (**3** to **5**) is not reversible.
9. For axial-selective Prins cyclizations, see:(a) Jasti R, Vitale J, Rychnovsky SD. J Am Chem Soc 2004;126:9904–9905. [PubMed: 15303848] (b) Lolkema LDM, Hiemstra H, Semeyn C, Speckamp WN. Tetrahedron 1994;50:7115–7128. (c) Miles RB, Davis CE, Coates RM. J Org Chem 2006;71:1493–1501. [PubMed: 16468798]
10. Keck GE, Tarbet KH, Geraci LS. J Am Chem Soc 1993;115:8467.
11. Davies HML, Boebel TA. Tetrahedron Lett 2000;41:8189–8192.
12. (a) Dahanukar VH, Rychnovsky SD. J Org Chem 1996;61:8317–8320. [PubMed: 11667826] (b) Kopecky DJ, Rychnovsky SD. J Org Chem 2000;65:191–198. [PubMed: 10813915]
13. Barry CSJ, Crosby SR, Harding JR, Huges RA, King CD, Parker GD, Willis CL. Org Lett 2003;5:2429–2432. [PubMed: 12841747]
14. Trifluoroacetic acid has been shown to have an exceptional combination of high solvolytic power and low nucleophilicity. See the following and references therein:Nordlander JE, Gruetzmacher RR, Kelly WJ, Jindal SP. J Am Chem Soc 1974;96:181–185.
15. Nucleophilic attack by trifluoroacetate is not reversible. Subjecting optically active tetrahydropyranyl trifluoroacetate to solvolysis conditions results in no reaction.
16. Grob CA. Angew Chem, Int Ed Engl 1969;8:535–546.
17. Lambert JB, Putz GJ. J Am Chem Soc 1973;95:6313–6319.
18. For similar racemization pathways in iminium ion cyclizations see:(a) Daub GW, Heerding DA, Overman LE. Tetrahedron 1988;44:3919–3930. (b) Castro P, Overman LE. Tetrahedron Lett 1993;34:5243–5246.

Figure 1.
Stepwise Prins Cyclization Reaction and 2-Oxonio-Cope Rearrangement

Figure 2.
Energetics of a Tetrahydropyranyl Cation

Figure 3.
Racemization as an Indicator for Grob Fragmentation

Figure 4.
Trifluoroacetolysis of Carbon Analog **19** and Tetrahydropyran **20**

Figure 5. Tetrahydropyranyl Cation **21** as a Common Intermediate for a Prins Cyclization, a 2-Oxonia-Cope Rearrangement, and a Grob Fragmentation

Scheme 1.
Synthesis of Mesylate **17**

Scheme 2.
Trifluoroacetolysis of Mesylate **17**

Scheme 3.
Trifluoroacetytolysis of Mesylate **25**