was voided with extreme pain and increased frequency. At length the flow was almost continual, and the pain so intense as to forbid an attempt at lithotomy; the extraction was, therefore, executed above the pubis. Here, however, a new difficulty arose—the stone, on being seized, was found to have become pulpy, and it escaped from the forceps, with the exception of a nucleus, which was brought away. The remaining portions were then extracted by means of the finger, aided by the *curette* and the dressing forceps.

The softening of this stone seems to have resulted from the highly inflamed bladder and the alkaline secretions, which are usually met with in similar conditions of that viscus. Ammonia dissolves the mucus of the bladder, which matter is the cementing bond of the saline substances which form the stone.

SPONTANEOUS LUXATIONS OF BOTH KNEES.

M. BOUVIER presented a very singular example of lateral incomplete luxation of the tibia, spontaneously effected after paralysis and articular pains. The patient, 60 years of age, is an inmate of the *Salpêtrière*. The two femoro-tibial articulations are constituted solely by the external condyle of the femur and the superior extremity of the tibia. The right knee projects inwardly, so as to form an angle of 150 degrees between the thigh and leg; but on the left side the axis of the leg has not deviated,—the corresponding surfaces of the tibia and femur being obliquely placed in respect to each other, as in the fractures designated by the term *bec-de-flûte*. This remarkable affection has not abolished the motion of the joints, nor has it greatly altered their solidity.

M. GERDY read a new memoir on the laws of vision; but as it contains merely the history of this physiological question, it is not of sufficient importance for analysis here.

GUY'S HOSPITAL REPORTS.

PRACTICAL HINTS ON THE TREATMENT OF STRICTURE OF THE URETHRA.

BY BRANSBY B. COOPER, F.R.S.

Mr. Cooper strongly recommends attention to the treatment of constitutional symptoms in stricture of the urethra, mild instances of which may be removed by general remedies alone, whilst severe cases are protracted if the aid given be solely instrumental. Mr. Cooper agrees with the best surgeons of the day in deprecating the application of force in the passing of instruments, except under certain circumstances, which he specifies, and which are rarely met with.

Strictures have been divided into mixed, permanent, and spasmodic; but Mr. Cooper discredits the existence of the latter, and does not believe the urethra to be muscular, and opposes the opinion of Howship, who asserts he has seen the urethra by its own contractile power dislodge a moderate-sized bougie. Mr. Cooper has watched for some such manifestation of muscular action; but has never observed it, except when the bougie was passed up to the *bulb*, to which we know a muscular apparatus is attached. He is aware that sudden obstruction of the urethra *does* occur, but attributes it to a distension of a portion of the corpus spongiosum, somewhat similar to the venereal intumescence of that structure, but produced by a morbid cause. The mistaken views on this point have been confirmed by the fact, that the same treatment relieves both this affection and muscular spasm; muscular fibre may, however, be affected secondarily, if the distension occurs at the bulb, or in the membranous proportion.

The Causes of Stricture.

The most frequent cause is a morbid action set up in the urethra itself; one of the most frequent kinds of which is inflammation from the long-continuance of gonorrhœa. Stricture of the urethra may also result from disease of the surrounding parts, the prostate rectum or bladder; or it

may be produced by local or general irritation, in which case it may be distinguished by suddenness of obstruction, by its tendency to bleed, and the diathesis of the patient. The most frequent seat of stricture is the most vascular portion of the urethra,—the membranous and bulbous sections of that canal.

Treatment of Stricture.

The sedative plan of treatment is adapted to the irritable stricture,—the warm-bath, opiates, or other narcotics, leeches, belladonna fomentations, and, in case of necessity, caustic bougies. Where there is a disposition to spasm, recourse should be had to bleeding, opium, and belladonna injections. When the stricture from its thickness resists the gentle introduction of bougies, the obstacle may often be overcome by injecting warm water into the urethra, through a canula, to which a syringe is fitted. In the irritable form of stricture, force should *never* be used; for it is frequently destructive of the organization of the urethra, and is sometimes fatal to the patient.

Force is justifiable in a very few cases only; the amount of force cannot be expressed in words, it must be measured by the tact and judgment of the practical surgeon, and should never be carried to an imprudent degree. If there be present severe symptoms of retention, calling for immediate relief, and the catheter can be brought to right angles with the position of the recumbent patient, being then, but not until then, checked in its progress,—it is plain that the stricture is situated in the membranous portion of the urethra, *where the operator may safely use force, judiciously applied*, provided the prostate gland be healthy. The danger which would be incurred in other portions of the urethra, is here prevented by the firm attachment of the canal to the surrounding parts of the deep fascia of the perinæum, and by the guidance of the instrument by the ossa pubis. If the surgeon has, in such cases, carried force as far as he thinks prudent, but has failed to pass the obstacle, he should cut down on the membranous portion of the urethra, for otherwise there is danger of perforating the prostate gland or rectum, or lacerating the canal, in which latter case the stricture is almost certain to recur.

In cutting upon the membranous portion of the urethra, an incision of two inches in length should be made along the raphe, which should be used as a guide in the place of the staff. The urethra is to be felt for whilst the patient strains, and opened; when a female catheter should be introduced into the bladder, and the urine withdrawn. The next step has for its object the division of the stricture; which, if the obstacle be (as is usually the case) behind the scrotum, should be done in the following manner:—A male catheter should be passed to the stricture, its point felt for in the perinæal incision, and the adventitious growth between it and the finger divided by the knife. The male catheter should then be passed into the bladder, and *retained* there; the catheter should be maintained in the bladder (although this measure has been condemned), for if it be withdrawn, urine may be extravasated in the urethra, and the stricture will certainly become more formidable than ever. If the stricture be in front of the scrotum, it is not safe to divide it with a knife; but its relief must be attempted by bougies, which are always more likely to effect the desired object after the urethra has been opened posteriorly.

The recommendation of caustic has been confined by Mr. Cooper to a few cases only, but we wish the employment of this agent had been still further restricted by him, more especially as he advises the use of potassa fusa, which is so very soluble, and which in the hands of young or incautious surgeons may prove most mischievous in lighting up violent inflammation.

DR. ASHWELL ON UNDUE LACTATION.

HAVING stated that undue lactation has not received a fair share of attention, and made honourable mention of Dr. Marshall Hall, "the only author who has bestowed upon it more than a few incidental remarks," Dr. Ashwell proceeds to assert, that exhaustion, generally attended with

symptoms of *reaction*, but sometimes of depression, is the essential feature of lactation when it has become a disease; that anæmia, irritability, and pallor, varying in degree, are as manifest as in chlorosis, and that the following propositions may be readily proved:—First, that lactation, to be morbid, need not be long; evil consequences may ensue soon after its commencement, occasionally within a few weeks,—more frequently within a period protracted beyond nine months. Secondly, that organic lesions may, although very rarely, ensue from undue suckling; and thirdly, that weaning the child is indispensable to the cure.

Persons usually attacked.

The period of suckling is, happily, in the majority of women, one of health and enjoyment; nevertheless there are exceptions to this rule, a circumstance to be expected in an artificial and increasingly luxurious state of society. Thus, robust and plethoric women rarely suffer from over-nursing, unless, indeed, they be weakened by insufficient food, anxiety, broken rest, or the protraction of lactation to fifteen months, or more. On the other hand, women originally of susceptible, weakly, and especially of strumous constitutions, whose minds have early and long been cultivated at the expense of their physical strength, who live in confined and unhealthy places, who before marriage have suffered from chlorosis, and who have since been weakened by hæmorrhage, and leucorrhœal discharges, or, indeed, by any undue secretion,—are most frequently the sufferers from undue lactation.

Symptoms of Undue Lactation.

Occasionally in a few weeks,—commonly in a few months, it will be apparent, from the imperfect nourishment of the infant, and the anæmia, debility, and pallor of the mother, that the injurious consequences of nursing have commenced. Among the earlier symptoms are, a heavy, dragging sensation in the back and loins; and directly between the scapulæ, when the child is at the breast; a feeling of peculiar emptiness and sinking at the pit of the stomach, and over the whole abdomen, and a deficient supply of milk. At this period much might be done by weaning entirely, or partially, and securing the mother's rest at night; measures which would remove the preceding symptoms, and restore the appetite and strength. But if nursing be persevered in, the morbid results are soon aggravated,—there is excitement or depression of mind, a proneness to hysteria, a frequent and compressible pulse, muscular debility, almost complete loss or fastidiousness of appetite, constipation, or flatulence, tormina, and looseness of the bowels, headache, vertigo, and impaired vision, pain between the shoulders, or below the cartilages of the false ribs, and, ultimately, frequent palpitation, cedema of the face, and swelling of the ancles.

Complications of Morbid Lactation.

1. Profuse menorrhagia. 2. Leucorrhœa. 3. Functional amaurosis;—this disease is often removed by weaning alone. Quickly-recurring pregnancy should be avoided; for the continuance of nursing, or the early occurrence of pregnancy, maintain a congested state of the eye. 4. Jactitation; this affection should not be considered a distinct disease, but rather a symptom of weakness and general irritability, and on this account rendering it imperative that the child should be weaned. 5. Epilepsy. 6. Insanity;—this disease rarely takes the acute form at any stage, and hardly ever at the beginning. The disease is usually marked by perversity of the disposition, and eccentricity of conduct; but, by an attempt to carry on lactation by the aid of stimulants (as porter, wine, and animal food), it is likely to be converted into the acute or wild form of mental aberration. This species of insanity is readily cured; but weaning (with mild diet, and tonics) is absolutely necessary. If the patient attempts nursing after another pregnancy, there is a great probability of a recurrence of insanity, which, in this respect, as well as many other points, resembles puerperal mania.

Undue Lactation may, although rarely, induce Organic Disease.

The brain may suffer in this way, and headache must be considered an alarming symptom, unless it be transient, general, and slight. Danger is to be apprehended if it be permanent in one region, intense, or preceded by rigour; and still more, if there be partial paralysis, mental peculiarity,—forgetfulness approaching to imbecility,—or impeded deglutition, impaired hearing, or deafness.

The lungs may be affected, and phthisis be induced.

The uterus may undergo organic change.

Dr. Ashwell thinks the organic diseases in many instances, consist in softening,—the result of a morbid state of the blood, and produced without the intervention of inflammation,—and quotes from Andral in support of his opinion.

Treatment.

In functional cases, total or partial weaning, tonics, country air, exercise, the securing to the mother unbroken rest, and nutritious unstimulating diet.

In organic disease, the same measures, in addition to counter-irritation, and, *perhaps*, cupping or leeching. If another pregnancy quickly supervene, the woman should not, on any account, attempt to suckle.

To illustrate his opinion, Dr. Ashwell records some cases.

1. An instance of insanity from undue lactation, which recovered, but again returned during the suckling consequent upon another pregnancy.

2. An instance of insanity from the same cause, but not marked by any peculiar circumstances; recovery in three months.

3. A case of phthisis occurring in a woman who had nursed her last two children for seventeen months each; during the last period of nursing, she has had (in vain) a full allowance of porter, &c. for the purpose of enabling her to continue the supply of milk.

4. A case of softening, and abscess of the brain. A lady had a tedious labour, and suffered from hæmorrhage, but recovered, and nursed her child for twelve months, when she began to complain of great weakness, and headache. The proper treatment was pursued, except weaning, which she declined. The symptoms became more serious,—unconsciousness, or epilepsy, or jactitation,—at first occasionally, then more frequently, and in an aggravated form. Death in three months. Dissection revealed the above-named changes.

THE LIMB SAVED, AND CONSIDERABLE POWER OF MOVING THE KNEE-JOINT PRESERVED, AFTER THE PATELLA HAD BEEN CARRIED AWAY BY A GUNSHOT WOUND, AND THE ARTICULAR CAVITY WIDELY EXPOSED.

(Communicated by Mr. Ward, of Huntingdon, through Mr. Bransby Cooper.)

Mr. E. M—, on his return from shooting, had received a gunshot wound of the right knee, in attempting to force his dog from its kennel, with the butt-end of a loaded gun; the contents struck the patella on the outside of the knee, carrying away the whole of that bone, except a small solid triangular portion, which still adhered to the ligament; there was a nearly circular wound of the integuments, completely exposing the joint, and sufficiently large to admit the whole hand into the joint between the tibia and femur, but the cartilages of these bones were uninjured.

The propriety of immediate amputation was the first subject which suggested itself; upon mature deliberation, however, it was resolved to endeavour to save the limb: first, because the exposed bones and their cartilages were uninjured; secondly, because the soft parts around the wound were not contused or lacerated, and, consequently, there was little danger of sloughing; thirdly, because the patient was young, healthy, and temperate.

Treatment.

The patient was placed on his back, with the knee slightly flexed; a poultice was applied, and a full dose of opium given. In the morning his spirits were good, his pulse

quiet, and he complained of nothing more than a slight aching in the knee.

An anodyne was given at bed-time, for a few nights, aperients occasionally exhibited, and the poultices continued until granulations began to rise, when the piece of the patella was removed, and the surface dressed with lint dipped in oil, whilst slips of adhesive plaster were employed to approximate the edges of the wound.

On the 21st January, 1839, the wound was healed, and the gentleman was so recovered as to be able to dress himself, and sit up in a chair. In March, he rode on horseback six miles to the house of Mr. Ward.

He has remained well to the present time, and has long thrown aside the splint and bandage, which he wore for some time after rising from bed. Mr. E. M.— can walk without a stick, dance, or run without much inconvenience.

Mr. Cooper observes, in reference to this case, that there is no kind of injuries the treatment of which is so likely to be improved as those of the joints. It is no longer deemed indispensable to sacrifice the limb; and it would seem that the same danger is not to be apprehended from wide as from small and punctured wounds.

MISCELLANEOUS FOREIGN INTELLIGENCE.

NEW MODE OF TREATING TUMEFACTION OF THE SPLEEN, WITH OR WITHOUT INTERMITTENT FEVER, OR DROPSY.

Dr. VOISIN, of Limoges, affirms that he has cured eight or ten cases of enlarged spleen, as also the ascites and icteric tint of the skin consequent to that affection, by means of a mercurial plaster (*vigo cum mercurio*), with six or eight grains of sulphate of quinine applied to the region of the spleen. The plaster is to be renewed every forty or fifty days. In three or four cases treated by this eminent practitioner, the diseased organ occupied about two thirds of the left half of the abdomen. The doctor adds that the deuto-sulphate of iron, taken in *the same dose as sulphate of quinine*, when the latter is administered internally, is equally efficacious with it in the cure of intermittent fever. The fault, in the administration of the sulphate of iron, is the too small dose in which it is generally prescribed.

THE NECK OF THE UTERUS, AND NOT THE VAGINA, THE SEAT OF THE BLENNORRHOIC DISCHARGE IN FEMALES.

THE *meatus urinarius* is unquestionably one of the seats, but Dr. Gibert, recently the physician of the Female Venereal Hospital, says that in every case where he has used the speculum, he has perceived a discharge from the uterus coexistent with that of the *meatus urinarius*; and as the uterine discharge continues after the urethral, the doctor considers the neck of the uterus as the principal source of the blennorrhagic flux. In the greater number of cases the vagina has no part in the secretion of the discharge. On the other hand, in every case the urethral and uterine fluxes coexist. In 216 patients observed in the wards of l'Ourcine, only 40 had vaginal affection of any kind, and from this number must be deducted several instances where redness was the only symptom, and which was promptly subdued. Hence the term *vaginite* for blennorrhagia is evidently unfounded. The affection of the vagina may possibly be the result of accidental and secondary irritation.

INJECTIONS INTO THE CAVITY OF THE UTERUS.

The French medical press contains a polemic on this subject, which, in the year 1832, formed the matter of an essay, published in the records of the Academy of Medicine. The opponents of the practice adduce instances of extreme pain in the loins having been produced by it, which have been accompanied by fever, so as to lead to a presumption of inflammation of the Fallopian tubes in some cases, and of the ovaria in others. M. Leroy d'Etiolles gives a case in which acute pain was instantly produced in one of the flanks, with rapid enlargement of the ovary on the affected side, and a degree of tympany which that author never met with in any other disease. No pain was produced by pressure, which would serve to distinguish

this affection from peritonitis. Dr. Vidal, of the Female Venereal Hospital de l'Ourcine, considers interior uterine injections to be not only a useful remedy, but free from the inconvenience above stated, if gradually introduced by means of a canula, which, by not filling up the orifice, would suffer the liquid to escape after having washed the sides of the uterus. The opposing party contend that this gradual injection does not wash away the tenacious mucus, which it is often important to detach.

A COURT APPOINTMENT.

M. GUERSANT, physician to the *Children's Hospital*, Paris, has been lately appointed consulting physician to the king of the French. Is his majesty, Louis Philippe, already making preparation for his second childhood?

JONES'S COMPARATIVE ANATOMY.

Recently Published, Price 2s. 6d., Part XII., containing 28 Illustrations, of

A GENERAL OUTLINE OF THE ANIMAL KINGDOM, and MANUAL OF COMPARATIVE ANATOMY. By THOMAS RYMER JONES, Professor of Comparative Anatomy in King's College, London.

This work will comprise, in one volume, the structure and economy of every class of living beings, and their adaptation to the circumstances in which they are severally destined to exist. The principles of zoological arrangement will be developed, and, in order to render the work as intelligible as possible to unscientific readers, a glossary of technical terms will be supplied in the concluding part.

"This work was much wanted, particularly by the younger members of the profession, whose attention we strongly direct to it."—*Medico Chirurgical Review*.

John Van Voorst, 1, Paternoster-row.

BOOKS RECEIVED.

Observations on the Surgical Practice of Paris. Illustrated by Cases. Being a Thesis, to which a gold medal was assigned by the Senatus Academicus of the Edinburgh University, at the Graduation of 1840. By W. O. MARKHAM, M.D. London: Highley. 1840. 8vo. Pp. 114.

[We shall make frequent extracts from this interesting Thesis.]

Elements of Natural Philosophy, &c. By GOLDING BIRD, M.D. London: Churchill. 1839. 8vo. Pp. 400.

A Treatise on the Diseases of the Eye and its Appendages. By RICHARD MIDDLEMORE, Surgeon to the Birmingham Eye Infirmary, &c. Longman and Co. London, 1835.

[We take this opportunity of stating, that one of the most distinguished ophthalmological professors in the provinces has undertaken to review for us all works on diseases of the eye, &c.]

Practical Remarks on the Discrimination and Appearances of Surgical Disease, &c. By JOHN HOWSHIP, Surgeon to the Charing-cross Hospital. Churchill, London, 1840. 8vo. Pp. 400.

A Letter to Sir B. C. Brodie, containing a Critical Inquiry into his "Lectures Illustrative of certain Nervous Affections." By WILLIAM GOODLAD, M.R.C.S. &c. Longman and Co. London, 1840. 8vo. Pp. 154.

Practical Observations on the Causes and Treatment of Curvatures of the Spine, &c. By SAMUEL HARE. Simpkin and Marshall, London, 1838. 8vo. Pp. 152.

On the Nature and Treatment of Stomach and Urinary Diseases, &c. By WILLIAM PROUT, M.D. Third Edition. Churchill, London, 1840. 8vo. Pp. 483.

A Practical Treatise on the Cure of Strabismus, or Squint, by Operation, and by Milder Treatment. With Plates. By P. BENNETT LUCAS. Highley, London, 1840. 8vo. Pp. 90.

Printed by RICHARD CLAY, of No. 5, Newmarket Terrace, Cambridge Heath, in the County of Middlesex, at his Printing-office, Nos. 7 and 8, Bread-street Hill, in the Parish of St. Nicholas Olave, in the City of London; and published by WILLIAM BALL, of 34, Paternoster Row, in the City of London, at No. 34, Paternoster Row.—Saturday, Oct. 17, 1840.