lids are cedematous, and completely cover the eye. I removed the sloughy mass of brain this morning; it was very dark, of a greenish-black colour, exceedingly fetid, and when shaken in water the principal portion hung together, as if it were sloughy membrane. Mr. Cock said this was attributable to the vessels of the mass. On removing it there was some little bleeding, which stopped on slight pressure. There is a sloughy spot over the right trochanter. He will not allow a poultice to remain on; bowels open twice to-day.

16. Has passed a better night, but cries out vehemently between his sleeps; is conscious, but unable to articulate; has more power in his right leg; he can turn himself in bed; right side of face much swollen, skin inflamed; bowels open in the night. Warm water dressing applied to face.

1 P.M. Has been very restless all the morning. Face erysipelatous; about two inches above the affected eye on the same side as the injury there was a collection of pus, and another commencing about an inch below the seat of fracture. I opened the anterior abscess, and expressed some ill-formed sanious pus; there was a great tendency to bleed, which was counteracted by pressure; the posterior abscess contained healthier pus; a mass of sloughy brain was protruding from the wound, as large as the former one, and very fetid. A poultice was applied over the whole, and water dressing to the face.

17. A collection of sanious fluid was liberated from the anterior abscess; the posterior abscess was also similarly distended; he passed a very restless night. On trying, without much force, to remove the sloughy portion of brain, there was much bleeding, stopped by lint and pressure. Is taking a good deal of nourishment.

18. Is now constantly moaning; has been screaming incessantly and violently during the night; is very passionate when he cannot make himself understood; bowels open freely; erysipelas diminished; motion and sensation tolerably perfect; tongue natural; pulse very weak.

19. Very restless, still screaming and moaning; perfect use of both legs; takes scarcely any nourishment; countenance contracted; pulse exceedingly weak; bowels open twice naturally.

20. 1 P.M. Opened another abscess beneath the scalp; pus tolerably healthy; there is vomiting. He is evidently sinking, constantly moaning, and, though not able to articulate, seems to retain consciousness. He died at half-past 6 P.M.

Post-mortem Appearances.

Integuments of the injured part sloughy. Trephine opening and that of another small angle removed, filled up with sloughy brain. Over the left side of the vertex, a little posteriorly, two unequal arches of bone, two inches long and half an inch wide, were much depressed. The deeper arch consisted only of the inner table. There were two ragged perforations of the dura mater half an inch wide; a thick pus, or feeble fibrin, was spread about these, and there were some detached spots, one as large as a half-crown, immediately within the spine of the occiput, broad layers beneath the dura mater on the left side, and one over the right anterior lobe, composed of pus superficially, and a granular adherent layer beneath; under these the brain was green. A mass of slough, as large as a pigeon's egg, with pus, was lodged in the brain at the point of fracture. Part of the walls of the cavity so formed were defined, firm, grainy and vascular, a line and a half thick. In other parts there was softening, varying in degree, and extending into the ventricle, which contained an unnatural quantity of clear fluid. The fornix and septum lucidum were extensively softened, and probably even perforated. The pons Varolii was covered with purulent serum in great quantity."

Now from the continuity of the posterior crus of the fornix with the hippocampus major, the softening of the former may probably tend to explain the loss of articula-

tion.—I find that it is now late, and as there are several points on this important subject worthy of further consideration, I shall postpone any discussion of the interesting case now in the house till our next meeting.

WORCESTER INFIRMARY.

PRACTICE OF DR. HASTINGS.

DIABETES MELLITUS.

WM. GOODMAN, æt. 40, was admitted on Jan. 15, 1820, under the care of Dr. Hastings. The following symptoms were observed on his admission to the infirmary:—

Pain in the epigastric region; urine very copious; urgent thirst; is affected with pain in the region of the left kidney, shooting up at times to his heart; also in the right arm and shoulder; constant pain in his head, with sickness at times; urine high coloured, copious and clear; tongue white; much thirst; pulse 108; appetite good; bowels not open. Has been ill two years with pain across his loins. Has been much worse within the last six months.

18. To have some cathartic mixture until the bowels are opened.

20. Three dejections yesterday, two this morning; pain relieved. Eight ounces of blood to be drawn by cupping-glasses from the painful part. Solution of sulphate of magnesia, one ounce and a half thrice a day; calomel, one grain, every second night.

22. Makes less water; pain much relieved; bowels regular; not so much thirst. Continue the remedies.

24. Made eleven and a half pints of urine in the last twenty-four hours; drank four quarts. Had leeches on the painful part yesterday, which have very much relieved it. Tongue still white; pulse 102; three dejections yesterday. Continue the remedies.

25. Has been making less water every day since he came in; eleven pints in the last twenty-four hours; tongue cleaner; bowels open. Continue.

28. Made on the 26th ten and a half pints, and yesterday eleven pints; thirst not so urgent; tongue still furred; bowels quite open; pulse 106. Has omitted the calomel in consequence of complaining of griping pain in the bowels. Urine does not taste sweet; does not deposit any saccharine matter on evaporation. Continue the sulphate of magnesia. Omit the calomel.

31. Made on the 29th only seven pints, and yesterday nine pints; thirst continues; three dejections yesterday, one to-day; has still some pain in the left hypochondrium; tongue cleaner; complains of pain in the head. Repeat the leeches; continue the solution; medium diet.

Feb. 3. Pain nearly gone; thirst abated; three stools yesterday, and one this morning; made eight pints and a half on the 1st, and yesterday four. Tongue cleaner; pulse 90; still complains of pain in the shoulder; pain in the head relieved; skin very dry.

4. A tepid bath every other night.

6. Urine, five and a half pounds; [drink, five pounds; skin not so dry; pain of the shoulder relieved by the bath; appetite good; bowels regular; not so flatulent.

8. Urine, six and a half pounds; drink, five pounds; flatulence better; thirst abated. Continue remedies. Extract of gentian, ten grains, thrice a day.

10. Urine, five pounds; drink, four pounds; bowels regular; sweats much at night.

15. Urine, five pounds; drink, four pounds.

17. Urine, six pounds; drink, four pounds; tongue cleaner; complains of weight in epigastrium. A blister to the epigastrium.

(Made an Out-patient.)

Mar. 1. Solution of sulphate of magnesia, one ounce, thrice a day; extract of gentian, ten grains, thrice a day; compound ipecacuanha powder, five grains, whenever the pain is severe.

15. Continue remedies; a blister to the epigastrium.

April 26. Complains of pain in the region of the kidney, and passes a large quantity of water. Blood to be drawn from the arm to ten ounces.

May 10. He passes thirteen pints of urine in twenty-four hours. Ipecacuanha, six grains, every morning.

(Made an In-patient.)

18. His thirst, and the proportion of urine to fluid taken into the system, have gradually increased since he has been out of the house. Compound rhubarb pill, ten grains, thrice a day.

19. Repeat the pill. One pound of meat a day.

23. To have the pill only twice a day.

25. Complains of being much griped. Carbonate of soda, fifteen grains; tincture of orange peel, one drachm; infusion of cascarrilla, eight drachms; to be taken thrice a day. Cupping-glasses over the loins.

29. Complains of pain in the back.

June 5. Complains of pain in the head and back. To be bled to eight ounces.

7. Pain in the head and back relieved; blood somewhat sisy.

11. Complains very much of pain in the back and head; he is stronger. To be bled to eight ounces; continue remedies.—15. Pain of the back continues. Repeat the bleeding; continue the remedies.—17. Headache better.—

19. Feels stronger; pain in the back continues, but is not so violent. Repeat the bleeding; continue medicines.—

21. Gains strength; pain of the back is better; he now perspires freely; sometime ago his skin was constantly dry.—23. Pain of the back returned. Repeat the bleeding; continue remedies.—28. Feels better; sweats freely.

July 5. Complains very much of pain in the back. Six ounces of blood to be drawn by cupping-glasses over the loins; continue remedies.—10. Pain relieved.—12. Headache. To be bled to eight ounces.—14. No pain in the back.

Aug. 10. He does not make more urine than before he was taken ill; it is quite natural in its appearance. He has gained a good deal of flesh, and is capable of a good deal of exertion; digests his food well; skin is not dry. Six ounces of blood to be drawn by cupping-glasses over the painful part.

Dismissed cured.

Dec. 22. Has returned, with pain in the back, and other aggravated symptoms; constant pain in the head; frequent sickness and nausea. In the course of about twenty-four hours, drinks ten pints of water, and the urine discharged in that time is about fifteen pints; water clear and very light-coloured; pulse 90, rather hard; tongue white. Twelve leeches over the region of the kidneys.

27. Calomel, two grains, thrice a day; to be bled to ten ounces. Flesh diet.

1821, Jan. 12. Severe pain in the region of the left kidney; pulse hard. To be bled to ten ounces.

13. Pain not abated; pulse hard; blood much buffed and cupped. Repeat the bleeding.

17. Pain in the loins abated, and he is in every respect better.

23. Repeat the bleeding.

24. Has been seized with shiverings, and cold and hot fits; pulse 100; pain in the limbs; much thirst. An emetic at once; antimonial saline mixture every four hours; calomel, one grain, every night.

Feb. 1. Pain in the epigastrium; has returned to meat diet, omitted on the 24th. Eight ounces of blood to be drawn by cupping-glasses from the epigastric region.

14. Infusion of calumba, one ounce; tincture of orange-peel, one drachm; to be taken twice a day.

17. Pain in the loins and side. Eight ounces of blood to be drawn by cupping-glasses from the loins.

21. To be bled to eight ounces.

26. Pain continues; is very low.

Mar. 24. The quantity of water since last report has been about four pints and a half, or five pints; pain in the

back continues, and tongue loaded. Calomel, one grain, every night.

1823, April 16. Has never been well since he went out of the house twelve months ago; but has been able to follow his usual employments; pain in the situation of the left kidney is very violent; health not much impaired; tongue furred; bowels regular; the quantity of urine voided daily is about twelve pints; he drinks about ten pints. Repeat the calumba mixture; calomel; flesh diet.

20. To be bled to six ounces.

1827, Feb. 3. Has never been well since he left the house three years ago. Headache; occasional pain in the stomach. The quantity of urine voided daily about thirteen pints, and drinks about ten. Appetite perfectly good; bowels regular; tongue furred; pulse 90; is much emaciated. Blood-letting to ten ounces; calomel and opium, of each one grain, every night.

1828, Dec. 6. Pain and tenderness in the epigastrium; has been much purged for several months; tongue furred; pulse 96; headache. Meat diet. Calomel and opium, one grain, at night; solution of sulphate of magnesia, one ounce, thrice a day; eight leeches to the epigastrium.

23. Passes nine pints of urine in twenty-four hours.

26. Appetite good; pain in the right and left hypochondrum much the same.

1829, Jan. 5. Drinks seven pints in the twenty-four hours; makes eight pints and a half of urine.

8. Ingesta, seven pints; egesta, eight pints. Compound spirit of ammonia, twenty drops; infusion of calumba, one ounce; thrice a day. Continue the pills.

April 8. Six leeches were applied to the side.

29. Passes about six pints of urine in the twenty-four hours; much pain in the side and lumbar region; urine sweet.

May 2. Complains of pain all over the body, but particularly in the region of the kidneys; feels very low; pain in the head; very thirsty; makes a great quantity of water. Compound spirit of ammonia, fifteen drops; tincture of gentian, one half drachm; infusion of calumba, one half ounce; thrice a day. To be bled to twenty ounces.

14. Injesta 9; egesta 11½; bowels confined.

15. Drank nine pints; made ten and a half pints; pain very severe. Had caustics put in his back to-day; leeches in the lumbar region.

27. No saccharine deposit; painful micturition; uses warm baths at night.

June 2. Taste of residuum on evaporation decidedly salt; the residuum very small in quantity.

9. Drinks nine pints; passes ten pints.

20. Saccharine deposit, two scruples in one quart; of a saline taste.

July 1. Gains strength; drinks eight pints; passes eight and a half pints.

9. Residuum one drachm from one quart of urine; lost eight ounces of blood.

Aug. 3. Passes about the same quantity he drinks; residuum three drachms one scruple from a quart of urine. Had a seton put in his loins, which was removed on account of hæmorrhage supervening.

Dec. 23. Great abdominal pain and spasmodic action; bowels have not been moved for nine days; urine scanty. Croton oil, two drops; crumb of bread, enough to make two pills; to be taken at once.

The patient died on the 24th.

On examination, the abdominal viscera were all healthy. The kidneys and liver presented no morbid appearances sufficient to account for the symptoms.

On opening the thorax, the pleura costalis presented firm adhesions, particularly on the left side. The substance of the lungs was completely gone on that side, being studded with tubercles.

The right lung was also tuberculated. There was discovered no cavity.

The heart was rather less than natural.

The following table indicates the relative proportions of *Ingesta* and *Egesta*, at various periods during this protracted case:—

Days.	Inges.	Egest.	Days.	Inges.	Egest.
	Pts.	Pts.		Pts.	Pts.
1820. Jan. 24	8	11½	1821, Jan. 5	10	12
25	8	11	14	8	9½
26	8	10½	29	6	7
Feb. 3	5	8½	Feb. 17—21	5	5
17	4½	6	March 4	5½	5
May 10	6	8	1823, April 17	10	12
23	9½	12	May 15	7	7
24	9	11	Sep. 10	7½	8
30	8	9½	1824, Aug. 18	6	8
June 1	8	10	1827, Feb. 6	10	12
7	8	9½	20	7	9
11	9	9½	23	6½	7½
12—21	7½	9	1828, April 29	7	9
23	7	8	1829, May 14	9	11½
28	8	8	18	9	10½
July 5	7½	8	June 9	9	10
12	7½	8	July 1	8	8½
21—Aug. 7	6	6	August 3	8	8

The quantity of saccharine matter in a pint of urine was also determined at different periods of the disease.

On the 10th of May, 1820, it amounted to fifty grains. From June, of the same year, to the middle of May, 1829, the residuum ceased to be saccharine. In June, 1829, and during the remainder of that year, from one half to three drachms of residuum, having a saline taste, were obtained from a quart of urine.

CONTRIBUTIONS TO THE PATHOLOGY OF CHILDREN.

BY P. HENNIS GREEN, M.B.

Lecturer on Diseases of Children at the Hunterian School of Medicine.

SCROFULOUS TUBERCLE OF THE BRAIN.

AMONGST the various diseases of the brain to which children are subject, there is none so insidious in its approaches, so difficult of detection from the diversity of symptoms by which it is attended, or so completely neglected by medical writers as scrofulous tubercle. Were we to judge from the works of English or continental authors on diseases of children, it might be inferred that scrofulous tubercle of the brain is an extremely rare affection, for although I possess every standard work on the subject, I am not acquainted with one in which cerebral tubercle is mentioned: it may not, therefore, be superfluous to direct the attention of medical practitioners to this disease, of the frequency of which long experience has fully convinced me. With this object, I shall first relate several cases, and then endeavour to point out the symptoms by which scrofulous tubercle of the brain may be distinguished from the other affections of that organ.

CASE I.

Vomiting and headache; acute sensibility of the skin; occurrence of small-pox; death; extensive tubercle of the cerebellum.

ALEXANDRE LEGROS, 3½ years of age, was admitted into the Children's Hospital on the 1st of July, 1835, whither he was brought by the nurse under whose care he has been since the age of three months. The mother of the child is consumptive. For the last month he complained of pain in the abdomen, for which a medical man ordered baths

and twelve leeches to the abdomen; he took some medicine (supposed to be vermifuge) and vomited it up; on the following morning he had some convulsive attacks (the first lasted ten minutes), during which the face was very pale, the eyes fixed, and the arms contorted; he vomited again, without having taken medicine, in a few days, and the vomiting has recurred since then several times. The child was frequently stupid; no delirium or pain in the head; no squinting observed; no diarrhoea. For the last eight days the boy has had pain in the back of the head, and since then squinted; he also evinced an unwillingness to move about; makes urine frequently. He remained in the hospital until the 10th with no symptom worthy of notice, except squinting and a slight movement of the lower jaw, and was removed by his friends.

On the 16th of July he was again brought to the hospital, with a slight eruption of small-pox, in the first stage. We now learn that on the 14th and 15th the child vomited (he also vomited this morning), and complained very much whenever the limbs were touched; he had a strong convulsion this morning, and several similar attacks since Wednesday last.

Present state.—The boy now lies quietly on his back; the face full, pale and cool, but the forehead excessively hot; general surface moderately warm; there is some convulsive strabismus of the right eye, the eyelid of which is more closed than the left one; tongue clean and moist; the lateral movements of the lower jaw are more marked than they were before, and he occasionally catches the bed-clothes between his fingers; answers clearly when spoken to; extremities neither stiff nor contracted; when the integuments of the hands or arms are touched he evinces the most acute sensibility; no pain in the abdomen; pulse small, 128; respiration free, at one time 28, at another 36; short, dry cough.

17. The general appearance is the same as yesterday; had no regular convulsive fit, but much agitation and delirium last night; the squinting and movements of the jaw continue, with jactitation of the hands; right eyelid nearly closed; pulse small, 130; respiration 34; skin very warm; abdomen contracted and free from pain; tongue red and covered with a number of white pips, like aborted pustules; the face and extremities are covered with small papulæ, most of which are flat and irregular, like those of measles; a few only are prominent.

Evening. Pulse 144; no motion of lips or jaw; thirst; deglutition perfect: some areola round the pustules on the body; on the thighs they are very confluent; the child has been seized with a peculiar trembling of the upper extremities.

19. Child appears to be dying. He lies on his back, with both hands elevated to the ears, and trembling violently; face pale, covered with a number of flat and pale papulæ; no convulsive motion of muscles of eyes or face; pupils natural; skin of forehead still excessively warm; pulse small and rapid, 140; respiration 46; lips and tongue dry and dirty; low muttering delirium; the pustules on body are assuming a bluish aspect. Died in the evening at eight o'clock.

Body examined 15 hours after Death.

Head: The superior surface of the brain is pale, and the membranes not injected; on the left hemisphere, along the course of a large vein, there is a sero-sanguineous effusion, about two inches in length; in the right hemisphere, about two and a half inches from the frontal bone, and a quarter of an inch from the medium line, there is a large solid tubercle embedded between the convolutions, with two or three small tubercles in the adjoining pia mater. The membranes and surrounding nervous substance are not altered; on the side of the left hemisphere, near the falx, there is a flat mass of tubercular matter, a quarter of an inch in diameter, partially embedded in the substance of the brain; the pia mater lying over it is not injected, but it is surrounded by a deep red circle of injection, about two