PROVINCIAL MEDICAL & SURGICAL JOURNAL.

No. 9. Vol. I.] LONDON, SATURDAY, NOV. 27, 1841-2.

PRICE SIXPENCE.
[STAMPED EDITION SEVENPENCE.]

CASE OF WOUND OF THE AXILLARY ARTERY.

TO THE EDITORS OF THE PROVINCIAL MEDICAL AND SURGICAL JOURNAL.

GENTLEMEN,—Wounds of the axillary artery being of rare occurrence, perhaps the following case will not be unacceptable to your readers.

Your obedient servant,
EDWARD SCRATCHLEY, M. D. P.

At about one o'clock in the morning of the 12th of May, 1840, I was requested to visit a person who had been wounded, as I was informed at the moment, in the arm, and was lying at a wine-shop in a village about two miles from Paris. Hastening to the spot, I found a young man, named Viscontet, lying on a mattress on the floor, pale and almost lifeless, his clothes and the mattress deluged with blood. One of his comrades was beside him holding a towel stuffed into the left axilla, from which, however, blood was escaping. On inquiry I found that between eleven and twelve a quarrel had taken place between Viscontet and another man, which ended in blows. Viscontet, whilst raising his left arm to strike his opponent, had been stabbed in the arm-pit with a sharp-pointed knife. Violent hæmorrhage immediately ensued, and he fell fainting. One of his companions had the presence of mind to tear up his shirt-sleeve, and to oppose the escape of blood by stuffing a towel into the arm-pit. Persons were immediately sent for a surgeon, but from various circumstances none could be found, and Viscontet remained without aid until half-past one, when I arrived.

Finding that I had a much more serious accident to deal with than I expected, and that immediate assistance must be given to the patient, my first step was to examine the wound; for this purpose I took an ordinary-sized door key, and, wrapping my handkerchief round its handle, compressed the subclavian artery above the clavicle, then raising with my other hand the arm, I perceived a wound parallel to the direction of the pectoralis major, from an inch to an inch and a half in length, immediately over the axillary artery; in fact, the wound was perfectly similar to an incision made by a surgeon for the ligature of this artery; on removing for a moment the compression on the subclavian, arterial blood gushed out of the wound: as I said before, the patient was pale, exsanguinous, with pulse hardly perceptible, with skin icy-cold. What was to be done next? Ligature of the axillary artery, even on the dead subject, is an operation of some difficulty; but placed as I was, without assistance, any attempt at the operation would have been wholly unjustifiable, when the

No. 61.

loss of a few ounces more of blood might have left the patient dead in my hands. The essential thing was to stop the hæmorrhage; for this I took a couple of towels, rolled them up into a thick pad, and placing them in the axilla, fixed them in this position by another towel brought from under the arm-pit, and tied over the shoulder of the same side; on twisting the towel tightly by means of a stick, I found that I had perfect command over the hæmorrhage. The arm was then bound to the side, and the substitute for a tourniquet fixed with tape to prevent its slipping. The patient was placed on a stretcher, and carried to the Hospital Beaujon, where he arrived about three. During the transport, so efficacious was the apparatus, no blood was lost. He was put into a bed, and, as no blood escaped, the parts were left in situ, and it was not considered necessary by the house-surgeon to send for Mr. Laugier, surgeon of the hospital, before five A.M., when a slight hæmorrhage recommenced, but was immediately arrested by tightening the apparatus. At half-past five, Mr. Laugier arrived, and the following description of the operation is taken from the "Bulletin Chirurgical" of Mr. Laugier.

"On arriving at the hospital," says Mr. Laugier, "I found Viscontet extremely pale, with pulse entirely suspended on the left side, and very feeble on the right, but intellectual faculties entire. Compression above the clavicle having been made by the fingers of an assistant, the towels were removed. Behind the lower border of the pectoralis major, and parallel to the direction of its fibres, is a wound fifteen lines in length, and penetrating deeply into the axilla. I began by enlarging the wound towards the chest; this first incision exposed a coagulum of dark-coloured blood, placed before the axillary artery and the plexus of nerves and veins. The incision was next carried downwards in the direction of the brachial artery. In this stage of the operation, one of the venæ comitantes was slightly wounded, and a small flow of venous blood was the consequence. As every ounce of blood was of importance to the patient, I thought it best to pass a thread round the opening in the vein.*

* "This circumstance," adds Mr. Laugier, "shows the necessity of following the wise precept, founded on experience, of dividing layer by layer, and on a director, the parts immediately surrounding large vessels. In this case the colour of the vein was confounded with that of the coagulum covering the vessels and nerves; and although but a small puncture was made into it, still, if death had not so rapidly supervened, its ligature might have had some influence on the result of the operation." It would be well if this precept were firmly impressed on the minds of all surgeons, notwithstanding the veto passed upon it by a celebrated operator, who recommends, in cutting down on large arteries, to use the knife with the hand unsupported. It is far better for them to appear slow and ungainly in their operations, than to run the risk, by some false stroke, of seriously injuring their patient. I have seen a celebrated surgeon and operator, who disdained the use of a director, open the femoral vein in ligature of the femoral artery. Another surgeon, in operating for femoral hernia, after dividing the integuments, to show off his dexterity, cut with one stroke into the hernial sac, and was on the point of opening the gut itself, when he fortunately perceived where he was. This cut-and-thrust surgery may do well for display, but it is scarcely scientific.

The wound having thus been sufficiently enlarged, and the coagulated blood removed with the finger, I found that the opening in the artery was situated deep in the axilla. The knife with which the patient had been struck, had passed between the median and ulnar nerves, without touching them, making a longitudinal incision into the artery, parallel to its direction and to that of the surrounding nerves and veins.

"In order to find the position of the upper orifice of the wounded artery, the compression above the clavicle was slackened two or three times, but the blood escaped so rapidly and in such quantities as to wholly impede my view. This showed to me clearly the existence of a large opening in the artery. Also, when, to expose more fully the axillary artery, the arm was raised from the body beyond the right angle, the clavicle was pushed upwards, and the fingers of the assistant had no longer command over the artery, and the hæmorrhage recommenced.

"Having thus perceived that if I persisted in attempting to tie the artery, above and below its wound, through the axilla, I stood the chance of seeing my patient die under my hands, I determined on following another plan, and that was to tie the artery above the pectoralis minor. I knew well that this would only be a temporary ligature, and could be considered simply as an auxiliary operation, in addition to the requisite one of tying the artery above and below its wound. In recent wounds of arteries, Anel's or Hunter's method is essentially faulty, and in this case the hæmorrhage would have been reproduced by the lower orifice; for large branches enter the artery beneath the spot where I intended placing the ligature. I dismissed quickly the idea of cutting through the pectoralis major and minor, in order to arrive more directly at the wounded spot; for although by this method I would have been enabled to place two ligatures, instead of three, on the artery, still this advantage, great in itself, would not have compensated for the great destruction of the axilla, and the risk attendant on it. Therefore, the artery being compressed above the clavicle and below in the axilla, I followed the method of M. Marjolin for tying the subclavian below the clavicle, by making an incision between the two portions of the pectoralis major. In this operation the artery is found above the border of the pectoralis minor, covered by the axillary vein, and placed between it and the brachial plexus of nerves; the latter being behind and somewhat above the artery. I arrived easily upon the plexus of nerves and vessels, and, having isolated with great care the artery from the surrounding parts, I carried a ligature around it by means of a curved blunt needle, with the eye at its point.

"Whilst searching for the artery, the following circumstance struck me as being worthy of notice. When the axillary artery is tied below the clavicle for aneurysm, or even when the operation is performed on the dead subject, the axillary vein is always found placed before the artery, and more or less distended, its size varying in different subjects, and also depending on the respiratory efforts of the patient. The presence and volume of this vein are considered as an obstacle to the isolation of the artery and its ligature. It is even recommended, in operating on the living subject, to oppose the entrance of blood into the vein, and

its over-distension, by the finger of an assistant. But to counterbalance the inconvenience resulting from the size and position of the vein, its colour makes its more evident, and easily distinguished from the artery. In our patient, however, the axillary vein was collapsed, and could not be distinguished, either by its volume or colour, from the fascia covering the vessels. The pulsations of the artery were felt distinctly through the vein, but, from the flattened state of the latter, some difficulty might have been experienced in pressing it downwards and inwards; and if its peculiar appearance had not been remarked, or if the needle had been to the slightest degree pointed, or great care had not been taken in passing it around the artery, the vein might have been mistaken for the fascia, and pierced through and through."

"The artery having been tied, I was enabled to return to the axilla, and look for the lower orifice of the wounded artery, from which a small quantity of blood was still oozing. Having tied this, the hæmorrhage ceased for an instant, but, as I expected, it reappeared, after a few seconds, at the upper angle of the wound. I attempted to place a ligature round the upper orifice, and then it was that I was able to judge of the length of the wound in the artery, which extended longitudinally upwards, and appeared to be from twelve to fifteen lines in length. The upper orifice being tied, all bleeding ceased, and the wound was united by a few slips of sticking plaster.

"The patient supported the double operation with patience and courage. It is true that he was in a state bordering on syncope; after the operation, the pulse was scarcely perceptible on the right side, but in a short time it acquired the same strength as before.

"Wine and water was given several times during the operation, and he was ordered to drink freely of cold broth, and take several spoonfuls of the medicated wine of Collioure, and a potion with syrup of cinchona.

"In the evening he was in the same state; pale, anxious, answering questions with precision, but, when his attention was not fixed, uttering unconnected phrases. No blood had been lost since the operation, and the broth and wine had remained on the stomach; his pulse, however, was weaker, and general anxiety increased; he passed the night in the same way, a nurse having been ordered to give him broth frequently; at seven in the morning he was still conscious, but shortly after he fell into a state of syncope, and died without convulsions.

"On opening the body, the pectoralis major and minor were divided so as to expose freely the axillary vessels and nerves; the first ligature was found to have been properly applied, and to embrace only the artery; inferiorly, and in the direction of the vessels, the cellular tissue was slightly infiltrated with blood; the space between the two ligatures, placed above and below the wound, appeared to be from twelve to fifteen lines in length, and was comprised between the origins of the two circumflex arteries; the upper ligature was immediately below the origin of the anterior circumflex, and the lower one above that of the posterior circumflex. The left upper extremity was of a dark colour, and putrefaction seemed more advanced in it than in the neck or right side.

Loss of blood was found to have been the only cause of death.

"In addition to the interest that is always attached to wounds of large arteries, the above case appears to me to possess a certain value from the necessity I was under of modifying my original plan of operating, caused by the great loss of blood before the operation, and by the height at which the wound in the artery was situated. It is evident, from this observation, that cases may present themselves in practice, when ligature of the subclavian above the clavicle, or, at least, ligature of the axillary artery, above the pectoralis minor becomes indispensable, even in cases of wounded axillary artery, which we are not the less forced to tie both above and below the wound. But as my principal reason for thus operating in the above case, arose from the difficulty experienced in compressing the artery above the clavicle, by means of the fingers of an assistant, when the arm was raised, shall I be blamed for not trying some other method of compression? It strikes me that I would have found it as difficult to have kept in position either a pad or the handle of a key, and that the compression would have been in no way more certain.

"I would, therefore, in cases analogous to this, make it a rule, as a preliminary step, to stop the hæmorrhage by first tying the axillary artery below the clavicle. This would almost entirely arrest the hæmorrhage, and would give the operator sufficient time for applying the necessary ligatures both above and below the wound."

The method recommended by M. Laugier appears to me objectionable in many points, but principally on account of the difficulty that would result from the application of three ligatures, in the return of circulation to the limb.

After ligature of the axillary artery, simply above the pectoralis minor, abundant means of carrying on the circulation exist by the anastomoses of the transversalis colli, the supra-scapular, and the internal mammary with the thoracico-acromialis, the subscapularis, the circumflex and the external mammary; but, in this case, by the application of three ligatures, one above the pectoralis minor, the other below the origin of the anterior circumflex, and the third above that of the posterior circumflex, it is evident that the above circle of anastomoses would no longer be serviceable, and that the circulation would have to be restored principally by the anastomoses between the posterior circumflex, the supra-scapular, and the profound arteries.

That the blood could flow pretty freely by this channel, was shown by the fact of the bleeding having recommenced from the lower orifice of the wounded artery, more abundantly than from the upper, after its ligature below the clavicle. But, as observation has proved that mortification is more likely to ensue in a limb, after the ligature of its principal artery for wound than for aneurysm, we should avoid causing a greater obliteration of a wounded artery than is necessary, for the freer the communication between the collateral branches is, the more likely we are to obtain a favourable result. This, then, seems to me to be a most serious objection to the plan recommended by M. Laugier.

By cutting through more or less of the lower bor-

der of the pectoralis major, M. Laugier might have arrived, as he himself admits, immediately on the wounded spot; but he did not think that "the advantage to be derived from it would compensate for the great destruction of the axilla, the risk attendant on it, and the necessary loss of blood from the division of one of the thoracic branches." And he also adds, "the situation of the wound might require total division of the pectoralis major, a circumstance that should be avoided if possible." In the first place, I can conceive no wound of the axillary artery *through the axilla* would require total division of the pectoralis major. If the wound should have been made from above downwards, through the substance of this muscle, we have nothing to do but to follow Scarpa's advice, and enlarge the wound. In the second place, the division of the pectoralis major, sufficient to have exposed the wounded spot, would not have been attended with greater risk than the operation for tying the artery above the pectoralis minor. Instead of having one large wound, and thus simplifying the operation, two distinct ones were made; time, a matter of life and death in cases of wounded artery, was lost, and it is probable, that before deciding on his plan of operating, and whilst searching, through an insufficient opening, for the upper orifice of the wounded artery, more blood was lost than if he had divided half a dozen thoracics, over which he had immediate command.

It therefore appears to me a safer and more expeditious plan, in cases of wounded axillary artery, to follow the example of Desault, Roux, Delpech, and others, and not to be deterred, by the fear of dividing a few muscular fibres, from enlarging the wound, even if we have to cut through half of the pectoralis major and minor. By this method we arrive more quickly on the wounded spot, and, what is of more importance, we preserve a greater number of collateral branches for the subsequent restoration of circulation, than if we followed the plan of M. Laugier of tying the axillary artery above the pectoralis minor, in addition to its ligature above and below the wound.

A CASE OF CROUP, SUCCESSFULLY TREATED.

By SIR ARNOLD JAMES KNIGHT.

1841. OCT. 10. Sunday, 9 A.M.—I was requested to visit a little boy about four years old. I found him in the advanced stage of a severe and neglected attack of croup. I was informed that about a fortnight before, his grandmother having gone from home, he cried so excessively that those about him could not pacify him, and they thought he had never been well since. On the Friday before I saw him, he had become so much worse, that his friends took him to a medical practitioner, who told them that he had got the croup, and gave him some powders, and a blister; they gave him the powders, but did not apply the blister: the boy grew still worse, and on Saturday night his symptoms were so much increased, that the friends sent for me on Sunday morning. The little patient's countenance was now pale; the lips livid; pulse quick; breathing and cough of that kind which is peculiar to the advanced stage of croup. Think-