PROVINCIAL MEDICAL & SURGICAL JOURNAL.

No. 18.—VOL. I.]

LONDON, SATURDAY, JAN. 29, 1842.

PRICE SIXPENCE.
[Stamped Edition Sevenpence.

CLINICAL OBSERVATIONS ON INCONTINENCE OF URINE,

Delivered at St. George's Hospital.

By SIR B. C. BRODIE, Bart.

By the term "incontinence of urine," I wish you to understand a dribbling away of the urine without any knowledge of it on the part of the patient. It is a different affection from irritability of the bladder, where the patient is obliged to use an effort to empty that organ. In the course of your practice you will find that in nine cases out of ten, where the urine dribbles away, it is from an overloading of the bladder. I have been called to patients whose urine has passed away very well during the day, and at night I have been summoned to them and have found their bladders distended as large as their heads; I have passed the catheter, and have drawn off a large chamber-pot full of urine. Now, such a state of things as this may arise from the presence of an old stricture, or from an enlarged prostate gland, or from paralysis. You will sometimes find in these cases that the bladder is not always distended in an equal degree; the moment the urine attains to a certain quantity, it begins to flow away involuntarily. In some of these cases you will find the symptoms very troublesome to combat; the coats of the bladder will, from long distension and irritability, become inflamed and thickened; and what is worse than this, you will eventually find that the kidneys will become diseased, and then when you draw off the urine you have not cured the disease, but only relieved one symptom; the disease in the kidneys becomes worse from day to day, but that must not deter you from drawing off the patient's water whenever he requires it.

Where incontinence of urine exists, as the result of paralysis, it will be found to occur in cases where the bladder is not overloaded. A gentleman had an attack of rheumatism, for the relief of which he was ordered to rub in a liniment, containing the tincture of cantharides; by some mistake he swallowed some portion of the liniment; appropriate remedies were immediately employed, but he became affected with paralysis, and had retention of urine; for one week he was obliged to have his water drawn off, and then he had incontinence of urine. In four years' time he was able to hobble about upon crutches, and his urine still continued to come away involuntarily; the catheter was frequently passed, but no urine followed, and the bladder seemed to have lost all tone, and not to be able to retain any fluid whatever; so that as fast as it reached that viscus it passed away involuntarily by the urethra. I have seen several cases of this kind.

There is a species of incontinence of urine occurring in those persons who are disposed on the occurrence of any slight or trivial cause to become hysterical; this cannot strictly be called incontinence of urine, or paralysis, but it may be more properly considered as a deficient or false action of the mental power of volition. A young lady was once brought to me who was said to have incontinence of urine; when she sat in a chair, or walked out, the urine would come away involuntarily. As you may imagine, all remedies had been tried, but without doing her any good, and she had become quite a nuisance to her friends and family. She had been troubled with this complaint so long that her mental powers became depraved, and she began at last to forget the disgusting indecency of the disease under which she unfortunately laboured. I tried several remedies, but they were all unsuccessful, and eventually her friends tried what effect the feeling of shame would have upon her. She was sent from home to lodgings in the country; this made her feel very uncomfortable and unhappy, and in the course of a comparatively short time she recovered sufficiently to be allowed to return home, and in time she became quite well. Now, in a case of this kind, there was clearly a defect of the power of volition.

The incontinence of urine, which you sometimes find occurring in children, cannot be correctly termed incontinence. The child passes his water whenever he has a desire to do so, simply because of that desire, and because he has not been taught that it is wrong to do so; and he becomes in time so accustomed to it, that he continues the habit. If he is not properly corrected, he of course does not care about it, and thinks it a thing of no consequence. There are some boys and girls who pass their urine in bed every night; some do so occasionally, some only at night, and some only at particular hours of the night or day.

Dr. Prout says that this affection most frequently occurs in those children whose urine is very acid. This may be very likely the fact, but I have not found that purgatives and alkalies will cure the disease. Some surgeons are in the habit of applying a blister to the sacrum in these cases, with the supposed view of stimulating the nerves of that part; but I am disposed to believe, with Sir Charles Bell, that in these cases the benefit of the blister results principally from its preventing the patient lying on his back, which most children do who are troubled with this affection. I have been long struck with this fact, and have, in consequence, had an instrument made for these little patients to wear, with a piece of steel projecting from the back, so as totally to prevent the recumbent position in that situation, and I have met with some very good results from the use of it.

In many cases of this kind I have observed that children are in the habit of wetting the bed at a certain hour of the night, and in such cases I have observed that

dered that the child should be awake just before the hour arrived, in order to make water, and such cases have by perseverance recovered. If the child be at all advanced in age it is a good plan to have an alarm clock in the room, and the alarm set as to awake him just before the hour arrives. This affection is sometimes the result of a weak and debilitated constitution in childhood, and in such cases you will find that the sulphate of quinine will prove a valuable remedy, not only as a tonic, but as an antiperiodic; for I have told you that this affection will often recur at stated periods of the day or night. Although the cure of this affection is in some cases as simple as I have described it, you will nevertheless gain great credit with your patients where you can effect a cure. One thing, however, is quite necessary, and that is, that the child should wish to be cured.

CASES FROM THE EARLY NOTE BOOKS OF THE LATE

SIR ASTLEY COOPER, BART.

Extracted with permission of Bransby B. Cooper, Esq., F.R.S.

No. XVII.

STRICTURE OF THE COLON.

Mr. —, who had been afflicted for years with costiveness, applied to my friend Dr. Lister on account of languor, feebleness, and wasting. The Doctor ordered him an emetic, purges, and some bark, and he seemed to be getting better; but a few days after, on drinking a small quantity of wine, the debility came on in a greater degree than before, and he seemed, without any other striking symptom, to be sinking rapidly.

Three days after he had drank the above liquor, he complained of some slight uneasiness in his bowels and stomach. The debility now became extreme, and in a few hours he died.

Dissection.—On opening the abdomen, the small intestines, particularly the ileum, showed marks of inflammation. These marks were but slight externally; on cutting the intestines open, there was more appearance of inflammation found within. The stomach, which externally had no appearance of being inflamed, internally exhibited all the signs of that condition. There was a great deal of blood extravasated in the mucous coat.

The large intestines, as far as the sigmoid flexure of the colon, were of great bulk; on the outside they showed no inflammation, but within were much altered in colour owing to it.

At the sigmoid flexure there was an obstruction in the intestine, so that at this part it would not admit more than a large bougie through it. Just above this the inflammation was most strong.

The intestines, from the sigmoid flexure to the cæcum, were full of feces, and much enlarged.

About two inches below the upper stricture there was another, through which I could just pass my little finger.

REMARKS.

The above case is an instance of a very rare disease—namely, stricture of the intestine, and that independent of any scirrhus complaint.

It also shows that there may be much internal and little external appearance of inflammation.

The symptoms in inflammation of the mucous coat do not seem so strong as those from the peritoneal.

INJURY OF THE HEAD.

Mr. — had a fall from a chaise on August the 31st

by which he received a blow on the back part of his head.

Sept. 1. Has had a great deal of pain in the head all day long.

2. Pain in his head violent; his face is flushed; his skin sometimes hot; sometimes he is in a profuse sweat, then chilly; his pulse is very quick and full; the carotid arteries beat violently; he has pain shooting into his limbs at each pulsation; his muscles tremble; his breath is sometimes so short as to give him the sense of suffocation.

Bled to twenty ounces. Gave him calomel, three grains; castor oil, one ounce. Produced a copious stool in half an hour.

Evening. His pulse better; his head free from pain; he feels very languid; his blood is very sizey.

REMARKS.

Symptoms of irritation, which arise from an injury to a part very important to life, are much higher than those produced by a part of inferior importance.

Means to lessen them.—They require very early and active means to prevent their proving fatal.

In the foregoing case, if the violence of action had not been quickly moderated, suppuration would have taken place upon the brain. He was, therefore, bled till the pulse was found to be deprived of its fullness.

3. Mr. —'s pulse is still full and quick; he is often sick; he has a hot sensation in his ears; his eyes are painful to the light; he has some stupor; there is less chilliness and heat. Bled to twelve ounces; ordered a blister; to take a saline draught, with antimony.

4. Mr. — passed a tolerable night from having taken a grain of opium, excepting that his dreams disturbed him.

His pulse, however, is too full; it intermits; his urine is suppressed, either from the blister or the use of opium. He has had no stool since last night, nor has he vomited. Half an ounce of castor oil.

The least hurry of circulation produces great intermission of the pulse. Bled again.

5. Mr. —'s sickness has been less from the use of the opium.

His pulse, however, still occasionally intermits; his breathing is interrupted by sighs; he is restless; he lost the use of one arm and leg yesterday for a short time, and his sight for a few moments.

Restless, and his pulse fuller. Ordered to lose more blood, to fourteen ounces; ordered Dover's powder, ten grains.

REMARK.

Blood-letting.—If a person is bled when arterial action is very strong, a pulsating motion is observable in the blood from the vein.

6. Mr. — passed a good night from the use of the pulv. Doveri, and is much better this morning.

7. Mr. — much better. Continues the pulv. Doveri.

8. Mr. — much better.

9. Ditto.

INJURY TO HEAD.

17. A boy of about fourteen years of age came into the hospital last night, with a fractured skull, from the kick of a horse.

Pulse irregular; comatose in a degree, but impatient if disturbed. Bled.

This morning pulse quick; not so irregular as last night; great beating of the carotids; no stool; no urine; more comatose, but sleeps easy at present.

CONCUSSION OF BRAIN.

Killed a rabbit, by giving it a blow upon the head so as to produce violent concussion. It struggled violently in dying.