cases peculiar to women are about the average number yearly on the books. The number of cases of hæmorrhoids is larger than usual, owing to the wet and cold season before alluded to. This is, however, an affection not much met with in country practice, where the people are generally healthy, and have good air and a great deal of vigorous exercise. The number vaccinated was 240; all were successful, except two doubtful cases, and two not afterwards seen. Scrofula is less prevalent than heretofore; but syphilis is more so than ever, owing to the introduction of the disease into the country by the men employed on the survey at present going on in Ireland. It has extended itself in the line of the travels of the sappers and miners in the mountains and vallies of the whole country.

NEW MODE OF ACCELERATING LABOUR.

TO THE EDITORS OF THE PROVINCIAL MEDICAL AND SURGICAL JOURNAL.

GENTLEMEN,—If the following practical hint merit a place in your Journal, I shall be glad if you will give it insertion, as it is calculated to shorten the sufferings of the tender sex during parturition, and save much time of medical men who are in extensive practice.

I am, Gentlemen,

Your obedient servant,

SAMUEL STANILAND.

Fareham, Hants, March 14, 1842.

Being a midwifery pupil of Dr. Blundell, I of course carried his valuable instructions with me into practice, and one (not the least—viz., that “meddlesome midwifery is bad”) had much influence on my conduct when called to attend the parturient female. At first the unnecessary calls which were made for my attendance, and the frequently slow progress of labour in others, where the true expulsive efforts had commenced, together with the circumstance of being urged, on various occasions, to assist labour if possible, made me reflect upon the nature of the physiological action of the uterus, and the expulsive action of the abdominal muscles. Much thought on the subject, connected with the latter action on the rectum, bladder, stomach, and lungs, when irritating substances need expulsion, led me to the consideration of the process of labour, which appeared to me to be an analogous action, and I at length found the principle of imitation of the natural stimulant power, which I have now ten years carried into very successful operation, not only with a saving of much suffering to the patient, but an immense saving of time to myself.

When I first practised it, I was not so conversant with the physiology of the animal economy (except by book and school knowledge) as at present; yet, having studied Nature's designs in her various sympathies, the principle of excitation and reflex influences came to my assistance, and taught me that, as the bladder, when distended with urine, and the rectum distended with fæces, call upon the abdominal muscles to expel their contents, so would an excitement, similar to that of the head of the child, or the unbroken membranes on the vagina in their passage, promote, under proper circumstances, the natural efforts, and lead to a more speedy and equally safe labour.

The principle being given, what is the practice, and under what circumstances, should the operation be performed?—for I desire never to forget Dr. Blundell's valuable maxim.

The practice consists simply in imitating the influence of the child's head or membranes on the natural passages, and thus producing a reflex and

wholesome contraction of the abdominal muscles (I say nothing of the independent action of the uterus), by introducing the forefinger, or fore and second fingers, as far as the point of the os coccygis, and passing them along the whole surface downwards of the vagina, so as to give the sensation of distension, not pain, just enough to excite the required action, and give new and more vigorous impulse, even after hot fluids, stimulants, ergot, &c., have been tried in vain.

What are the circumstances, then, which warrant its application?

1. In cases of first labour, where the os uteri is fairly dilated, the head of the child has been in the passage from eighteen to twenty-four hours, the patient much fatigued, and the pains, without assistance, comparatively ineffectual, the womb acting.

2. In cases after the first child, where the passages are rather confined, or the head of the child unyielding, although in the passage, and the natural efforts unavailing, especially when warm fluids and stimulants have been tried without benefit.

3. Where there is much rigidity of the perineum, on which the head rests, but the natural efforts either greatly exhausted or the perineum unyielding; the practice here serves two purposes—viz., dilatation and stimulation.

Lastly. Where the womb and abdominal muscles, after a severe labour, sink into a collapsed state, and are indisposed to do more work, and the placenta, though in the upper part of the passage, does not excite the abdominal muscles to action. With this principle at my command, I have never seen such a thing as the old retained placenta in any form, and much doubt whether, with it, I ever shall.

There are other minor circumstances which it will be necessary to apply, but the observant practitioner will readily find them out when conversant with the application of the principle, and the above will be sufficient to guide him in most cases. If, however, with these directions, there should be any doubt on his mind, I would say, rather wait the result of the old practice, than urge assistance when unnecessary or “meddlesome.”

I shall feel much pleasure in hearing from any gentleman, either personally or through your Journal, the results of their practice, and the more so if they will favour us with new aphorisms for its general application.

Dr. Marshall Hall will at once perceive in this principle a practical application or illustration of the excito-motory nervous sensation; but it was not in consequence of his suggestion that the principle was applied by me, but after a study of the great sympathetic nerve, for I had practised it nearly five years before I read his excellent work.

CASE OF TRACHEOTOMY.

TO THE EDITORS OF THE PROVINCIAL MEDICAL AND SURGICAL JOURNAL.

GENTLEMEN,—I send you the following case of tracheotomy for insertion in your Journal, if you deem it of sufficient interest to occupy a place in its columns.

I am, Gentlemen,

Your obedient servant,

HENRY ALFORD,

Surgeon to the Taunton and Somerset Hospital.

Taunton, March 23, 1842.

William Marsh, aged 24, a tall-chandler by trade, July 22, 1841, was admitted into the Taunton union workhouse as a vagrant. He stated that he had been suffering some months from an affection of the throat, which had been brought on by cold, and had been

kept up by exposure and insufficient clothing; he had had a syphilitic attack four or five years since, but had not suffered from secondary symptoms.

His symptoms were—difficult inspiration, hoarse voice, frequent croupy cough, with some expectoration, redness, slight swelling, and some superficial ulcerations of the tonsils and fauces; the pulse was weak, and rather frequent; he was emaciated, and looked anxious and out of health. I gave him aperients and diaphoretics, applied blisters to the throat, kept him in bed, and restricted him to a milk and farinaceous diet.

July 29. About the middle of the day an attack of urgent dyspnoea came on, and rapidly increased; I saw him about five o'clock; he was then in great distress; the inspirations were performed with great difficulty and labour; his countenance was livid and very anxious. I gave him immediately an emetic composed of ipecac., twenty grains; tartar emetic, one grain; this was followed by copious draughts of warm water, but it did not produce vomiting; the dyspnoea increased with sonorous croupy inspiration; he was bathed in a profuse perspiration, and he was rapidly getting into a state of asphyxia.

I now proceeded to open the trachea, which I did about midway between the cricoid cartilage and the top of the sternum; the thickening of the integuments from previous blistering was rather an impediment, and I was somewhat embarrassed by a process of the thyroid body, but there was no hæmorrhage requiring a ligature; I attempted to use the tracheotomy trocar, having no other tube but the canula fitted to it, but I could not make the point of the stilette enter the trachea, as the cartilages bent under the pressure; I then opened the trachea with the knife, and introduced the tube alone, fixing it in the usual way. By the time this was done, although it only occupied a few minutes, the man had become insensible, and respiration had almost ceased, being only performed by occasional gasps. He now, however, quickly rallied, consciousness returned, and in a few minutes he breathed tranquilly through the tube, all the symptoms of dyspnoea and distress being relieved. For the first few days I had some trouble in clearing the tube of mucus, and was obliged several times to take it out for that purpose; but I had so much difficulty in replacing the tube, from its circular shape, and from the collapsing of the edges of the opening, that after a short time I contrived to clear it without removal, and in a week or two he was able to manage it himself. All the wound, except the part occupied by the tube, quickly healed, and various plans of treatment were tried, with the view of removing the disease and obstruction in the larynx. I applied blisters to the nucha and behind the ears, rubbed in iodine and mercurial ointments over the larynx, put him under a mild course of mercury, used purgatives pretty freely, and finally gave him tonics and a full diet; under this treatment his health improved, and he gained flesh, but he was still unable to breathe when the tube was closed more than a few seconds without distress; indeed, on one occasion, when the tube was removed for some purpose, and accidentally mislaid, he was in great distress from urgent dyspnoea in the four or five minutes that elapsed before it could be found and replaced, the opening having been nearly closed after a few inspirations by atmospheric pressure.

He left the workhouse October 8th, much improved in health and appearance, intending to go to his own parish, Bideford, in Devonshire.

I saw him again yesterday, nearly eight months since the operation; he was on his way to Bristol, where he intends working at his trade. He has worn the tube constantly; indeed, it has not been removed since he left the Taunton workhouse. He can now breathe by the glottis rather more easily, and for a longer time than he could, but is unable to do so without distress more than a minute or two at a time.

He has suffered no inconvenience or irritation from the presence of the tube, except for the first few days after its introduction. He has scarcely any cough, is in pretty good health, and is able to walk twenty miles a day without distress, except slight dyspnoea when walking against a hill.

PROVINCIAL

MEDICAL & SURGICAL JOURNAL

SATURDAY, MARCH 26, 1842.

It will be observed, by a reference to our notices of the proceedings in Parliament bearing upon medical questions, that the evils arising from the interment of the dead within the precincts of the metropolis, and other large towns, have at length attracted the attention of the legislature. The revolting abuses connected with this practice, arising partly from insufficiency of room in the burying-grounds, partly from other causes, have become known chiefly, if not entirely, through the instrumentality of Mr. Walker. To the publications of that gentleman's "Gleanings from Graveyards," and a pamphlet on the state of the burying-grounds of London, subsequently issued, we have before called attention. From the information collected, with much research, from various sources, and brought together in these works, it would appear that the indecent violation of the remains of the dead, in some establishments of this description, is almost without parallel either in ancient or modern times. Mr. Walker deserves the thanks of the public for the energy and zeal which he has displayed throughout the course of his inquiry into this subject; and it cannot but be highly gratifying to him that there is at last a prospect of substantial benefit accruing from his researches.

The question of the interment of the dead within the precincts of towns, and in the midst of a dense population, presents two separate and distinct features, each of which will be found to involve matters of deep import for reflection. This custom has its bearings as well upon the moral as upon the social condition of the people, and, followed as it now is, must prove deleterious both to the healthy state of body and mind. From the earliest periods, and amongst all nations, the disposal of the remains of those who were in life objects of veneration or regard, has been considered a religious duty. The Jew and the gentile, Christian and heathen, the man of social ties and civilised life and the roving inhabitants of the desert or the forest, have alike been accustomed to pay something like respect to the material framework, once animated and dwelt in by an intelligent essence similar to their own. The Hebrew, a passing sojourner in the land, seeks a place wherein he may bury his dead in safety out of his sight; the Egyptian endeavours to preserve the perishable form by curious arts; the Jew of later times commits the body to sepulture