WESTMINSTER MEDICAL SOCIETY.

Saturday, March 19, 1842.

Mr. H. J. JOHNSON, President.

PARALYSIS OF THE PORTIO DURA.—INFLAMMATION AND WASTING OF THE DELTOID MUSCLE.

There being no paper to be brought forward for discussion

Mr. STREETER drew the attention of members present, to some cases of paralysis of the portio dura from cold; he had on a former occasion brought three cases of the same kind before the society. Since Christmas, he had been called in to attend another case of a similar nature; it was very singular that many of these cases recovered by care and attention. The case to which he had alluded, had occurred in a child five years of age, who had been subject to fits. There was some swelling of the right side, and the nervous branches of the orbicularis palpebrarum were slightly affected. When called in, he at once gave his opinion that the cause of the affection was from cold. There was no pain in the ear, nor any cerebral disturbance. He viewed the case as one of inflammation of the nervous sheath, and treated it by leeches, poultices, fomentations, and warmth. The muscles had now recovered their functions. Many of these cases did not, however, recover as favourably. If the inflammation of the nervous sheath was in that part of it enclosed within a bony canal, it might be permanent. The child had now recovered for about five weeks, and for the last two weeks it had been quite well. The little patient had at one time shown a disposition to hypertrophy of the brain, which had been subdued by proper remedial means combined with change of air. As there was no specific subject before the society he would beg to draw their attention to the subject of whooping-cough, which, according to the recent returns of mortality, had been fatal in between 2,000 and 3,000 cases.

Mr. SNOW observed that, since Mr. Streeter had brought forward his case of paralysis of the portio dura, he had seen one case of the kind which went to prove the truth of Mr. Streeter's observation, that when uncomplicated with any organic alteration, this affection might be considered almost a harmless one. In the case to which he alluded, there were many dyspeptic symptoms present, and the orbicularis palpebrarum muscle was paralysed, the eyeball was kept constantly moistened with a warm cloth to prevent desiccation of the eye. To a question from Dr. V. Pettigrew, Mr. SNOW replied that there was no pain or enlargement of the parotid gland.

Dr. CHOWNE remarked that the general issue of these cases of paralysis of the portio dura was not so favourable, he only recollected one case in which the paralysis had gone off. Many serious evils resulted in these cases from the open eyelid; dust, air, and smoke would cause the eye to suffer very much. The muscles of the face were very liable in these cases to waste; the disease was a harmless one, and would not destroy life. He saw a case of the kind occur in an elderly man who was apoplectic. The diagnosis was easy, but the pathology was obscure, and he believed that it most frequently resulted from cold.

Dr. V. PETTIGREW observed, that these cases depended either on cold, or pressure, or the predominance of acidity in the stomach. The latter cause was a very common one. If a portion of bone pressed upon the nervous sheath in its passage through the Fallopian canal, deafness might be combined with the other symptoms, either from an inactive state of the auditory nerves, or from disease of the chorda tympani. He recollected one case in which paralysis of the portio dura and tic douloureux were caused by acidity of the stomach. He knew one case in which a spasmodic action through one of these nervous branches was

brought on whenever the levator labii superioris muscle was put into action; when the patient laughed, he was sometimes suddenly obliged to stop from the severe pain, which would last for several minutes; in this case he had advised the patient to have the muscle divided, but as he had a great horror of the knife, he would not consent to it. The effect of muscular action upon the arteries was very great; the arterial action was first quickened, the nervous power was then impeded, and lastly the circulation in the cutaneous veins became more full and rapid. It was a theory of Mr. Wardrop's, that the cutaneous veins were the safeguards of the lungs.

Mr. STREETER observed, that he should regret if any observations which had fallen from him should lead members to suppose that the state of a patient labouring under paralysis of the portio dura, was one devoid of danger; he believed that such cases were generally accompanied by inflammation, effusion, and pressure upon the nervous trunks. Such cases generally required a twelvemonth for their perfect removal. He generally watched such cases narrowly every day; if the prognosis was favourable, he left the case more to the power of nature; but, generally speaking, even these cases required the frequent application of blister to promote absorption.

Dr. CHOWNE considered that it was difficult always to ascribe a cause for these cases. The most common symptoms were those of an inflammatory character; the pain was sometimes very acute, and the paralytic state speedily followed it. The degree of immobility of the muscles of the face was very extraordinary; the paralysed side was still and stolid, whilst the other side was cheerful and animated; the immobility was greater than in ordinary cases of paralysis.

Mr. ELLIOTT inquired whether worms were ever met with in such cases, as tending to cause muscular irritability. He believed, with Dr. Pettigrew, that acidity of stomach, and derangement of the digestive organs might produce some of the phenomena of paralysis; he believed that the numbness of the arm alluded to by Dr. Pettigrew was of a very anomalous kind.

Dr. CHOWNE was desirous of knowing whether the President had met with any cases of paralysis of the portio dura of a permanent kind?

The PRESIDENT replied that the first case which he had met with, was of a physician of Naples, who, in travelling in a stage coach, had been exposed to a severe draught of cold air, which had brought on this peculiar form of paralysis; he went through the most approved plan of treatment, but in the course of a month he was very little better. In two other cases there had been inflammation of the conjunctiva, with great irritation of the eye, and the muscles of the face had wasted. In the third case the paralysis had been caused by the patient standing near an open window, and taking cold. The symptoms were subdued by leeches and purgatives. In neither of these cases was there any lesion of the brain or spinal marrow. He recollected a case in which there was inflammation of the sheath of the parotid gland in a gentleman subject to gout; depletion subdued this, and as the inflammation subsided the paralysis improved.

Mr. CHANCE alluded to a case occurring in the Middlesex Hospital, in which inflammation was supposed to be the cause of the paralysis; leeches and blisters were employed to subdue this; the result of the case, however, he did not know. In another case, which lasted four months, leeches, blisters, and purgative medicines had been employed; the case was much benefited by the treatment, but the recovery was not quite perfect. He alluded to a case formerly quoted by Mr. SNOW, in which severe and almost immediate stupor had followed the application of a grain of the acetate of morphia to a blistered surface; he (Mr. Chance) had never met with such rapid and decided effects from following the same plan of treatment; he believed that the endemic application of

narcotic medicines was frequently preferable to their administration by the mouth. He had administered as much as two grains of morphia in some painful nervous affections; yet even in this large quantity, relief had only been obtained after an interval of two hours.

Mr. SNOW remarked that, in the case to which he had formerly alluded, he was led to believe that the nerves of sensation only were affected, or, perhaps, the severe effect upon these nervous trunks might have caused a sympathetic action in other nervous branches.

Dr. A. B. THOMPSON had seen cutaneous ulceration ensue as a consequence of applying morphia over an abraded surface. In a case which was under the care of Dr. Graves, in Dublin, one-sixth of a grain of strychnine had been applied to a blistered surface, but no untoward results had occurred. He considered the application of morphia in this manner to be a very dangerous one.

Some conversation here occurred, in which Messrs. Chance, Snow, and Clarke, took part, which was terminated by the inquiry of the President, whether in Mr. Snow's case there had been any difficulty in raising the arm?

Mr. SNOW replied in the affirmative, adding, that the patient could not throw the arm forward without pain.

The PRESIDENT observed that he had been induced to ask the question, as he had seen several cases of a similar kind. One occurred to a navigator, in St. George's Hospital; many remedies had been tried, but without avail; moxa had been tried, and so effectually as to cause the patient to decamp. He had under his care a gentleman who, when with his regiment in Ireland, had been harassed by very severe and dangerous service, and exposed to severe atmospheric changes. The result of this was, that he one day found himself unable to move his arm. He could neither carry it backwards, nor raise it to a horizontal position. There was much pain accompanying motion, and the deltoid muscle was wasted. Pressure over the head of the humerus gave great pain. When first he saw these symptoms, he came to the conclusion that the sheath or fibrous tissue of the deltoid muscle was inflamed, and the circumflex nerve participated in the mischief. Blisters, leeches, colchicum and calomel, and opium, were employed, but the pain and wasting of the muscle continued. Mr. Keate saw the patient, and took the same view of the case; the same remedies were again tried, but without avail, and Colonel Coffin went down to Brighton. There he was under the care of Dr. Hare, who employed electricity—a remedy which he (the President) did not think an advisable one in such a case. No good, however, resulted from it; and now the other arm became affected, the same remedies (leeches, blisters, colchicum and calomel, and opium), were applied as in the former affection, and with somewhat better results; for though the muscle wasted, there was still some motor power present. The patient had thus lost the use of one arm, whilst the other was only moderately serviceable. In reflecting upon such cases as these, he was induced to consider that they required very strong and active treatment, which should consist of such remedies as those he had named. He did not use strychnine or morphia in these cases; he had never seen any good or bad effects from either of these remedies.

Dr. V. PETTIGREW observed that such cases as those described by the President were not uncommon. The guard of a mail coach was attacked with rheumatism, and was unable to raise his arm. All the means that had been previously named had been resorted to, but without any good effect occurring; but an old woman's remedy cured every symptom—it was the application of hot vinegar to the part for three hours, for seventeen successive days, at the end

of which time he was well. He had also a gentleman under his care with a similar affection, and in which the same remedy had succeeded. It was a custom at the opera, when the dancers sprained their ankles or feet, to plunge them into hot vinegar, which immediately relieved them.

After some further observations from Dr. Leonard Stewart, Mr. Snow, and others, the society adjourned.

PATHOLOGICAL SOCIETY OF BIRMINGHAM.

March 5, 1842.

FREDERICK RYLAND, Esq., in the Chair.

DISEASE OF KIDNEY.

A specimen of dilated pelvis of the right kidney, which had been sent to the secretaries by Mr. Hadley, was exhibited to the society. The person in whom the disease occurred was a man of 65 years of age, of spare habit, and suffered much from asthma; he had not been under any medical gentleman's care; he died suddenly on the 26th of February. On examination of the body on the 1st instant, effusion was found underneath the pia mater, which was thickened in patches. On the right side of the chest elongated adhesions were found in the pleura, and the lung was much engorged with dark blood, considerably softened, and did not crepitate on pressure. The right pleura was healthy. In the upper portion of the right lung there was a small cavity, the parietes of which were somewhat ossified; the lung itself was much engorged with blood. The heart was generally dilated. The mucous membrane of the stomach presented red and inflamed patches; the other contents of the abdomen were healthy. The left kidney was found healthy. The pelvis of the right was dilated so as to be capable of containing a pint of turbid urine, by which it was much distended. The substance of the kidney was nearly obliterated, but its artery and veins were of the normal size. The ureter of this kidney was described as being nearly obliterated.

Dr. FLETCHER believed that the dilatation of the pelvis of the kidney was the result of obstruction, existing in some part of the course of the ureter; and illustrated this position by a preparation, which he exhibited to the society, in which the ureter was obstructed by carcinoma at its inferior portion, which had caused dilatation of the whole of the ureter above it, dilatation of the pelvis, and absorption of the structure of the kidney; and he thought that death had taken place in Mr. Hadley's case from the absorption of urea.

FUNGOID DISEASE.—RUPTURE OF GALL-BLADDER.

A specimen of fungoid disease was brought forward, which had been sent by Mr. Ryan, of Sutton Coldfield. He had taken it from just above the left mamma of a female, aged about 37. It had been treated by caustics by an uneducated quack practitioner. Constitutional irritation, miscarriage of a seven months' child, and death, had taken place.

Dr. FLETCHER then brought before the notice of the society a ruptured gall-bladder, which had been presented to him, some months previously, by Mr. Hazlehurst, of Claverley. On July 3, 1841, the patient in whom it was found was taken suddenly ill with pains in the abdomen, after fighting, so as to render him unable to walk home. Peritonitis succeeded, and he sunk, and died in the afternoon of the 5th. There were no marks of external violence in any part of the region of the gall-bladder.

Mr. HONGSON said he had heard something of the case now before the society, from Mr. Hazlehurst, and believed that the gall-bladder was not healthy. He knew of another case, from which, and from experiments which were performed in consequence of