27. Has passed a very restless night ; shivering and constant sensation of cold, accompanied by cold perspiration : no increase of pain, but cough more difficult ; expectoration quite clear, frothy, and less viscid ; pulse 104, weak, and tremulous ; bowels natural. Continue pills. To have half an ounce of port wine, as required ; beef-tea, sago, arrow-root.

29. No more shivering ; feels more comfortable ; expresses a desire for animal food ; no pain of chest ; cough free ; pulse 84, much firmer. Continue pills. Mutton chop.

December 2. Continues to improve, and is apparently gaining flesh ; has slept well ; cough easy, and very rare ; expectoration much diminished, chiefly mucous ; respiration puerile in clavicular region of left side ; still entirely absent, with dulness on percussion in the lower portion of right side. Continue remedies.

5. Still continues to improve, and is evidently gaining strength ; pectoral symptoms ameliorated. Continue remedies.

8. Is improving, and sits up during the day.

16. Reports himself quite well, and wishes to leave the infirmary.

20. Has complained for the last three days of some increased pain, and to-day all the acute symptoms of pneumonia have recommenced ; cough frequent, small, and causes considerable pain ; expectoration brown, viscid, and copious, mixed with striæ of blood ; pulse 84, small, and soft ; tongue quite clean. Omit pills.

Tartar emetic, twelve grains ;

Water, twelve ounces ;

Tincture of opium, one drachm. Two tablespoonfuls every three hours, the dose being gradually increased to four every three hours.

A blister to the affected side.

22. Symptoms relieved.

It would be useless to continue the report of this case ; sufficient to say, this poor man had subsequently two separate attacks of pneumonia, and in both the symptoms were alleviated by the use of tartar emetic solution, but, in the last instance, not until vomiting was produced.

On the 12th of February (having apparently become nearly well) he was seized with symptoms of pleuro-pneumonia of the right side, which, however, were subdued by small doses of calomel and opium, blister, &c. On the 16th the pulse became suddenly weak and fluttering, the expectoration being of a putrid character, and the breath very fetid. On the 17th he died.

Autopsy Twelve Hours after Death.

On opening the thorax, the heart, left lung, and pleura, were found quite natural. The right pleura was almost universally adherent by strong bands of lymph and false membrane, and was with difficulty separated from the ribs and lung. It also contained a small quantity of dark colored serous fluid, and underneath the third rib a considerable opening, of a somewhat valvular appearance, was discovered, from which escaped dirty, unhealthy pus, of a putrid smell. This opening was connected with a small cavity in the upper lobe of the (right) lung. The whole of the right lung was in a hepatised condition, but more particularly the upper lobe, which sank very readily in water. Near this portion was found the abscess,

of a jagged appearance internally, and of the size of a pigeon's egg. The whole lung was beautifully covered with a dense false membrane, upon which were distinctly visible the impression of all the ribs, and the marks of which could not be removed by dissecting off the false membrane.* The intercostal spaces were observed projecting on the right side ; on the left they were natural. The bronchial lining membrane was much injected and congested. No trace of tubercular deposit could be found in either lung. Permission could not be obtained to examine the other viscera.

CONTRIBUTIONS

TO

OPHTHALMIC SURGERY.

By JOHN WALKER, Esq.,

Assistant-surgeon to the Eye Hospital, Manchester.

PRIMARY CAPSULAR CATARACT—COMPOUND OPERATION.

S. A. Hyam, aged twenty, applied to me on account of a defect of vision of both eyes, especially of the left, which latter was entirely useless. The affection had been coming on during the last twelve months, and had gradually increased in intensity. She had never had any inflammatory affection of her eyes, nor had they been the subject of any accident.

On examination, the eyes presented a generally healthy aspect ; but on looking into the pupil of the *left*, evidences of opacity of the capsule were distinctly visible, principally in its posterior hemisphere, which had a peculiar spotted or streaked appearance, perfectly white, stretching across the posterior chamber at a considerable distance from the iris. There was also a single white spot on the anterior hemisphere of the capsule, near its centre, and another about double the size, but not larger than a pin's head, immediately behind the former, and apparently seated in the outer lamina of the crystalline body, which was otherwise perfectly transparent. In the *right* eye the opacity was limited to a single white spot on the centre of the anterior hemisphere of the capsule. The vision of this eye was not sufficient to enable her to attend to her employment ; and as it appeared to be getting worse she was very desirous that something should be done for the restoration of that of the other. She was accordingly admitted into the hospital for the purpose of undergoing the usual operation.

June 25, 1842. The pupil of the *left* eye having been previously dilated by the use of the extract of belladonna, a straight needle was employed to perforate the sclerotica and break up the lens and capsule. A plaster of the moistened extract of belladonna was reapplied and the patient removed to bed.

30. The usual antiphlogistic treatment had been resorted to, and there had been but slight appearances of inflammation since the operation. The pupil had been kept well dilated and the lens appeared as a soft pultaceous mass, a considerable portion of it pressing forward into the anterior chamber.

July 2. For the last two days she had suffered much pain about the eye and head, and there was consider-

* Je n'ai jamais vu pareille chose, et je crois même qu'elle est impossible.—*Laennec Traité de l'Auscultation*, p. 397, édit. 3.

able vascularity of the outer tunics, the pink vessels predominating. Considering that the position of the lens in the anterior chamber and in immediate contact with the iris had excited and was calculated to keep up the irritation, and looking, also, to the future success of the operation, it was determined to extract the softened lens through a small aperture in the cornea. That operation was accordingly performed this day; the extraction knife was passed into the anterior chamber at the outer margin of the cornea, the point of the knife being carried into the centre of the pupil, the aqueous humor and more fluid portion of the lens escaping through the aperture; the scoop was then passed through the wound of the cornea into the pupil, when the remaining portion of the lens was evacuated, leaving the pupil clear, with the exception of a few fragments of opaque matter and a part of the capsule. Strips of plaster were then applied to keep the lids in contact, and the patient put to bed.

3. She had considerable pain yesterday after the operation, which continued all night, but was much abated this morning. The vascularity of the eye was not increased; the pupil was well dilated and a considerable part of it clear; she could distinctly observe surrounding objects, but owing to the irritation could not bear to exercise the organ. A little belladonna plaster was ordered to be applied, and a mild purgative administered.

9. The patient had gone on very favorably since last report, having had but little uneasiness, and the vascularity much diminished. There were still two or three fragments of opaque lens in the pupil and anterior chamber, but they had a soft flocculent appearance; and a large central aperture was visible in the opaque capsule.

Nothing untoward occurred after this. The opaque matter was gradually absorbed and very good vision restored, a small portion of capsule only being observed around the margin of the pupil when much dilated.

In this compound operation, as well as when the needle only is employed, we are generally called upon to resort to an after operation for the removal of the opaque capsule. It occurred to me that when using the scoop for the purpose of removing the more solid portion of the opaque crystalline, that instrument might in most instances be so employed as to make a large central aperture in the posterior part of the capsule, and thus obviate the necessity for ulterior proceedings. Such happened in this case. When the pupil is well dilated the end of the scoop may be passed through the pupillary aperture and so used as to perforate the posterior part of the capsule, which will afterwards contract to such an extent as to leave a sufficient opening for the admission of the rays of light, and consequently render unnecessary any operation for the removal of the capsule.

CAPSULAR CATARACT FROM INJURY—ABSORPTION OF CAPSULE.

Mrs. H., a married female, about thirty years of age, had lost the vision of her left eye in consequence of a blow received nine years before. A white pearly opacity was seen to block up the pupil, which latter contracted and dilated with the utmost regularity. No change was observable in the condition of the iris or any other texture of the eye, and she could readily

discern a light or any shining object. The case appeared to be one of capsulo-lenticular cataract, and on the 22nd of October last, at her request, I performed the posterior operation for solution, but, on endeavouring to break up the lens, I found that there was nothing but opaque capsule, which was very tough and parchment-like, the lens having evidently undergone absorption soon after the receipt of the injury before mentioned. I attempted to divide the capsule into fragments, in which I partially succeeded, and then pushed the entire body out of the axis of vision, into which, however, it very speedily returned. The moistened extract of belladonna was then applied over the orbit, and the usual antiphlogistic treatment enjoined.

The case went on favorably for the first ten days, but little inflammation having resulted, and, there being an aperture in the capsule, she had now regained some power of vision. At the expiration of this term, however, she began to complain of severe pain in the eye, and, on visiting her, I found that a rather sharp attack of inflammation had come on, and I was also surprised to find that the entire of the opaque capsule had come forward into the anterior chamber, lying irregularly across and nearly filling it, and that the pupil had contracted upon it. My first impression was that I ought to puncture the cornea and evacuate the irritating body, but the patient was exceedingly averse to any further operation, and had been very restless and unmanageable at the first, so that I decided on waiting to see if the opaque body would be acted upon by the aqueous humor and removed by the absorbents. I accordingly contented myself with the employment of leeches, blisters, and the ordinary remedies, with a view of diminishing the inflammatory action as much as possible in the mean time. These, however, did not seem of much avail, and the pupil remained closely contracted, notwithstanding the continued use of the belladonna, until a considerable portion of the capsule had disappeared, when the inflammatory symptoms gradually subsided and the pupil expanded. It was about two months from the time of the capsule passing into the anterior chamber before it had become entirely absorbed, and I carefully observed its gradual disappearance under the action of the absorbents. I may here observe that no mercurial preparation was given at any time during the progress of the case. The inflammatory action has now entirely subsided, leaving the pupil slightly contracted, but clear, and possessing the power of contraction and expansion under the varying influence of light. There is no other morbid change visible, except a slight central opacity of the cornea, the consequence of the inflammatory action having extended to that texture, and she is now regaining her vision. Here, then, was an instance in which the removal of a large portion, if not the whole, of the opaque capsule was indisputably removed by the process of absorption—a fact, the possibility of which has been disputed by most writers on ophthalmic surgery.