there was a tolerable breeze. The light thrown over the country can only be compared to that existing during a partial eclipse of the sun; this could not rationally be regarded as depending on the relative position of the sun, cloud, and earth, as the sun was nearly at its altitude whilst the cloud was nearly north-west. The electrical origin of cholera acquires further support from the well-known property of electricity, not only to temporarily localise itself, but also to pursue a particular and undeviating course in its changes from place to place. Here, again, if we compare the course of cholera, a striking analogy will be found. Let me now, in conclusion, pass to what I regard as pathological proofs of the justice of the view now advocated. They are the following:—

The sudden, and in some cases almost immediately, fatal impression made on the system, which had but a moment before been in a perfectly healthy and vigorous state.

The condition of the blood abstracted during life, more particularly as regards the slowness of its coagulation and its dark grumous appearance.

Finally, the semifluid state in which it is found on dissection after death—and, in many cases, the flaccid state of the muscular system, and peculiar discoloration of the skin.

If the above phenomena do not strictly accord with the effects produced by lightning in its mitigated and concentrated form, I confess myself at a loss where to look for analogies, or to find a justification for attaching any importance to them.

In my next, a few practical remarks will be offered on the treatment of the disease.

I have the honor to remain,

Your most obedient servant,

GEORGE FIFE, M.D.

Sunderland, Feb. 23. 1843.

OBSERVATIONS

ON THE

TOPICAL PATHOLOGY OF THE NEURALGÆ.

By Dr. BLACK, Manchester.

[From a Paper read at the Manchester Medical Society, March 1, 1843.]

The pathology of those severe afflictions—the neuralgæ—is, perhaps, the least understood of all the numerous diseases that are chiefly characterised by pain, and the inquiries that have been made into the matter have been more directed towards their constitutional and exciting causes than into the intimate or proximate lesions which constitute their essence and being. Without adverting to the literature on this important class of diseases, which is very amply found in the works of Hutchinson, Martinet, Swan, Elliotson, Teale, Halford, Piorry, Scudamore, and Valleix, with a few others of late date, I shall proceed to analyse the phenomena of the disease, and submit a solution of the intimate nature or topical causes on which neuralgia, especially of the intermittent species, more immediately depends.

The chief characteristic of the disease is pain, and that of a peculiar sharp, poignant, lacerating, or excruciating nature. The pain along the course of a nerve, or at its issue through some texture or other of the body, is also very often sudden in its invasion,

temporary in its duration, or intermittent, and receding or departing sometimes as quickly, but more frequently with a gradual decline. Its recession generally leaves the part previously affected somewhat numb, as if it were contused or bruised, and the integuments often suffused. If the complaint explodes itself on the face or head, the senses remain after an attack for a while blunted, and the powers of the encephalon more or less exhausted and enervated.

One great diagnostic feature of neuralgia is its intermittence, which is sometimes very regular, as much so as a normally constituted quotidian or tertian fever, leaving the body or part in the interval free from any pain or sensible derangement of health. At other times the occurrence of the pain is very irregular, or at most remittent, and when the affection is long established, the diagnostic pain is only marked by a periodical or indeterminate exacerbation. The intermittent pain may also be engrafted on a deranged constitution or a diffused state of chronic rheumatism or gout. The next observation that may be made regards the condition of the system under which this affection obtains; and again, the several parts, organs, and tissues, in which it elicits its ruling character. Preceding or accompanying this complaint we frequently witness several deranged states of the nutritive and chylipoetic organs and of the nervous system, such as mucoenteric complaints, biliary obstructions or vitiations, hæmorrhages, lesions or weaknesses of the cephalic, spinal, or ganglionic nerves, from mental or moral excitement, distress, or exhaustion. We moreover find that the complaint has originated from sudden exposure to severe cold; or to one less intense, conjoined with damp or malaria; and especially if the exposure have been topical, while the rest of the body has been warm, perspiring, or inactive. The condition of the parts affected are, however, more particularly the object of our observations, and to these we shall confine ourselves in the course of this paper, leaving the above enumerated subjects as being sufficiently recognised to appreciate the few simple but more precise data and inferences to be produced.

The principal parts on the head and face, where these peculiar attacks of pain are manifested, are—the supraorbital foramina or notches, the suborbital, mental, stylomastoid foramina, and the suboccipital region, the focus of which latter pain is between the base of the cranium and the atlas vertebra. On the trunk these pains are more particularly observed to be along each side of the spine, constituting spinal neuralgia; and again, midway along the periphery of the ribs, and next anterior to their osseous termination, giving rise to what are called intercostal neuralgia. These pains also manifest themselves in some transversely parallel lines over the abdomen, also towards the rectum, uterus, testes, and penis, on or about the knee from the anterior crural nerve, and also along the course of the sciatic nerve, extending even to the peroneal and plantar nerves. I may also advert to the irregular location of what are called the painful nervous subcutaneous tubercle, situated, according to my observation, more frequently on the trunk or upper extremities than elsewhere.

Our next inquiry is, What is there peculiar in these several parts or foci, whence irradiate those pungent or lacerating pains and other lesions of sensation and

motion? Anatomy shows us that at these points the several nerves pass through narrow bony canals, or between bones lined with hard cartilaginous surfaces, or else through apertures in firm aponeurosis or fibrous tissue, from all which they emerge upon the skin or outward integuments. Where such anatomical structure does not obviously obtain in the course of the affected nerve, we find that the composition of the neurilemma of the nerve itself goes far to supply the absence of any unyielding channel or aperture, through which, in addition, other nerves affected with neuralgia usually pass. It must, however, be remarked, that those nerves which proceed through osseous or fibrous canals are those which are more liable to be affected with intermittent neuralgia, or *tic douloureux*, while those which are less characteristically so exhibit painful affections of a more remittent or permanent type.

Another coefficient element in structure is, that those nerves which are more particularly afflicted with intermittent neuralgia, through whatever species of canal or aperture they pass, are accompanied at least with one artery and a vein, which are invested with little or no cellular tissue, and fill up with the nerve the whole canal or aperture, which they conjointly traverse. We have, then, a nerve, an artery, and a vein, passing together through a hard, unyielding, or firmly resisting, aperture, capable of no dilatation nor contraction, at least, in the absence of sufficient time, for molecular accretion or absorption to operate. The normal or physiological condition of the nerve and tissue of vessels is a relative and mutual adaptation and consentaneousness among the functions of the whole, permitting the free play of nervous action and sensation, and an undisturbed egress and ingress of the circulating fluids. Suppose, however, what has often been discovered,* that an osseous spicula or growth should intrude on the calibre of any of these bony canals or foramina, pressure on the nerve will follow, and according to the extent and mode of pressure so will the nature and degree of the pain or paralysis be. As the nerve cannot thus escape or be extended to one side, it must, from immediate pressure, be subject to a lesion of its functions. This special cause of neuralgia, therefore, requires no further notice of its structural pathology.

In the case, however, where the canal or aperture is perfect, and only lined with its periosteal, fibrous, or cartilaginous membrane, we must look either for some periodical or some more or less permanent causes of pressure on the transmitted nerves, the seat of the *tic* or the neuralgia. These causes, according to the character of the affection, may be physiologically attributed to the temporarily or chronically dilated diameters of the blood-vessels that are enclosed in the same unyielding canals or apertures along with the nerves, and of these vessels the arteries are those principally effectual in producing this pressure. We shall, for instance, take the infraorbital foramen, which is about one-tenth of an inch in diameter, and allot one-third of the area to each tissue—namely, the nerve, artery, and vein—all in the healthy state, performing their separate functions freely, and undisturbed by the action of one another; but when the artery, which is an elastic tube, happens to be

impaired in its muscular tension, from topical cold or any enervating cause, and especially if there is an increased *vis a tergo*, its diameter becomes disproportionately increased; and since the canal which it traverses is unyielding and resistant, the nerve will suffer its share of the encroachment and pressure from the artery. The extent of this pressure will occasion a proportional amount of pain, according to the sensibility of the nerve or of the constitution at large; and this pressure, for the time, acts in a similar manner to what would be induced by the intrusion of a harder body. The vein will also be impeded in its current, and give rise to fulness of its distal branches, and to integumental suffusion. As the nerve here instanced is one of sensation, the pain is often very exquisite and poignant, without paralysis, while such morbid categories occurring in the transit of the facial or *portio dura* nerve through the stylomastoid foramen occasion paralysis, without much or even any pain in many instances, this nerve being one principally of motion and expression.

The resolution of an attack of this intermittent neuralgia commonly takes place by an effusion of secretion from the nares and the lachrymal organs; while the climax or extremity of the pain reacts upon the brain, exhausts the nervous energy, and thus diminishes the propulsive action of the heart, by which the affected and offending artery is allowed to regain its normal elasticity and diameter, when all the several parts resume their healthy and reciprocating functions. The renewal of this morbid catenation depends, as the return of an ague, on the periodical enervation of the muscular power of the part or system, from the accumulation of debilitating impressions, conjoined with the recurring excitement of the heart from diurnal stimuli or from ganglionic irritation.

We have often very illustrative examples of this painful access, climax, and decline in attacks of toothache. The paroxysms of odontalgia are often excited by some debilitating cause, as cold or damp; and the pain becomes so gnawing, throbbing, and excruciating, with swelling and suffusion of the gums and jaws, that the dentist's aid is forthwith summoned for extraction. Before, however, the operation is performed, the fear of its infliction possesses the mind to such a depressing extent that the heart becomes weakened through the emotion, so that the pain is resolved and the extraction of the tooth is often delayed.

Now, in this special category, we have a carious or phlogosed lesion of the sensible tissue of the tooth, which is a stated cause of muscular enervation. We have also accompanying the nerve a nutrient artery, entering together with it at the apex of the fang to supply the tooth, all equiposed in their structure and functions in a state of health; but in these paroxysmal attacks the artery becomes debilitated and dilated, and yields to the normal impulse, and much more to any increased *vis a tergo* of the blood. The associated nerve is thus compressed and pinched; and though its denuded extremity may appear to be the seat of the pain, I believe the pressure on its transit, through the unyielding canal of the fang, gives that intermittent exquisiteness of suffering which is so often felt. If the acute pain depended solely on the morbid lesion of the extremity of the nerve in the crown, I do not see how the pain would become so

* Sir Henry Hallford's *Orations and Essays*, 1830; Dr. Abercrombie on *Diseases of the Brain*, 1834.

suddenly alleviated, as it often is in the circumstances above noticed. We are cognizant of no change that could take place in the nerve in such a short time, if it is not treated topically or otherwise by medicine; but the quantity and force of the vascular current admit of very sudden changes, arising from either physical or mental emotions, and we can easily appreciate them in all their extent, from syncope and fear up to the effects of joy and anger, as producing vascular activity.

The unbalanced physiological conditions which we have endeavoured to illustrate in the asthenic dilatation of the artery, and consequent pressure on the accompanying nerve in the bony canals above cited, will apply to all the other nerves and arteries similarly situated, and also to those points of transit between separate bones, as between the vertebræ.

In those cases where the affected nerves do not make their trajet through unyielding canals or apertures, as through the intercostal fasciæ at the points noted by Valleix, we infer the pressure on the nerve to arise from the vessels which surround and accompany them through these anatomical apertures; and in other cases, from their being enclosed in the same firm sheath or neurilema. These morbidly dilated vessels are often found in cases of sciatica—for instance, to consist of veins tortuous and enlarged from chronic dilatation. The pain is hence either persistent or far less intermittent than when the affected nerves traverse, together with the vessels, a bony canal. The vascular dilatation may become organic, and the disease be chronic from a deposit of lymph in the interstices of the nervous fibrils, or from a varicose state of the vessels themselves; all, however, contributing to the efficient cause of this peculiar pain—viz., pressure or counterpressure.

I shall now proceed to inquire how far the views which have been propounded are borne out by a just inference from the effects of our therapeutical measures in these disorders.

We have two classes of remedies which are generally administered in these affections, and found more or less alleviating, and, in many instances, productive of a cure. These are, irrespective of other preparatory or auxiliary means, narcotics and tonics. The beneficial action of narcotics, as in many other cases of pain, is derived from their direct agency on the *sensorium commune* or the nerve, by impairing or benumbing their sensibility to the injury inflicted by pressure, or by allaying the irritability of the heart or the muscular fibre in general, and thereby assuaging the abnormal impetus of the blood through the vessels that are previously weakened and implicated. But I would draw my chief corroboration of the views I have stated from the physiological effects of tonics—as bark, steel, arsenic, and mercury. If, not omitting the decided effect of ol. terebinth. in many cases of neuralgia, any of these are exhibited, in tonic doses, during the accession of intermittent neuralgia, they are found at the time rather to aggravate the pain than otherwise. This arises from their immediate action, by increasing the already over-excited sensibility, and unduly stimulating the heart, thereby urging a greater *vis a tergo* on the already atonically dilated blood-vessels of the part implicated, and adding, for the time, to the unbalanced condition of

things that exist, not to omit the resistance to absorption during this period of over-excited sensibility. When, however, these tonics are exhibited in the interval of a paroxysm, or of a severer access of pain, or while the system is less in a state of resistance, they not only serve to obtund the sensibility of the nerves of the *primæ viæ*, but, from being absorbed into the current of the circulation, they impart increased tonicity to the blood-globules, and greater contractility to the fibrinous coats of the vessels, so that the capillary circulation is made more active, and the diameters of the vessels are thereby diminished. Thus, pressure against or around the associated nerves is either moderated or provided against, in the event of any aberrations or disturbances in the system tending to reproduce neuralgic phenomena. The beneficial effect of the oil of turpentine very much comports with its remarkable property of increasing the capillary circulation, and contracting the coats of dilated blood-vessels.

That such is the principal curative action of these substances on the living animal fibre and blood-vessels, is what many observations have confirmed, and that the morbidly affected nerves are not directly influenced by their exhibition. If they were so, I do not see why their painful lesions of sensation or paralysis are not relieved by them during the paroxysms, as they are frequently by the exhibition or application of narcotics. Besides, we know that the gallic acid and tannin in the cinchona, and other astringent barks, have a great tendency to consolidate gelatinous matters, and that arsenic and mercury have also a determinate affinity for albumen, forming with it a firmer compound. Now, there is no doubt but that a diminished action from these substances, within the confines of healthy physiological life, will tend to consolidate and to give firmness, or what is called tone, to the gelatinous and albuminous matters in the tissues of the body.

Local bleedings, as by leeches, often alleviate the exacerbations of pain, but they seldom afford any permanent relief; for, as they only serve to take off the local pressure for the time, they impart no permanent tension to the vessels implicated. The benefit which blisters have occasionally afforded seems to proceed from their direct stimulus to the vascular fibre, producing contraction of the dilated vessels; but as no invigorating power is permanently imparted, their utility is temporary, if they are not from time to time renewed.

The idea that the morbid affection originated in, and was confined to, the nerves themselves, and their branches, formerly gave rise to cross sections of the nerves, with the view of cutting off the communication with the brain. It is needless to say how often disappointment followed these incisions, however completely performed, because the seat of the neuralgia was not in the distal branches of the nerves, but somewhere central to them in the course of some of the bony or unyielding outlets of the cranium or spinal canal, if not in the brain itself. The operation spilled a little blood, and so took off the temporary pressure, while some irritative stimulus was given to the vascular tissue around, but no molecular or vital tone was imparted; on the contrary, after the stimulus of the operation was over, the paroxysms of

pain were often renewed with more violence than ever, the lesion which the parts had suffered only adding to the previous enervation and debility. We have no correct idea how mercury or arsenic, taken internally, acts upon the affected or other nerves; but we can fairly observe and appreciate their effects upon the blood-vessels, by their increasing capillary action, and giving more contractile vigor to congested and dilated vessels. This energetic effect relieves the associated nerves from pressure, wherever situated; but it will be more palpably displayed where the vessels are closely bound up or imprisoned with the nerves. The benefits resulting from moxas, or perpetual blisters, seem also to be owing to their continued diversion of the fluids from the subjacent or neighbouring vessels that are injuriously pressing upon the affected nerve, and not from any direct curative action which they have upon the nerve itself. It is also to be remarked, that in the great majority of cases of neuralgia the pain is very much increased, if additional pressure be made by the finger at the several points affected, where the nerve is emerging from a bony or other unyielding aperture, thereby corroborating the doctrine that a less pressure is the cause of a lower degree of pain. A severer pressure by the finger will also, in some cases, produce numbness and insensibility.

These, then, are shortly the views which I beg to offer concerning the topical nature of our well characterised neuralgiæ, especially those of the intermittent type. Irrespective of constitutional or other derangements, I conceive that they primarily consist in a hyperæmic and an atonic dilatation of the blood-vessels, which either surround or accompany the nerves affected through unyielding canals or apertures in bone, cartilage, or else through fibrous sheaths or fasciæ of more or less density or resistance—that the pain is occasioned by pressure or counterpressure, varying according to the phlogosis or fulness of the dilated vessel—that the intermittence depends on the diurnal or periodical nervous exhaustion or accumulation of excitability in the system—and lastly, that the above pathological view is confirmed by the effects of such tonics as bark, steel, arsenic, and mercury, taken internally, the action of all which is principally confined to the vascular system.

EFFICACY OF VACCINATION.

[The following letter, from the superintendent registrar of the parish of Birmingham, has been forwarded to us by Dr. Edward Johnstone. It contains some important facts relative to the effect of proper vaccination in preventing the occurrence of small-pox, and is, moreover, very creditable to the zeal of its author.—Eds.]

DEAR SIR,—You are aware that in the month of July, 1840, the royal assent was given to an act, 3rd and 4th Vic., c. 29, “to extend the practice of vaccination.” By that act, guardians of the poor in every parish are directed to contract with legally qualified medical practitioners for the vaccination of all persons resident in their respective parishes; and to remunerate such medical officers according to the number of persons who, not having been previously successfully vaccinated, shall be successfully vac-

nated by them. By the same act, any person convicted of producing, or attempting to produce, by inoculation or any other means whatsoever, the disease of small-pox, is subjected to imprisonment in the common gaol or house of correction for any period not exceeding one month.

The guardians of the poor of the parish of Birmingham were not slow to obey the injunctions of the Vaccination Act; they made arrangements for commencing operations with the year 1841 in a most efficient manner. Of the importance of the Vaccination Act to the welfare of the country at large, I presume there can scarcely be a difference of opinion; of its importance in the parish of Birmingham, and the necessity for its being promptly acted upon by the guardians of the poor in this parish, my position, as superintendent registrar, afforded me an opportunity of judging, which was not so open to other persons. At the time I am alluding to I deemed it prudent to confine my observations on the subject to a few professional gentlemen, lest an unnecessary alarm might be excited; it may now, however, be useful to record them.

During the quarter ending December 31, 1840, small-pox prevailed here to what appeared to me a frightful extent; it amounted to full 11 per cent. of the entire number of deaths in the parish; and in the most central district in the parish, out of ninety deaths in the same quarter, twenty-one deaths, or 23½ per cent., were registered as caused by small-pox. From the 1st of January to the 7th of February, in 1841, nine deaths from small-pox were registered in the same district, being equal to 20 per cent. of the whole number of deaths in that district during the same period; by March, however, this plague seemed to have been stayed. I now proceed to my chief and pleasing object in making this communication, which is to notice the apparently very successful result of carrying out with vigor the provisions of the Vaccination Act in this extensive parish.

During the first three months of 1841 more than 700 cases were successfully vaccinated by our medical officers, and the number so vaccinated in the whole year 1841 was 1481; in the year ending December 31, 1842, the number so vaccinated was 850.

Now, Sir, you will learn with pleasure that in the whole year, ending 31st December last (1842), only 10 deaths were registered in the entire parish of Birmingham as caused by small-pox, being about 1 in every 358 deaths instead of 1 in every 9 deaths, as was the case in the quarter ending December 31, 1840, to which I have before alluded. And I desire further to notice that not one of the ten deaths from small-pox, registered in the year 1842, has occurred since the month of April last, so that for many months it would appear that no death has been caused by small-pox in this parish. I will not, as a non-professional man, indulge in conclusions, or attempt deductions from these facts; I remember the adage, “Do not shout till you are out of the wood;” but surely there is very great encouragement to guardians of the poor, to the medical profession, to parents, and the public in general, to persevere in a course promising such important results, and to keep up a kind, although by no means a disinterested, watchfulness over the families of their poorer neighbours.