supervene, but after many days of anxious suspense they subsided. He did not, however, regain his strength for several months, but at length he got quite well, and had the perfect use of his limb without any lameness.

Some difficulty arose in passing the catheter from the position of the patient, and still greater in preventing the urine from running about the bed, which inconvenience was remedied by inserting the point of one elastic catheter into the mouth of the other. By thus lengthening the tube and bending it over the thigh, the contents of the bladder were received in a basin, and future discomfort avoided.

CASE II.—A robust young woman was thrown from a horse, and suffered a compound fracture of both bones of the leg, about three inches above the ankle. Nine months after the accident she was placed under my care. The fibula only had united, and a piece of detached bone was wedged firmly between the broken ends of the tibia, which prevented any union. Her health had suffered much from long confinement, and amputation had been proposed, which she refused, but was willing to submit to any treatment which offered a chance of restoring her limb. I divided the integuments over the tibia to the extent of four or five inches, and sawed off the ends of the bone with Mr. Hey's saw, which enabled me to remove the intervening portion. The operation was more tedious than painful. In the evening she was attacked with erysipelas, which spread rapidly over the thigh and leg, and was accompanied with constant sickness and vomiting. Half a grain of opium with an effervescing draught every two or three hours, allayed the irritation of the stomach, and the application of a spirituous lotion lessened the inflammation of the limb so much, that in two or three days the immediate effects of the operation had subsided. The wound soon healed, but there was a very considerable space between the ends of the tibia, and the limb was still useless. A case made to fit the leg, of the same kind of leather as is used for the flaps of saddles, strengthened by some narrow plates of iron, and fastened under the knee and around the ankle, was applied, which supported the limb so effectually that she was soon able to walk about with the assistance of a stick only. The ends of the bones gradually approximated each other, so as to form a firm union and give her a useful limb.

C A S E

OF

NEURALGIA OF THE INFERIOR DENTAL NERVE.

By JOHN WATERS, M.D.

A lady, aged about forty, of an anxious temperament, came to town last August to consult me concerning an affection of the lower maxilla, from which she had been suffering during the last three years. She described it as coming on slowly at the time mentioned, and considered the pain as the result of toothache; but on consulting her medical attendant, he pronounced it to be a case of neuralgia of the inferior dental nerve. She underwent a variety of treatment, but with little or no permanent effect; tried change of air, but to no purpose; and had six or

seven teeth extracted, in the hope of experiencing relief, and also as some suspicion was entertained that the fang of a tooth might be diseased. Her sufferings still continuing on the increase, at the recommendation of a friend, whom I had attended some short time before, she came to London to see me.

Her general health seemed much impaired, from the excessive pain that she had been laboring under so long; countenance expressive of anxiety; catamenia regular; appetite in general not impaired; not subject to headache; never had rheumatism, but had considerable anxiety and fatigue in her attendance on her husband previous to his death; the pain, which was of the most excruciating nature, was solely confined to the ramus mentalis of the inferior dental nerve as it issues from the foramen, and could be almost covered with the point of the finger, but about two years previously it extended along the entire course of the nerve; there was no swelling, heat, or discoloration of the part, except a slight thickening of the skin from the different topical remedies that were previously applied, and the incessant friction and pressure that she made use of to allay the paroxysms of pain; there was no permanent muscular contraction. When I first saw her the pain was so incessant (coming on about every three or four minutes) that her hand was continually pressing the part; she described it as an insupportable dragging pain, a kind of electric commotion, which, without precursory signs, arrived at its maximum of intensity, and as suddenly departed; and sometimes these lancinating pains arrived so rapidly at their highest degree of development that she was forced to scream aloud; at each attack there was an involuntary movement of the whole body, but pain was only felt in the part described. She had tried the different preparations of iron, morphia, veratria, blisters, leeches, and the usual category of remedies employed in similar cases, without experiencing relief. Prescribed the following pills:—

Strychnine, one grain and a half;
Ioduret of iron, twenty-four grains;
Extract of conium, twenty-four grains. Mucilage enough to make into twelve pills; one twice a-day.

Also applied a small blister over the foramen mentale, and dressed the abraded surface with the following ointment:—

Hydrochlorate of morphia, sixteen grains;
Suet, one ounce;
Hydrochloric acid, three drops. An ointment to be applied thrice a-day.

On continuing this treatment for a week, I was obliged to administer the pills but twice in the day, as the twitching from the strychnine was much complained of. She described the pain as now totally changed since she felt the twitching in the affected nerve, been more lancinating, but the intermissions were longer and the duration of pain somewhat abridged. She also took five grains of diluted sulphate of quinine each day at noon.

Extract of aloes, fifteen grains;
Sulphate of quinine, twenty grains;
Sulphate of iron, eight grains;
Dilute sulphuric acid, eight drops. Make into eight pills; one twice a-day.

Having continued this treatment for fifteen or sixteen days with little improvement, I detailed the case to my friend Dr. Hennis Green, and on his recommendation I adopted the following treatment, which was eminently successful :—

I gave the arsenical solution twice in the day, commencing with five drops, and gradually pushed it to forty. On the sixth day there was a marked diminution in the paroxysms, and the pain was now so much mitigated that she ceased to press or rub the part as she was accustomed to do during the paroxysms; her general health began to improve; sleep not disturbed; and within the month from the time of commencing the treatment she was entirely free from pain; and last week I received a letter from her, wherein she states that there has not been a single recurrence of it.

62, Torrington-square,
Dec. 15, 1842.

PROVINCIAL MEDICAL JOURNAL

SATURDAY, JANUARY 28, 1843.

In our last Number we published an important letter from Mr. R. Cartwright, Secretary to the Shropshire Medical Association. We call this document an important one, inasmuch as it furnishes evidence that medical practitioners are at length becoming sensible of the truth conveyed in the ancient, though by them long-neglected maxim, "Union is strength." There is no class of men for the protection of whose interests unity of purpose and action is more imperatively requisite than the members of the medical profession; yet our want of union in public and the discords which distract us in the relations of private life have become almost proverbial. It were easy to trace the causes of this absence of harmony to their real source. Medical men are not of themselves more disposed to jealousy and strife than other classes of the community; their faults are not of their own creation; they have been forced on them by the vicious constitution under which we live—by the selfish views of the medical corporations. Instead of working together for the common good, and endeavouring to reconcile conflicting interests, the corporate and chartered bodies study exclusively their individual interests, leaving the commonwealth of medicine to look after itself. This selfish and short-sighted policy is daily becoming more prevalent; it has extended from one extremity of the land to the other. In the North we have an ancient university conferring degrees for the sum of twenty-five shillings; in the South we establish a counterpoise by deteriorating the quality of the material to the lowest standard. Surgeons are now sent forth from Lincoln's Inn something after the fashion of printed goods from Manchester. Without

the slightest regard to the state of the market, or the least consideration for the quality of the article supplied, surgeons and stuffs are manufactured merely to keep the mill going. Hence are we like lean kine, ready to devour one another. Physicians and surgeons, emanating from different sources, and forced to submit to the factitious distinctions imposed on them by their respective governing bodies, are on about the same terms of friendship as cat and dog. The practice of medicine is cast between them as a bone of contention.

The surgeon, again, looks with jealous eye on the general practitioner, while the apothecary wages open and deadly war on his neighbour the druggist. Even the sub-species of doctors, surgeons and apothecaries, are split up into factions and arrayed against each other. Glasgow and Edinburgh are as much opposed to each other in the manufactory of doctors as Calais and Boulogne ever were in the importation of passengers. The holder of a certain Scottish degree is treated almost as a Jew. Oxford and Cambridge, indeed, seem resigned to their fate; but the Pall Mall serpent, like the Egyptian one of old, has suddenly raised up its head, and threatens to devour all competitors.

Under such a state it is impossible that anything save discord should prevail in the medical profession. We therefore look with exceeding satisfaction at any signs which indicate that medical men may be induced to unite together for the redress of injury or the attainment of good.

The successful opposition of the united profession in Ireland to the scheme of the poor-law commissioners is an encouraging example of what may be effected by zeal and unanimity in a good cause; and we look with almost equal pleasure on the union of the Shropshire practitioners in the assurance offices affair. Of 115 medical practitioners resident in Shropshire, no less than 112 have signed a pledge "not to answer the inquiries of assurance offices unless accompanied by a fee of one guinea."

We have already expressed *our own* opinion upon this point, and shall not recur to it here; one sentence, however, of Mr. Cartwright's letter contains a hint which may lead to a satisfactory adjustment of the question. It is conceded on all hands that the medical man should be paid for his certificate; but the rub is, who is to pay him? One party say the assurance offices, the other would throw the onus on the person applying for the certificate. With this paltry dispute the medical practitioner is mixed up in an unnecessary and unseemly manner. The question should be settled between the office and the applicant. Let every one who makes an application to an assurance office be compelled by the company to deposit the sum of one guinea for the purpose of obtaining a medical certificate, and let the offices arrange with