fourchette, so as to depress it as much as possible, whilst on each side the index and ring fingers are employed to keep the labia separate. The body of the speculum is seized in the right hand, and its extremity presented in a perpendicular direction to the back of the middle finger, on which it is to be made to rest. The instrument is then slowly brought to a parallel direction with the vaginal axis, its diameter being rendered slightly oblique, so as to avoid the meatus, and it will then glide on without difficulty; the handle is then placed in the patient's left inguinal region, and confided to her care.

In this position the valves correspond not to the lateral, but to the anterior and posterior walls of the vagina, so as effectually to prevent protrusion of the mucous membrane, which might otherwise occur between the valves, and interfere with the investigations.

To clear away the secretions of the part, a little instrument may be used, which I here present to you, and which I have invented to replace those disgusting little wooden brushes at present used for the same purpose. It is merely a three-pronged forceps, which opens or closes by means of a ring, and which can be loaded with cotton, and fixed on the female catheter, to increase its length.

A surgeon, Gentlemen, who desires to succeed in practice should be very neat in his instruments; in them, particularly, perfect cleanliness is indispensable, and should be combined, if possible, with elegance. I recommend, therefore, the use of gilt specula; they do not become stained like common tin instruments, their aspect is much more pleasing, and their cost not very considerable. Besides, they keep a long time, especially when they are enveloped in a velvet bag, embroidered, like this I now show you, by the fair hands of a grateful patient.

ON THE RAPID CURE OF GONORRHŒA.

TO THE EDITORS OF THE PROVINCIAL MEDICAL
JOURNAL.

GENTLEMEN,—I shall be obliged if you will permit me to contribute the results of my experience in the treatment of gonorrhœa to our general fund of knowledge on the subject.

From the unsettled opinion of the profession in the treatment of this complaint, no system has been proposed which the practitioner could regard with confidence and satisfaction. Cubebs, copaiba, liquor potassæ, tartar emetic, iodine and its preparations, injections of zinc, nitrate of silver, have each in their turn been vaunted as a specific, or greatly praised for the relief it was capable of affording. Yet, on none of these could much dependance be placed; they have all disappointed the practitioner, for, however valuable they may be in themselves, and however especially local their action, they generally fail to effect a speedy cure, from the variety of symptoms which usually attend the severe forms of the disease; and the reason is obvious: gonorrhœa can be divided into three distinct stages, and, if so, the treatment must undergo a similar division. Now, with the exception of the tartarised antimony and iodine, the whole of the medicines mentioned act in a similar

manner on the genito-urinary mucoid surfaces. Beyond a certain point, they fail to produce a beneficial effect, and, therefore, are not at all adapted to meet the new state of things, which the complaint, together with their action, has produced.

During the period when the Great Western Railroad was in the course of formation, the larger proportion of illnesses from which the "navigators" suffered consisted of gonorrhœa, and other forms of the venereal disease. The dissolute habits of these men caused the inflammatory symptoms to rage with great intensity; and, without acting on any "system," I was obliged frequently to resort to very energetic measures, and whenever sharp antiphlogistic means were adopted, before exhibiting the *usual* copaiba and liquor potassæ mixture, the complaint invariably ran a milder as well as a shorter course. This was but in accordance with the first principles of the practice of medicine; and it seemed, in my humble opinion, only necessary to make the whole treatment accord with such principles, in order to subdue the disease at once safely and rapidly.

Before the plan is submitted to the profession, I must own that the following combination of medicines—very similar in appearance to a nostrum called the "specific solution"—produced a cure, although one neither so speedy, *scientific*, nor safe, as the measures which I shall presently bring forward:—

Hydriodate of potash, one drachm;
Oil of cubebs, half an ounce;
Balsam of copaiba, half an ounce;
Liquor potassæ, one ounce;
Hydrochlorate of morphia, two grains;
Spirit of nitric æther, two ounces. Two teaspoonfuls to be taken three times a-day in water.

The diet moderate, and an occasional calomel pill and draught administered. The cure was generally completed in ten or fourteen days. Occasionally a gleet discharge remained, which a weak zinc injection was sufficient to remove.

Satisfactory as were the effects of this combination of medicines, the treatment was not exactly what the state of our science demanded. It was no system; and, therefore, as the lines of demarcation between the inflammatory, suppurative, and atonic stages were sufficiently distinct, there was every reason to conclude that, by anticipating or attacking them with appropriate means when developed, the best results would follow.

The treatment was consequently divided into three parts; the first had reference to the inflammatory stage, the second to its effects, and the last to the atonic state of the lining membrane of the urethra, and the vessels with which it is supplied. It consisted of ten grains of calomel, divided into three doses, one to be taken every four hours, and of the following mixture:—

Sulphate of magnesia, half an ounce;
Jalap, two scruples;
Scammony, fifteen grains;
Potassio-tartrate of antimony, three grains;
Water, eight ounces. Three tablespoonfuls to be taken every four hours, beginning an hour after each powder.

Diet low, and drink barley water, linseed tea, &c. The first dose of the mixture was generally rejected,

but the stomach soon became accustomed to bear the tartar emetic. The genital organs to be frequently soaked in tepid water, and suspended. In four and twenty hours the suppurative stage may be anticipated, if not already established. The mixture to be omitted, and the following substituted :—

Liquor potassæ, one ounce;
Oil of cubebs, half an ounce;
Balsam of copaiba, half an ounce;
Hydriodate of potash, half a drachm;
Hydrochlorate of morphia, two grains. A teaspoonful to be taken in two ounces of barley water every four hours.

The parts to be rolled with a bandage, and kept wet with an acetate of lead lotion; barley water to be drunk very plentifully, and the diet improved. This treatment to be continued for three days; on the evening of the third a dose of calomel to be given, and a gentle aperient the next morning. The following injection may now be used every two or three hours for two days :—

Acetate of lead, one drachm;
Water, eight ounces. Make an injection.

This powerful lotion speedily arrests the remains of the secretion, and appears to restore the lining membrane to its normal condition. The treatment is now merely palliative, comprising a dose of the drops once a-day for a few days, and an injection of cold water twice every second day for the same time.

In no instance has a single unpleasant symptom arisen, nor the slightest contraction of the urethra supervened; and, unless the patient is very rebellious, and will drink spirits, beer, or stout, the cure is completed within the week.

The above plan was modified according to age, temperament, and symptomatic violence, but the principle in every case was the same. In my next paper I shall have much pleasure in demonstrating how rapid is the cure of chancre under the power of nitric acid, tincture of iodine, and a very slight course of mercury.

I am, Gentlemen,

Your obedient servant,

CONWAY T. EDWARDS, M.R.C.S.L.

Bath Easton, July 10, 1843.

TREATMENT OF UTERINE HÆMORRHAGE BY GALLIC ACID.

The following cases, showing the value of gallic acid as a remedy in cases of uterine hæmorrhage, are abridged from a paper by Dr. Stevenson :—

CASE I.—J. Johnston, aged thirty-two, plethoric, having menstruated irregularly for two years. On January 18, was attacked with pains in the abdomen, continuing at intervals until the next morning, when she passed an unusually large quantity of blood *per vaginam*. At eleven o'clock she was lying on her back in a very exhausted state; eyes sunk; pupils dilated; hands and feet covered by cold perspiration; abdomen tense; tenderness in the hypogastric region; about a pound of blood had passed from the vagina. Cold to the hypogastric region, castor oil, and frictions to the legs, were the remedies employed, and on the next day the pains had ceased, but the countenance

was more anxious and paler than before; the pulse 110, regular, feeble, and scarcely perceptible at the wrist. Blood has continued to escape from the vagina throughout the whole night, in quantity not less than twenty ounces. A glass of port wine was given, and ordered to be repeated every four hours, and the following :—

Gallic acid, eight grains;
Powder of cinnamon, five grains. Make a powder, to be given every third hour.

Three, p.m., of the same day (19th). Great exhaustion; pulse 130, feeble; fits of fainting; six ounces of blood, of highly offensive smell, had been passed since the morning. Wine omitted; powders to be continued.

20, Nine, a.m. Pulse 110, stronger and regular; countenance flushed; heat of skin moderate; three ounces of blood passed since yesterday, not offensive, and having been expelled in small clots. Seven powders had now been taken. Ordered to repeat them at large intervals.

Five, p.m. Pulse 120, feeble; countenance more anxious; hæmorrhage decreased; smell not so offensive. Twenty-four minims of laudanum were ordered to be given at bed-time, and the powders to be continued.

21, Twelve, a.m. Pulse 110, full and regular; passed a quiet night; countenance animated; a few drops of blood passed during the night, but towards morning about half a drachm; skin warm. Repeat the powders.

After this period the powders were given once daily; the hæmorrhage did not recur, and in a few days the patient was convalescent.

CASE II.—Mary Slater, aged thirty-six, was seen on February 2. This patient had for three years menstruated more or less irregularly, the secretion having occasionally been profuse. Six months ago had had an attack of hæmatemesis, since which time the catamenial function had been suspended.

At the time of her being visited, on February 2, she presented the following symptoms :—General uneasiness, fulness in the head, lassitude, pains in the lower part of the abdomen, and a discharge of blood had taken place from the vagina, in quantity about a pound. She was unconscious; pulse quick, feeble, and irregular; respiration quick; heart's action feeble, accompanied by an indistinct *bruit*.

Nine, p.m. Reaction had taken place by means of the remedies employed; but two ounces of blood, mixed with offensively smelling mucus, had passed from the vagina. Symptoms of collapse again occurred, but were successfully met by appropriate remedies, and the following prescription was then ordered, plugging of the vagina having been previously had recourse to :—

Gallic acid, eight grains;
Sugar, ten grains. To be taken every third hour, and half a glass of sherry every second hour.

3, Ten, a.m. Pulse 120, small; respiration easy; plug came away from the vagina at eleven o'clock, covered with coagulated blood; hæmorrhage to the amount of six ounces since the last report. The powder to be continued.

Seven, p.m. Pulse 100, small, regular; countenance anxious; much thirst; gnawing pain in the