MR. FLETCHER'S REPLY TO "CLIO."

TO THE EDITORS OF THE PROVINCIAL MEDICAL JOURNAL.

GENTLEMEN,—As I state facts, and facts only, and have been in extensive midwifery practice upwards of thirty-eight years, after studying under the first accoucheurs of the day—Dr. Osborne, Dr. John Clarke, and Dr. Haighton—I felt surprised at my veracity being called into question by Clio.

Controversial statements in any way are unpleasant, but most so when anonymous and vindictive; which latter term I think duly applied to your correspondent for his attempt to misrepresent the facts of my case of difficult parturition; which misrepresentation he has thought right to strengthen by calculations in no way applicable to substances which are compressible, but to bodies hard, thick, and non-resisting—such, for instance, as must be his own well ossified cranium. How Clio could make out that the body and arms of the first child were contained in the vagina from my statement, that "the feet of the child presented, which with the body and arms were brought down, but in trying to get my fore finger into the mouth of the child, I found the apex of a head in the pelvis preventing any further delivery," is only to be accounted for by the thickness of his comprehension, or of his complete ignorance of the process of parturition, or of the manual operations made use of to facilitate the exit of the child; for it is very far from probable that an accoucheur would try to get his finger into the mouth of the child when its body and arms, as Clio supposes, were in the bony pelvis; indeed, the operation described ought to have indicated to your correspondent the exact stage of parturition—namely, that the body and arms of the child were born; that the head was supposed to have descended, with the face towards the cavity of the sacrum, into the bony pelvis, and only required the manual operation, which all accoucheurs are acquainted with, of putting the finger into the mouth of the child, and drawing the face into the convexity of the sacrum, to accomplish the exit of the head under the most favorable circumstances. But notwithstanding Clio's knowledge of the measurement of the dry, articulated pelvis, he must have forgotten the measure of the anterior wall of the symphysis pubis, the part against which the first child's neck rested, whilst the head and body of the second child passed by it in the bony pelvis; a state which every one must have understood from my description, though short, unless his head was endowed with that dulness of comprehension which has induced your correspondent Clio to suppose that wilful misrepresentation of a case, accusation of want of veracity of its reporter, and laughing satire, was the proper way to ask for "some further illustration." Perhaps, when your correspondent has proceeded further in his studies, he will find that it is possible for the dimensions of the young female pelvis to be increased during parturition; but more particularly he will become acquainted with the accommodation of the heads of children to the pelvis; and he will, also, perhaps be led to take into consideration the difference of the length of the child's neck previous to respiration and after that function is fully established,

by which the neck is very much shortened, from the expansion and rising of the chest and shoulders.

On referring to the case it will be found that the neck only of the first presenting child could be in the vagina, and not the chest and shoulders, the arms having been brought down; and the forceps were applied without difficulty, laterally of course, and not in the antero-posterior direction, as your correspondent supposes from his calculations; and further, it was stated that the children were small.

Having given the case in sufficient detail for all practical purposes, I shall not further enter into a discussion with one under a fictitious title.

I have the honour to be, Gentlemen,

Yours, very respectfully,

JOHN FLETCHER.

Shifnal, Shropshire, August 23, 1843.

* * Did Clio ever hear of the Siamese twins?—or of a case quoted by Dr. Rigby, of two children united by the anterior surfaces of the thorax, whose diameter was between seven and eight inches?—ED.

ANOMALOUS VACCINE ERUPTION.

GENTLEMEN,—I beg to forward the enclosed communication, which I shall feel obliged by your inserting in the Provincial Medical Journal.

I remain, Gentlemen,

Your obedient servant,

CLEMENT JAMES HAWKINS.

Surgeon to the Cheltenham Dispensary.

Cheltenham, August 22.

In the year 1832, during the time I was a pupil at the Dispensary and Casualty Hospital in this town, I vaccinated an apparently healthy child, about five years of age, residing in a district in which small-pox had proved very fatal; two of the inmates of the house in which the child lived had recently died of the disease. The vaccination was successful, and on the seventh day the late Mr. Minster, surgeon of this town, vaccinated the only child of a lady of rank from the arm of this child, and I also vaccinated an infant from the same source. On the evening of the same day I was requested to visit the child from whom he had taken the lymph, the mother alleging that the child had fallen ill after "he had lanced" the arm. I went immediately, and found the child hot and feverish; on examining the body, to my great dismay, I found an eruption appearing on the face and neck, with which I was too familiar to be deceived; it was small-pox. I prescribed a mild aperient, and went home. On the following morning I communicated the intelligence to Mr. Minster, who wisely said we must hold our tongues, and wait the result, which I did with great anxiety. Mr. Minster's patient took the vaccine disease, which went through its usual course. The other child did not receive the disease, which I attributed to a general disorder of the constitution which it was laboring under at the time of the operation. The patient from whom the vaccine lymph was taken had the small-pox mildly, and recovered favorably; the vaccine vesicles were retarded and matured with the small-pox.

The case was witnessed by the late Messrs. Minster and Whitmore, and other medical practitioners.