opinion, if not erroneous, at least very defective. He ascribes it solely to the cough, and observes, "The violent and full expirations occasioned by coughing are, of course, immediately followed by inspirations correspondingly ample, by which the air is introduced in large quantities into the lungs. When ten times the amount of air, not only in individual acts, but in a given time, enters the lungs, these organs are expanded in a correspondingly increased proportion. The air tubes and cells are under the necessity of doing ten times the amount of functional duty; and it is a law of the animal system, that the development or enlargement of an organic apparatus is according to the exercise of it. We know of no exception, nor can we imagine one." This is certainly a law of universal operation, but it must have a sound and proper application. It is one thing for organic structure to develop from increased action, but an increment of space is another thing. This law refers *only* to structure, and in its operation only to the development of structure. It is true that the bronchial tube does dilate from increased action, but not from increased action of the tube; indeed, the air tube becomes more than usually a passive agent. The air tubes are dilated from the increased quantity of air that is brought into the lungs, also into particular parts of the lungs. But before even this can operate, other conditions are required. There are two classes of tissues in the structure of the air tubes—the muscular and non-muscular. It is the inflammation of these that is the primary cause of the dilatation. In the case of the non-muscular, from continued irritation, there is a loss of elasticity, and the tube is exposed to the distending influence of the air; and in the instance of the muscular there is a loss of tonicity of the part, followed by innervation and paralysis similar to that of the intestine in ileus. We thus can account for the difficulty of expectoration, the contractile power being gone, also the dyspnoea—one very important adjunct in expiration being lost.

What, then, is this disease?—is it phthisis? Dr. Holland treats of the disease under the name of phthisis; but the cases he enumerates, and the pathological results which he details, lead to no such conclusion. If we are to understand by phthisis the literal meaning of the word—wasting away or decay—the term may not be objected to; but if we are to understand by that term tubercular disease, the question may be answered in a decided negative. No doubt grinders' disease may be conjoined with phthisis, and in a strumous habit the irritation in the bronchial membrane from the dust will easily call into play the morbid principles of that peculiar diathesis. In a case which terminated fatally the other day (August 21), and which we had hoped to have introduced on the present occasion, but that an examination was denied, there was every symptom of tubercular disease, but there was a prominently strumous habit. In another case, chronic laryngitis with tubercular disease was present, and where the patient had worked forty years as a penblade-grinder—one of the worst branches of the trade. But such cases are the exception rather than the rule.

The consistence of many of the morbid deposits resembles that of tubercle, but the proof that it is not tubercle will be found in the absence of all such con-

stitutional symptoms as are observed in the history of phthisis. There is a gradual decay of the system, but no irritability, no fever, or sweats, or any symptom indicating extensive sympathetic action. Where there is strumous diathesis, we have observed that the disease has commenced earlier in life, and been more rapid in its progress. Indeed, it is natural to suppose that the disease will take its aspect from the nature of the constitution; and that on such will depend, as in the case of the operation of other external agents in causing disease of the lungs, whether tubercle or disease of the air tubes is to be the exponent of morbid action.

Another and very interesting question that presents itself, with regard to these morbid deposits, is as to the mode of their formation. Are they simply inhaled, and there remain, or are they deposited from the blood? The late Mr. Thackrah, in his observations on a statement made by Diemerbroeck, that he found heaps of sand in the lungs of masons after death, inclines to the supposition that they were calcareous deposits, and were not inhaled. But tracing, as we can, by a decided connection, the daily inhalation of a foreign body into the lungs during life, and the existence of a similar body after death, surely it is more easy to suppose that the substance found after death is the same that was introduced during life, especially when the contrary supposition is attended with so many difficulties. The same view will apply to the question under consideration. In the daily occupation of grinders, there is the expectoration of black, gritty sputa, composed of the metallic and other particles inhaled. If the whole is not expelled, it must either remain in the state in which it entered, or must be dissolved and absorbed into the blood. The latter we cannot conceive. Nevertheless, it must be allowed that we are placed in a dilemma with regard to the bronchial glands. The gritty particles could not enter them by inhalation, and must have been deposited. The subject is evidently beset with some difficulties, and will require further study and opportunities of investigation.

We may state, in conclusion, that we have at present ten or twelve cases of this interesting disease under treatment; and although there is much difficulty in obtaining post-mortem examinations, we hope to be able to lay several of the cases at a future time before the readers of this Journal.

CASES OF DIABETES.

By CHARLES F. FAVELL, M.D.,

Physician to the Sheffield General Infirmary, &c.

(Continued from page 479.)

CASE II.—William Parke, aged fifteen, was admitted into the infirmary December 5th, 1842. He has been employed, for the last three or four years, as a farmer's servant; has worked a good deal in the fields, and consequently been exposed to the inclemencies of the weather. Until about four months ago he always enjoyed good health, and was able to get through his daily work with ease and comfort. His health then began to fail, but he is unable to give any accurate account of the commencement of his illness. His appetite for liquids and solids became

very great, and he was sent into the infirmary because his parents were unable to satisfy his craving for food. Both his parents are living, and in good health. At present there is considerable emaciation, which has been gradually increasing since the commencement of his complaints. His size is below the average of youths of his age. Countenance not expressive of suffering, slightly florid, dull, and unintellectual; hair thin, poor, and very light colored; scalp tender, and in one or two places slightly abraded; complains of great weakness and languor; skin dry, never perspires; pulse 80, of tolerable strength; bowels regular; no pain either in stomach, bowels, or back, but has sometimes a sense of oppression at the stomach after eating; tongue red and dry; thirst very great; appetite voracious; passes twenty pints of urine in twenty-four hours, of a pale straw color, and very sweet taste, specific gravity 1.040; slight cough, unaccompanied by pain or expectoration; lies with equal facility on either side; on percussion, slight dullness is perceived along and immediately below the right clavicle; inspiration in the same situation feeble, but no abnormal loudness on expiration.

He was put on a meat diet, and was not allowed any vegetables, bread, or pudding; dry biscuit was to be taken with the meat. The following medicine also prescribed:—

Tincture of opium, one drachm;
Tincture of sesquichloride of iron, three drachms;
Sulphate of quinine, eight grains;
Water, eight ounces. To take an ounce three times a-day, and four grains of mercury with chalk.

After continuing this treatment for six days, the quantity of urine was increased to twenty-two pints in twenty-four hours, but the specific gravity had slightly diminished. The following alterations were then made in the treatment:—He was to have meat as before, but no biscuit; to drink weak brandy and water; to have brocoli with his meat twice a-day, and an egg night and morning; to omit the medicine previously prescribed, and take the following:—

Infusion of Calumba, eight ounces;
Sesquicarbonate of ammonia, two scruples. An ounce three times a-day, and use a warm bath every night.

It was on the 12th of December when this plan was adopted, and on the 14th the quantity of urine had diminished to twelve pints. On the following day the quantity of urine passed was nine pints and a half. The thirst was greatly diminished; he drank five pints in twenty-four hours, and perspired freely during the night; specific gravity of urine 1.035. From the 15th to the 23rd the quantity of urine varied from nine pints to fourteen, the average being about twelve pints. On the 23rd it amounted to thirteen pints, specific gravity 1.040; quantity of solid extract in a pint, two drachms; quantity of fluid drunk in twenty-four hours, six pints. A scruple of carbonate of ammonia and a drachm of tincture of opium were added to the mixture. On the 31st the quantity of urine passed was ten pints and a half; specific gravity 1.040; a pint yielded three drachms of solid extract. During the intervening period there had once been only seven pints passed, and on one day it amounted to fifteen pints and a half. On the 1st of January he

passed nine pints of urine; on the 3rd, ten pints. He then complained of considerable pain at the stomach, which was increased by pressure; no sickness; thirst rather diminished. He was ordered to omit the brandy, to take two minims of hydrocyanic acid in water three times a-day, and to have a dozen leeches applied to the epigastrium. By these means the pain was speedily relieved, and he went on without much variation till the 10th, when he had an attack of fever, with a pulse at 120; but, although there was considerable febrile excitement, the thirst was not increased; the quantity of urine was eight pints. The prussic acid was omitted, and a saline aperient mixture ordered. The feverish attack did not last long, and on the 19th he again returned to the infusion of Calumba. He was then passing ten pints of urine, specific gravity 1.038; and the appetite for solid food was decidedly diminished. From the 19th to the 28th the urine ranged from ten to eleven pints, and he was then ordered to take one ounce of the compound iron mixture three times a-day. On the 31st he began to take three grains of antimonial powder, with ten grains of Dover's powder, every night. On the 7th of February the quantity of urine had again risen to twelve pints, and the thirst was so much increased that he drank eight pints of water in the twenty-four hours. He was ordered to omit all the other remedies, and to take eight minims of the hydrosulphuret of ammonia three times a-day. From this period the quantity of urine gradually diminished, until it vibrated between five pints and five and a half. In the meantime the dose of the hydrosulphuret was gradually increased, till it reached fifteen drops three times a-day. The appearance of the patient became very much improved. He was lively and active, and was so much stouter and more healthy in appearance, that his friends scarcely knew him. The skin acted freely; the thirst was moderate, and the appetite very much diminished; in fact, he was no longer able to eat the very plentiful allowance on which he had been placed. Under these circumstances, his friends were anxious that he should return home, and he was, therefore, discharged from the infirmary March 30, very considerably improved.

I heard nothing of Parke after his discharge for about a month or six weeks, when a young man from the same neighbourhood (a considerable distance from Sheffield) was admitted into the infirmary, and informed me that my poor patient was dead. He was unable, however, to give me any particulars respecting his death, and I, therefore, wrote to a gentleman who resided near Parke, and took great interest in his case; but the information I received from him is by no means so precise as I desired. After stating how greatly improved in health and strength the poor boy was when he returned from the infirmary, my correspondent observes, "but this improvement did not continue for more than a week, when thirst returned, which he quenched by milk, water, or what he could get, and the secretion of urine also increased in proportion. In this state he continued for nearly a week, when he lost all appetite for solid food, and complained of great weakness and debility. About five days before he died he drank a great quantity of newly-churned milk, from which

time he ate scarcely any solid food. A day or two after this time his mind was for a short period affected, and he became incoherent in conversation. A dulness or heavy stupor never seemed to leave him." His parents, however, did not anticipate his removal, as they had previously seen him very similarly affected. It would have been exceedingly interesting to have made a post-mortem examination of the body, but this was impossible.

REMARKS.

We should scarcely be justified, with the very limited amount of information we possess on the subject, to enter into any speculations respecting the immediate cause of death in the foregoing case; but there are one or two points connected with the treatment on which it may be proper to make a few very brief observations.

In the first place, the administration of the tincture of the sesquichloride (muriate) of iron, sulphate of quinine, and tincture of opium, was decided upon in consequence of the cases published by Mr. Clay, of Manchester, in which very considerable amendment appeared to result from this treatment. In my case the patient certainly derived no benefit; on the contrary, the quantity of urine was somewhat increased. It may certainly be said that the trial given to the medicine was not of sufficient duration. But to this it may be replied, that there was very little encouragement to perseverance, inasmuch as the symptoms gradually grew worse.

Secondly, When the treatment just noticed was discontinued, that suggested by Dr. Barlow was adopted, and apparently with decided advantage. When Parke commenced eating the cruciferous vegetables and taking ammonia, he was passing twenty-two pints of urine in the twenty-four hours. In two days afterwards the quantity fell to twelve pints (a diminution of ten pints), and on the third it fell to nine pints and a half. It is true that subsequently he passed a larger quantity than that last mentioned on several occasions, but it never rose to nearly the same amount as it had attained before the change of treatment was adopted.

Thirdly, The compound iron mixture of the Pharmacopœia was substituted for the vegetable tonic and ammonia, in consequence of the symptoms being, as nearly as possible, stationary, whilst the quantity of urine still continued to be very much above the healthy standard. The change appeared to be, to some extent, beneficial, inasmuch as there was a slight diminution in the secretion of urine.

Fourthly, The hydrosulphuret of ammonia was finally substituted for Griffith's mixture, in order to see what influence a medicine which has a powerful sedative action on the heart would exert over the secreting power of the kidneys. The details of the case furnish an answer to the inquiry. When he commenced taking the medicine the quantity of urine had again increased to twelve pints, and had for ten days been oscillating between eight pints and a half and ten pints and a half, but it afterwards rapidly diminished. He continued taking the medicine for very nearly two months, during the whole of which period the quantity was only, on one single occasion, ten pints and a half, and at the termination of it did not exceed five pints and a half. For some days

before leaving the house it remained steadily at five pints.

Fifthly, I may observe that, so long as the secretion of urine continued so decidedly above the healthy standard, although the general health of the patient was vastly improved, and the quantity of urine had diminished fifteen pints in the twenty-four hours, I should not have discharged him had it not been for the earnest and persevering entreaties of himself and friends.

CASE III.—On Sunday evening, August 13, 1843, I was requested to visit Mr. W. C., who had long been under the very judicious care of an able medical friend, and had also consulted Dr. Prout, on account of suffering from diabetes. It is not necessary that I should enter into the details of the case, especially as I did not see it till the period just mentioned. It will, however, be proper to remark that he had been subjected to the treatment recommended by Dr. Barlow, which for a time appeared to exercise considerable control over the secretion of the kidneys. The quantity of urine, however, remained steadily above the healthy standard, and the specific gravity several degrees higher than natural. When I visited him, I found him sitting up in his bed-room, considerably emaciated, and with all the usual symptoms of diabetes. He was cheerful, and wished to know my opinion respecting a temporary removal to the sea coast. On the previous day he had been out attending to some matters of business, but had passed a restless night, in consequence of severe pain in the back, which was the immediate cause of my attendance being requested. He represented the pain as not being stationary, but moving up and down, and being particularly severe on moving the body either backward, forward, or laterally, and also on coughing, but not increased by pressure over the spine. He had for several weeks been troubled with a short cough, attended with but little expectoration, but it had been worse for two or three days. The feet and ankles were considerably œdematous. The quantity of urine was about the same as it had been for several weeks, averaging about six pints, or rather more, in the twenty-four hours. It had a pale color, and a faintish, disagreeable smell. It reddened litmus, was not affected by heat, and had a specific gravity of 1.035. The pulse was moderate in strength and frequency, and there was no heat of skin. I was of opinion that the pain in the back was of a rheumatic character, and consequently recommended that he should be well fomented with a decoction of chamomile and poppy-heads, and take two grains of opium at bed-time. On visiting him next morning, I found that he had derived no benefit from the means employed. He had passed a very restless night from the severity of the pain, which was now more fixed over the region of the right kidney. In all other respects he was as before, the secretion of urine not being in the slightest degree diminished. I recommended the application of six leeches over the seat of pain, a saline camphor mixture to be taken every four hours, and some antimony and Dover's powder at bed-time.

In the evening he was considerably relieved. On the following day I did not see him, but on the Wednesday morning I was requested to pay him an early visit. When I arrived at his house, between nine

and ten o'clock, I learnt that during the whole of Tuesday he had been very comfortable, but became somewhat uneasy towards the evening. He passed a very restless night, and when I saw him was unable to speak, but he had a violent convulsive struggle, and then fell into a state of coma. I saw him again in the evening, but he was evidently sinking rapidly. The pulse was almost imperceptible, the breathing stertorous, insensibility perfect. During the whole day the secretion of urine continued in such quantity, that it ran through the bed. It was impossible to ascertain the quantity passed, but it must have been very considerable, as it completely saturated the bed and bed-clothes, and mattress, and the thicknesses of linen which were repeatedly applied to him. He died during the night.

Now, I have ventured to call this case interesting, "from the mode of death," and that it is so, I think will be generally admitted. This mode of death is unusual in diabetes. It does not frequently happen in such cases that patients are cut off suddenly, whilst the general health is not so much impaired as to render them unable to attend to their usual avocations. But such was the fact here; Mr. C. was out on business till rather a late hour on Saturday evening, and he would have been out again on Monday (two days before his death) had it not been for my interdiction.

It is interesting to inquire whether the severe pain in the back, of which the patient complained, had any connection with the cause of death. When first I saw him, it has already been stated, I considered the pain to be of a rheumatic character, and attributable to cold. But the result of the case has materially altered my opinion, and I now believe that the lesion which caused death also gave rise to the anomalous pains under which, for two days, the patient so severely suffered. I take it for granted, from the symptoms which preceded death (unfortunately I had not the opportunity of making a post-mortem examination), that there was effusion about the base of the brain. Now, it may be regarded as an established law in pathology, that a cause of irritation seated in a nervous centre often produces its first and most painful effects on the peripheral extremities of the nerves proceeding from the part. Hence we find that distant organs frequently give the first indication of the most formidable and fatal diseases. Great irritability of the stomach, for example, is very often the precursor of the most serious affections of the brain. I have known such irritability continue for a week or more uncontrolled by any remedies, and at the end of that time a sudden convulsion has carried off the child, or a fit of apoplexy, if the patient were an adult, has put a speedy termination to his sufferings. On other occasions I have known the cerebral seizure follow close upon the sickness; but the irritation is sometimes reflected to organs still more remote, and then the character of the anomalous complaints, under which patients suffer, can only with certainty be known, after some formidable but unequivocal indication has revealed the condition of the central organ. Experience has amply confirmed the truth of these observations to every one who has been long engaged in medical practice. But I must not enlarge; I only add, that the view I am disposed to take of the case just recorded is, that

a cause of irritation (perhaps capillary congestion) existed about the base of the brain or commencement of the medulla oblongata, and occasioned by reflexion an abnormal condition, first, of the posterior, and, secondarily, of the anterior nerves proceeding from some of the ganglia of the spinal cord, and that the same cause led to sanguineous effusion, and ultimately issued in coma and death.

In conclusion, I would suggest that the following four inferences may be fairly deduced from the cases now detailed.

1. That there is no necessity to impose upon patients laboring under diabetes an abstinence from all vegetable substances.
2. That the improvement effected in diabetic patients is often extremely transitory. A discontinuance of remedial agents, or a departure from diabetic regulations, too frequently occasions a speedy aggravation of the symptoms.
3. That the sudden invasion of severe pain, whatever may be its precise locality, in advanced cases of diabetes, is calculated to occasion great anxiety in the mind of the medical attendant.
4. That during the course of diabetes, serious disease may suddenly affect a vital organ, and prove fatal, without the abnormal quantity of urine being diminished.

Sheffield, September 5, 1843.

CONTRIBUTIONS

TO THE

PATHOLOGY OF CHILDREN.

By P. HENNIS GREEN, M.B.

HÆMOPTYSIS IN PHTHISIS.

It is a well known fact, though one not easily explained, that hæmoptysis and pulmonary hæmorrhage very rarely accompany any of the forms of phthisis to which children are extremely subject. Sir James Clark, in his work on Consumption (p. 61), observes, "Hæmoptysis is a rare occurrence; at least, I do not recollect to have met with any case in which it was present."

In a series of 118 cases of pulmonary tubercle in children, observed consecutively and in the same year, with the object of forming a ground-work for the history of phthisis in children, I saw only five examples of hæmoptysis,* although a few other cases presented themselves, in which the parents or patients alleged that spitting of blood had occurred at some period previous to the child's admission into hospital. This fact demonstrates the truth of Sir James Clark's remark: hæmoptysis is a rare symptom in the pulmonary consumption of children, and hence the following examples may prove of some interest to the profession. Before entering on a detail of the cases, I may, however, observe that the absence of hæmoptysis may in part be accounted for by the circumstance that children (especially those of tender years) do not expectorate, but swallow, everything which may be brought up into the mouth from the pulmonary mu-

* In 70 other cases observed, not consecutively, but at different periods, hæmoptysis did not occur once, at least while the patients were under observation.