and ten o'clock, I learnt that during the whole of Tuesday he had been very comfortable, but became somewhat uneasy towards the evening. He passed a very restless night, and when I saw him was unable to speak, but he had a violent convulsive struggle, and then fell into a state of coma. I saw him again in the evening, but he was evidently sinking rapidly. The pulse was almost imperceptible, the breathing stertorous, insensibility perfect. During the whole day the secretion of urine continued in such quantity, that it ran through the bed. It was impossible to ascertain the quantity passed, but it must have been very considerable, as it completely saturated the bed and bed-clothes, and mattress, and the thicknesses of linen which were repeatedly applied to him. He died during the night.

Now, I have ventured to call this case interesting, "from the mode of death," and that it is so, I think will be generally admitted. This mode of death is unusual in diabetes. It does not frequently happen in such cases that patients are cut off suddenly, whilst the general health is not so much impaired as to render them unable to attend to their usual avocations. But such was the fact here; Mr. C. was out on business till rather a late hour on Saturday evening, and he would have been out again on Monday (two days before his death) had it not been for my interdiction.

It is interesting to inquire whether the severe pain in the back, of which the patient complained, had any connection with the cause of death. When first I saw him, it has already been stated, I considered the pain to be of a rheumatic character, and attributable to cold. But the result of the case has materially altered my opinion, and I now believe that the lesion which caused death also gave rise to the anomalous pains under which, for two days, the patient so severely suffered. I take it for granted, from the symptoms which preceded death (unfortunately I had not the opportunity of making a post-mortem examination), that there was effusion about the base of the brain. Now, it may be regarded as an established law in pathology, that a cause of irritation seated in a nervous centre often produces its first and most painful effects on the peripheral extremities of the nerves proceeding from the part. Hence we find that distant organs frequently give the first indication of the most formidable and fatal diseases. Great irritability of the stomach, for example, is very often the precursor of the most serious affections of the brain. I have known such irritability continue for a week or more uncontrolled by any remedies, and at the end of that time a sudden convulsion has carried off the child, or a fit of apoplexy, if the patient were an adult, has put a speedy termination to his sufferings. On other occasions I have known the cerebral seizure follow close upon the sickness; but the irritation is sometimes reflected to organs still more remote, and then the character of the anomalous complaints, under which patients suffer, can only with certainty be known, after some formidable but unequivocal indication has revealed the condition of the central organ. Experience has amply confirmed the truth of these observations to every one who has been long engaged in medical practice. But I must not enlarge; I only add, that the view I am disposed to take of the case just recorded is, that

a cause of irritation (perhaps capillary congestion) existed about the base of the brain or commencement of the medulla oblongata, and occasioned by reflexion an abnormal condition, first, of the posterior, and, secondarily, of the anterior nerves proceeding from some of the ganglia of the spinal cord, and that the same cause led to sanguineous effusion, and ultimately issued in coma and death.

In conclusion, I would suggest that the following four inferences may be fairly deduced from the cases now detailed.

1. That there is no necessity to impose upon patients laboring under diabetes an abstinence from all vegetable substances.
2. That the improvement effected in diabetic patients is often extremely transitory. A discontinuance of remedial agents, or a departure from diabetic regulations, too frequently occasions a speedy aggravation of the symptoms.
3. That the sudden invasion of severe pain, whatever may be its precise locality, in advanced cases of diabetes, is calculated to occasion great anxiety in the mind of the medical attendant.
4. That during the course of diabetes, serious disease may suddenly affect a vital organ, and prove fatal, without the abnormal quantity of urine being diminished.

Sheffield, September 5, 1843.

CONTRIBUTIONS

TO THE

PATHOLOGY OF CHILDREN.

By P. HENNIS GREEN, M.B.

HÆMOPTYSIS IN PHTHISIS.

It is a well known fact, though one not easily explained, that hæmoptysis and pulmonary hæmorrhage very rarely accompany any of the forms of phthisis to which children are extremely subject. Sir James Clark, in his work on Consumption (p. 61), observes, "Hæmoptysis is a rare occurrence; at least, I do not recollect to have met with any case in which it was present."

In a series of 118 cases of pulmonary tubercle in children, observed consecutively and in the same year, with the object of forming a ground-work for the history of phthisis in children, I saw only five examples of hæmoptysis,* although a few other cases presented themselves, in which the parents or patients alleged that spitting of blood had occurred at some period previous to the child's admission into hospital. This fact demonstrates the truth of Sir James Clark's remark: hæmoptysis is a rare symptom in the pulmonary consumption of children, and hence the following examples may prove of some interest to the profession. Before entering on a detail of the cases, I may, however, observe that the absence of hæmoptysis may in part be accounted for by the circumstance that children (especially those of tender years) do not expectorate, but swallow, everything which may be brought up into the mouth from the pulmonary mu-

* In 70 other cases observed, not consecutively, but at different periods, hæmoptysis did not occur once, at least while the patients were under observation.

cous membrane. Mr. Murdoch relates a striking example of this fact in his thesis. A child, two years old, died suddenly from the rupture of a blood-vessel which traversed a tubercular cavity in the right lung. A very small quantity of blood had been discharged through the mouth, for the child had swallowed the greater part of it, and on examination after death the stomach and upper portion of the small intestines were found full of enormous clots of blood, which were moulded over their internal surfaces.

CASE I.—*Death from Hæmoptysis*.—Caroline L., nine years of age, was admitted into the Children's Hospital, Paris, on the 25th February, 1834. The parents of the child are healthy. According to the mother's account the cervical glands have become enlarged within the last six weeks only. The child has been feeble and delicate for a considerable time, and has gradually lost flesh during the last six months; since then, also, she frequently complained of pain in the right side of the chest, which was observed to become more bulging than the left side. About two months back she expectorated a quantity of purulent matter, with increase of pain and dyspnoea, and since that period the enlargement of the chest has gradually subsided. She has slight febrile paroxysms from time to time, but no regular hectic, and suffers from constant diarrhoea.

Feb. 26. On examining the chest this morning the following physical signs were noted:—There is dulness on percussion over the whole of the right side of the chest, in front, particularly at the upper part; some dulness, also, behind; the respiratory murmur is excessively feeble in those points of the chest, but there is no souffle, gargouillement, or pectoriloquy. The child coughs frequently, and expectorates a quantity of foetid, purulent matter, but no round masses in the sputa; pulse 120; respiration 40; moderate thirst; some appetite; has passed three fluid stools. She was ordered a tonic mixture with sulphate of quinine. On March 1st the same stethoscopic signs were noted.

5. The expectorated matter more abundant and firm; diarrhoea increased. Quinine omitted.

12. The sound at the upper part of the chest on the right side is more dull; behind, near the middle of the scapula, is heard a cavernous souffle with pectoriloquy. On the 18th the child was suddenly carried off by violent hæmorrhage from the lungs, the blood issuing in quantity through the mouth and nose.

On examining the body on the following day, the whole of the right lung was found to have been transformed into a substance of a dirty white color, traversed by white bands; it had altogether the appearance and consistence of firm cheese. The lung adhered extensively to the walls of the chest, and was extremely friable. In the centre of the caseous mass, just alluded to, was a large anfractuous cavity, partly filled by coagulated blood. The bronchus distributed to this portion of the lung and the trachea likewise contained clotted blood. The left lung contained a few tubercles, some of which were beginning to soften. The tubercular infiltration of the right lung was interspersed with lenticular tubercles, from three to four lines in diameter; they were continued from the lung into the intercostal muscles and diaphragm. The abdominal viscera were healthy, with the exception

of the lower portion of the ileum, which contained numerous ulcerations.

CASE II.—*Hæmoptysis*.—A. Foucault, fourteen years of age, a cotton carder, was admitted into hospital on the 14th December, 1833. The father and mother of the child are healthy, and he has six brothers and sisters also healthy. About five months back, at Lille, he was attacked by ague, which lasted for three months, and since then he has been much subject to fatigue and exposure to the weather while walking. Four days back, without any apparent cause, he was seized with hæmoptysis, and according to his own account spit up about an ounce of blood within half an hour. The spitting of blood has persisted since then, but in smaller quantity. Immediately on admission eight ounces of blood were abstracted from the arm.

15. During the night the child spit up one or two sputa of pure, grumous blood, and several others of blood mixed with mucus; each expectoration was preceded by a rattling sensation in the thorax and a pricking in the larynx; skin hot; pulse 112; respiration 28; the respiratory sound, posteriorly, is pure, and without any râle; in front and beneath the right clavicle there is some resonance of the voice; tongue moist; no pain in abdomen; no diarrhoea. He was ordered to take the decoction of ratanhia and to have sinapisms to the legs. There was no recurrence of the hæmoptysis during the day; cough moderate.

16. Only a few striae of blood in the sputa; pulse 96; respiration 24; skin cool. Six ounces of blood were drawn from the arm.

17. No trace of inflammatory crust on the blood; the sputa are still tinged, but there is no fever, oppression of breathing, or pain in the chest. No diarrhoea.

On the 18th and 19th the sputa continued to be slightly tinged with blood. Four leeches were applied to the feet.

On the 20th, 21st, and 22nd there was no trace of blood in the sputa.

23. The cough to-day is somewhat worse; the boy complained of a sense of tickling in the throat, and soon afterwards spit up a small quantity of pure blood; pulse natural, 64; respiration not accelerated; skin moderately warm; appetite good; no perspirations at night. To continue the decoction of ratanhia.

On the 25th there was a tinge of blood in the sputa, but this disappeared on the 26th, and the child soon afterwards left the hospital, apparently well.

CASE III.—*Death from Hæmoptysis*.—Lucrece Rey, a girl, eleven years of age, was admitted into hospital on the 12th of May, 1835. This child had been subject to cough for a few days only, when, on the 9th of May, she was suddenly attacked by severe shivering, which compelled her to return home from school; during the course of the same night violent pain in the side of the chest supervened, with fever and oppression of the respiration; and these symptoms continued up to the day of her admission. No medical treatment had been adopted.

13. The little patient presents the following symptoms when examined this morning:—The face somewhat flushed; cough without expectoration; dyspnoea; pain under the left mamma, increased on coughing and by the movements of inspiration; the respiratory

sound is feeble on the left side of the chest, with dullness on percussion, but there is no œgophony; some sibilant râle at the summit of the left lung; skin hot; pulse 116; respiration 42; tongue covered with a white fur; loss of appetite; no pain of abdomen; has passed only one stool for the last three days. Eight ounces of blood were abstracted from the arm, and twelve leeches applied to the painful side of the chest.

This treatment had the desired effect. The pain in the side disappeared in a few days, and the child was able to walk about the wards, when she was again suddenly attacked by pain in the left side, with considerable dyspnœa, the presence of effused fluid being indicated by œgophony. She was again relieved by antiphlogistic treatment, and was sitting up in bed on the morning of the 13th of June, when violent hæmoptysis came on, and terminated life in a few minutes, the blood issuing from the mouth and nose in great quantities.

Body Examined Twenty-six Hours after Death.

The right lung is free from adhesions, and the pleural cavity from effusion; in its tissue are several crude tubercles, with numerous grey, semi-transparent granulations, and some lenticular tubercles underneath the pleura costalis; on dividing the tissue of the lung, there appears a number of round, dark red spots, completely isolated from each other, and produced by effused blood; the tissue surrounding these spots is healthy.

On the left side the pleura is thickened, and the opposite surfaces extensively adherent by false membranes, which contain some tubercles; the cavity of the pleura on this side contains about two ounces of bloody serum. In the interior of the lung are numerous spots of effused blood, of the size of a sixpence, similar to those in the right lung. The tubercles are more numerous in this (the left) lung, and chiefly seated in the superior lobe. There is a good deal of coagulated blood in the larynx, trachea, and bronchi, and their lining membrane is of a deep red color, from imbibition. The bronchial glands are tuberculated. The heart is healthy, but pale and flaccid; underneath the serous membrane which covers the muscular substance of the heart there is a small lenticular tubercle, and seven or eight miliary tubercles near its base.

There is a small clot of blood in the œsophagus, and a second, as large as a hen's egg, in the stomach; some submucous tubercles, with ulcerations, near the termination of the ileum; large intestines healthy. The liver is united to the diaphragm by false membranes, containing tubercles, some of which exist in the substance of the liver, also in the spleen and mesenteric glands. The brain and its membranes are healthy; but near the middle of the left hemisphere and in the cortical substance is a small tubercle of the size of a pea, the surrounding nervous tissue being perfectly healthy.

CASE IV.—Hæmoptysis without Tubercle.—In connection with the preceding cases I may relate the following very rare example of hæmoptysis independent of tubercular deposit in the lungs:—

Clementine Scallier, a girl, thirteen years of age, was admitted into hospital on the 18th of June, 1834. The patient had been dismissed six days previously, convalescent from scarlatina. On the 17th, without

any apparent cause, she was attacked by shivering, with some febrile symptoms, and pain under right mamma, followed by expectoration of blood. On the 18th the symptoms were aggravated; the hæmoptysis more frequent and abundant, pure grumous blood, of a bright red color, being expectorated. Immediately on admission into hospital she was bled from the arm, with considerable relief, and passed a quiet night; the pain in the side was much less severe; no cough; expectoration of blood diminished.

19. The skin is desquamating extensively in small scales; face swollen, and eyelids slightly œdematous; skin moderately hot; pulse 132; respiration 48; dyspnœa; tongue foul, and somewhat dry; slight thirst; no nausea or vomiting; has passed several fluid stools; expectorated this morning some sputa of pure blood; the respiratory murmur is more feeble on the right than on the left side; no œgophony or dullness on percussion; the whole of the right side of the chest is painful. To be bled to twelve ounces, and sinapisms placed on the legs; milk diet.

20. Face more swollen than yesterday; pulse 121, regular, but very compressible; respiration anxious, jerking, with dilatation of nostrils, 56; has spit up a few clots of blood; there is some trace of *souffle* at the summit of the right lung, with bronchophony; cough, attended by pain along the course of the sternum; skin very hot; has passed three fluid stools; tongue dry and foul; great thirst. To take fifteen grains of the oxide of antimony in gum mixture.

21. Pulse 128; respiration 60; voice faltering; pain under the edge of the false ribs increased by pressure; skin very hot and dry; child agitated, and some *subsultus tendinum*; has vomited once, and passed three fluid stools; abdomen tender to the touch; has expectorated several clots of dark blood; *souffle* tubaire and bronchophony under the right scapula; the sputa frequently tinged with blood; abdomen tympanitic; no sudamina or lenticular spots. Continue remedies.

22. Pulse 120; respiration 40, less difficult; skin still hot; the diarrhœa continues, five or six fluid stools being passed during the day; abdomen painful on pressure, and tympanitic; has expectorated several sputa of pure blood; the pain in the side continues; to-day and yesterday a few indistinct papulæ were noticed on the face and arms.

23. Pulse 120; respiration 48; pain in chest persists; sputa slightly rusty; abdomen very painful; has passed several fluid stools; sound of chest, on percussion, equal on both sides; some sub-crepitant râle on the right side, the respiratory murmur pure in other parts; the eruption is now evidently small-pox, but it has made no progress; the vesicles are pale, and there is no trace of inflammatory action around them. To have a gum mixture, cataplasms to the abdomen, and the white oxide of antimony, as before.

25. The pustules are now more developed on the face, but still pale, and without areolæ; on the limbs they are irregular, imperfect pustules being mixed with papulæ and vesicles; the cough remains unchanged; expectoration of viscid, transparent sputa, without any blood; pain continues in the right side, where a crepitant râle is now distinguished, with feeble respiratory murmur, especially at the summit of the lung;

tongue cleaner; thirst abated; voice hoarse, and child complains of some pain in the throat, with difficulty of deglutition; three or four fluid stools; vomited once yesterday after taking her mixture; pulse firm, 128; respiration less anxious, 48.

26. The pustules are now more fully developed on the abdomen, which is less painful on pressure; cough and expectoration the same as before; pulse regular, 112; respiration 36; skin still warm, but a little moist. To have eighteen grains of the oxide of antimony.

27. Pulse 124; respiration 44; skin warm; the pustules on the face are beginning to suppurate; the *souffle tubaire* is heard in a very confined point, at the summit of the right lung, with crepitant râle, more extensively diffused; the sound on percussion is more dull on this than on the left side; the cough is unabated, but there is much less pain in the side; no expectoration; diarrhœa continues; tongue clean. Mixture, with twenty grains of the white oxide of antimony.

28. Cough and diarrhœa persist; no expectoration; child complains more of the pain in the side; pulse 120; respiration 48.

As my object in relating this case is simply to illustrate the occurrence of hæmoptysis in children, it is unnecessary to pursue the daily account of the symptoms any further. It will suffice to say that the thoracic and abdominal symptoms continued to increase to the 10th of July, when the little patient sunk under them.

Body Examined Twenty Hours after Death.

Chest.—The larynx, trachea, and bronchi, are healthy; the left lung is crepitant throughout, free from adhesions, and does not contain any trace of tubercle; the posterior and lower portion is congested. The two lower thirds of the right lung are in a state of grey hepatisation, and friable, the upper portion only being capable of admitting air; at the posterior part there is a considerable portion of the tissue, of a black color, softened, and infiltrated with serum, resembling the pulmonary tissue in an incipient stage of gangrene. The upper part of the lung is united to the walls of the chest by old false membranes, and the remaining part is also united to the costal and diaphragmatic pleuræ by false membranes of recent formation. Heart very large and soft; pericardium healthy.

Abdomen.—The stomach is large, its mucous membrane healthy; that of the small and large intestines is pale, and free from softening; near the end of the colon, however, there is some arborisation, with thinning of the mucous tissue; no tubercles in the mesenteric or bronchial glands. The kidneys are large; on dividing them there appeared several spots of a straw-yellow color, which pass from the cortical substance into the tubular in many points; these yellow spots are greasy to the touch, and give to the kidney much of the appearance peculiar to the fatty liver.

Head.—The arachnoid lining the upper part of the hemispheres is thickened and opaque; the cerebral tissue throughout healthy, but very pale; no serum effused in ventricles, or at the base of the brain.

CASE V.—*Sudden Death from Hæmoptysis.*—A. Audiffret, fourteen years of age, was admitted into

hospital on the 25th of February, 1833. The child had been previously in hospital for an inflammatory affection of the chest, and had been dismissed cured in December last. His parents are healthy, and he always lived in the country until eighteen months back, when he came to Paris. Since then he was subject to cough, with occasional pain in the side; within the last two months has had symptoms of disease of the hip-joint. He lost flesh considerably within the last six months, but never had any diarrhœa or spitting of blood; has been subject to night sweats for some time, with frequent cough and copious expectoration. On examining the patient, a few days after his admission, the following symptoms were noted:—He is excessively emaciated, but the chest is well formed; face pale; skin dry; pulse 100; respiration 32, not oppressed; cough very much abated; does not complain of any pain in the chest; expectorates a few mucous sputa; the voice is considerably changed—almost, indeed, gone—but he does not complain of any pain in the region of the larynx (the change of voice had existed for four months); the respiratory murmur is feeble on both sides of the chest, both anteriorly and posteriorly, but there is no râle or evident dullness on percussion; tongue clean; no thirst; appetite good; no pain in abdomen. During three weeks the child remained nearly in the same state, the attention of the physician being chiefly directed to the disease of the hip-joint, when, on the 27th of March, violent hæmoptysis set in, and speedily terminated existence.

Autopsy, Nineteen Hours after Death.

Chest.—The whole of the upper portion of the right lung adhered to the walls of the chest; the tissue of the lung was healthy, but it contained numerous small crude tubercles, none of which were softened; there was no trace of a cavern; the inferior lobe of this lung was congested, and the bronchi, much reddened from imbibition of blood, contained a few clots. On the left side the same old adhesions of the lung to the parietes of the chest existed, with deposit of crude tubercular matter, disseminated through different portions of the lung; the bronchi were dilated, and their lining membrane of a dark red color; the same color extended along the membrane of the trachea to the larynx, but there was no trace of ulceration in the latter organ. In the inferior lobe of this lung were numerous dark points, giving it a marbled appearance, and several small, circumscribed collections of effused blood; bronchial glands tuberculated. The heart was healthy; the pericardium contained about an ounce of clear serum.

Abdomen.—The stomach contained a good deal of blood, partly coagulated and mixed with mucus; its mucous membrane was firm, although of a deep red color in many places, but this was removed by washing. Near the end of the small intestines were two ulcerated patches, containing some *débris* of tubercular matter, and in the colon were found seven or eight small, circular, tubercular ulcers. The rest of the abdominal viscera healthy.

Head not examined.

CASE VI.—*Hæmoptysis.*—H. Landier, a boy, fourteen years of age, was admitted into hospital on the 1st of February, 1833. The parents of this child are healthy; he has been long subject to catch cold; six

months back had an attack of the chest, which lasted three or four months, with cough, pain of chest, and expectoration of blood. Within the last fifteen days the cough has become more severe; he has been frequently subject to bleeding from the nose.* On the day of admission (Feb. 1) the hæmoptysis returned with considerable violence, a large quantity of florid blood having been expectorated.

Feb. 2. Pulse 100; respiration 30; cough frequent, with expectoration, sometimes of pure blood, sometimes of blood and mucus; complains of pain behind the sternum; voice weak; the respiratory murmur clear behind, but under the right clavicle there is cavernous souffle with pectoriloquy; no gurgling; sound here dull on percussion; skin moderately warm, moist; is subject to night-sweats; no diarrhœa. To have a mixture containing the extract of ratanhy, with an opiate.

3. The expectoration of blood continues; pulse 84; respiration 24; no change in the general symptoms or physical signs. Ratanhy increased to five grains.

On the 4th the hæmoptysis was not abated; a strong decoction of ratanhy ordered. On the 5th there was no recurrence of the symptom, but on the 6th the expectoration of blood was very copious, sufficient to fill two of the spitting-pots used in the hospital; pulse 92; respiration 40, not oppressed.

7. Hæmoptysis somewhat abated; no fever; appetite good; allowed spare diet.

On the 9th there was merely a trace of blood in the sputa, and on the 12th this had disappeared, but was replaced by epistaxis. The child was removed by his parents on the 14th.

CASE VII.—*Hæmoptysis*.—Louis Klein, thirteen years of age, admitted August 28, 1833. Had a very long attack of hooping-cough when seven years old; is a child of scrofulous constitution; his mother died of phthisis when thirty-nine years old; father also died of some disease of chest. In the month of February last the child received a violent blow on the chest, and since then has complained of pain in the chest, with constant cough and expectoration striated with blood; never had any copious hæmoptysis; no diarrhœa.

On the 28th the child was again attacked by hæmoptysis; he spit up about a dessertspoonful of blood with a small quantity of mucus; the skin is rather warm; has copious night sweats; each expectoration is preceded by a sensation of gurgling in the right side of the chest; pulse 92; respiration 28; no diarrhœa or pain in the abdomen; the sound of the chest, on percussion, is very obscure on the left side, but clear on the right; there is cavernous respiration in the middle third of the left lung, with gurgling and pectoriloquy. The respiratory sound is audible in other parts of the chest.

On the 29th the hæmoptysis did not recur; pulse 88; skin cool; asks for something to eat.

Sept. 2. The child vomited two or three times, and complained much of pain in the left side of the chest; soon afterwards hæmoptysis occurred, and he expectorated a considerable quantity of blood. He was bled to four ounces.

* I have observed that children affected with phthisis are frequently subject to bleeding from the nose. The occurrence of epistaxis is nearly as frequent as that of hæmoptysis in adults. Perhaps the one hæmorrhage may replace the other.

3. The spitting of blood, which had ceased for a few hours yesterday, returned immediately after the bleeding, but the cough was much alleviated; pulse 88. Bleeding to four ounces to be repeated.

The blood drawn from the arm is very serous, and the clot not firm; pulse 104; respiration 28; no hæmoptysis; no diarrhœa. The stethoscopic signs are the same as already mentioned; night sweats very abundant. To have a mixture containing quinine and four grains of digitalis.

In this state the child continued, with little improvement, occasionally spitting blood, to the 1st of October, when he was removed from the hospital.

CASE. VIII.—*Hæmoptysis*. — Mouly, twelve years of age, had been dismissed from the hospital on the 21st of July, 1833, convalescent from a bowel attack; was readmitted on the 20th of August; while out of hospital he labored under diarrhœa, but his intellect is so dull that it is not easy to obtain any information from him. He has been subject to cough for the last five months and is much emaciated.

August 21. On examining the chest there is evident dullness of sound on percussion under the right clavicle, with cavernous râle, but no gurgling; abdomen tumid and painful; diarrhœa; no febrile symptoms. On the 3rd of September slight hæmoptysis occurred; pulse 104; night sweats not abundant. During the next few days there was no expectoration of blood, but on the 13th the child experienced a sensation of bubbling in the chest, and soon afterwards spit up a considerable quantity; pulse 84; skin hot; cough very troublesome; respiration slightly oppressed; cavernous râle and pectoriloquy heard behind the right scapula; night sweats copious; expectorates a quantity of muco-purulent matter, occasionally tinged with blood; diarrhœa continues.

During the next few days no change worthy of notice occurred; the diarrhœa had ceased after the discharge of two ascarides.

18. Face pale, and very emaciated; night sweats distressing; no pain in chest; cough frequent; expectoration of thick, yellow-green sputa; *gargouillement* and pectoriloquy evident in the left side of the chest, and dullness on percussion behind; on the right side mucous râle; tongue red; abdomen free from pain; pulse 80; respiration 24. From this period the child sunk rapidly, and died on the 4th of September.

Body Examined Twenty Hours after Death.

Brain.—The dura mater, arachnoid, and pia mater adhere at two points—on the right and left sides—to the hemispheres; it is impossible to separate them without removing a portion of the nervous tissue, but the latter is not altered in color or consistence. There is a great number of granulations in the fissures of Sylvius on both sides; about two ounces of clear serum in the ventricles; cerebral substance healthy throughout.

Chest.—The summit of the *right* lung adheres to the chest by old false membranes; the whole of the upper lobe is indurated, and contains a vast number of miliary tubercles; near the top is an anfractuous cavern, containing a dark fluid; the other two lobes are crepitant throughout, and contain a few semi-transparent granulations.

The *left* lung is free from adhesions; it crepitates

throughout the greater part, and contains a single tubercle, in the centre of which is a very small cavity. The right bronchial glands are tubercular; larynx, trachea, and bronchi, healthy. Heart healthy; about two ounces of clear serum in the pericardium.

Abdomen.—Mucous membrane of the stomach pale, very thin, and softened in many points; the first half of the small intestines contains an immense number of small, round, tubercular ulcers; the mucous lining of the cœcum is deeply injected, and also the seat of several ulcers; other viscera healthy.

As my object in relating the above cases is merely to place them on record as examples of the occurrence of hæmoptysis in children, I shall not append to them any remarks. I must, however, observe, that I have omitted to notice one form of pulmonary hæmorrhage which is peculiar to children, because I have not been able to find my notes of the only case which I had observed. This form is invariably and instantly fatal; it occurs in bronchial phthisis, and arises from the perforation of one of the large vessels of the chest by a tuberculated bronchial gland. Sudden death may thus occur from perforation of the pulmonary artery. Of this I have seen one example, and in a second case the artery was so completely flattened between two masses of the bronchial glands, that the passage of blood was interrupted, and death took place, as from disease of the heart.

London, September, 1843.

PROVINCIAL MEDICAL JOURNAL

SATURDAY, SEPTEMBER 16, 1843.

Our readers, as members of a profession, the exercise of which is a daily training in works of benevolence, cannot fail to have watched with anxiety and interest the investigations which have been pursuing of late years into the physical and moral condition of the poorer orders of the community. And, truly, these investigations have unfolded an appalling picture of wretchedness, ignorance, and profligacy, in the very heart of that country which, with honest pride, we are accustomed to look upon as the earthly abode of enlightenment and happiness.

But there was one conclusion which appeared fairly deducible from the evidence brought forward at an early period of these investigations, which must have been beyond all others painful to every Englishman. For the manufactories—those very means to which this country is indebted for so much of its wealth at home, its commerce abroad, and its precedence in the rank of civilised nations—were shown apparently to be inseparably accompanied with the seeds of social degradation, and were denounced as the sources of every evil, personal and political. The slow destruction of health—the sure abridgment of life—the procreation of a degenerate race, stunted, misshapen, bearing within them the seeds of consumption and scrofula—the entire benumbing of all the moral feel-

ings and intellectual powers, through the monotony of incessant labor and physical exhaustion—habits of sensual gratification rendered inevitable by the extinction of both opportunity and capacity for higher and purer pleasures—the accumulation of masses of a depraved population, utterly unamenable to restraint from priest or magistrate—these were set forth as the certain consequences of the exercise of those arts which conduce most to the comforts of civilised life; and it was said that the foundations of the commercial wealth and manufacturing prosperity of England were erected on the ruins of both the bodies and souls of myriads of our fellow-creatures.

We need not tell our readers how much of distortion and of venom these exaggerated statements derived from feelings of personal and political animosity; and, as was quite natural to expect, a totally opposite set of statements was advanced by those whose character and interests were affected, which deviated from the truth quite as much as the former ones, although in an opposite direction. Thus, by this party, employment in the factories was represented as the means by which (doomed as we are to eat our bread in the sweat of our brows) that sweat is deprived of all its bitterness. The inmates of a factory, we are told, breathe an ever genial atmosphere, free alike from the incursions of cold and the accumulation of close unwholesome vapor. Their labor affords them exercise without fatigue. Their wages suffice for all the reasonable comforts of existence. They are cared for during sickness. In fact, if we were to believe these panegyrists, factory labor is the greatest boon to the population; it is a preventive, if not a curative, of consumption and scrofula; and it secures to them the means of subsistence with the least amount of bodily strength, and with the smallest amount of exposure to the inclemencies of the seasons, and to the risk of injury or mutilation.

Now, in these conflicting statements we have no more than the inevitable result of investigations entered into by men, who, however honestly bent upon discovering the truth, are filled with preconceived ideas of what might reasonably be expected, and who, therefore, can see only the facts which support their own view of the case. We have also the results of that reaction which follows infallibly when men have been led by exaggeration to expect a frightful amount of evil. Finding that things are not quite so bad as was represented, they rush into a totally opposite conclusion, and can see no evil at all. This is most amusingly exemplified in the conversion of Dr. Cooke Taylor, a recent philosophical tourist in the manufacturing districts. “Thus,” to use the words of Mr. Noble’s most praiseworthy pamphlet,* “he

* Facts and Observations relative to the Influence of Manufactures upon Health and Life. By Daniel Noble, Member of the College of Surgeons, &c. &c. London, 1843.