PROVINCIAL MEDICAL JOURNAL

And Retrospect of the Medical Sciences.

No. 148.]

LONDON, SATURDAY, JULY 29, 1843.

PRICE SIXPENCE.
[Stamped Edition Sevenpence.]

CLINICAL LECTURES

DELIVERED AT THE

METROPOLITAN FREE HOSPITAL

BY MR. BENNETT LUCAS,

Senior Surgeon to the Institution, Lecturer on Surgery at the Aldersgate-street School of Medicine, &c.

Lecture X.—April 21, 1843.

GENTLEMEN,—When I last had the pleasure of addressing you, we were engaged in the consideration of the third stage of gonorrhœa, or that of gleet, and I endeavoured to impress upon you the great advantage you will derive, in the treatment of this obstinate affection, by making yourselves, as soon as possible, acquainted with the condition of the urethra. You may adopt one of two methods to ascertain this; the first, and the best, is by the introduction of a bougie, of which I have already spoken; the second is less satisfactory, but, notwithstanding, may be had recourse to either in conjunction with the bougie, or when our patient obstinately refuses the introduction of that instrument. I allude to pressure made with the finger along the urethra, first along its pendulous portion, next in the perineum, and lastly, with the forefinger in the rectum. If pressure, applied in this way, gives uneasiness or slight pain to the patient at any particular part of the urethra, it will be desirable to take blood, either by opening the dorsal vein, by leeches to the part, or by cupping in the perineum; to keep the bowels open by castor oil, electuary of senna, or some mild aperient of a similar kind, and to have recourse to the warm bath. You will also find it advantageous to administer every night, or every other night, some Dover's powder in due doses. This plan of treatment you should only persevere in as long as the urethra is pained upon pressure, or when the pain or uneasiness is increased, or remains undiminished after the bougie has been introduced. When irritability and pain are not present, these remedies will be worse than useless. Our great object, when a discharge from the urethra has become chronic, is to subdue the inflammatory condition of the canal, if such exist. Sometimes the mere introduction of a bougie accomplishes this; how it does so, it is difficult to explain, but sometimes it certainly does. For instance, on introducing the bougie, the patient will complain of great pain, and at the end of a few days all pain and discharge will be at an end; in short, he will be cured; but this happy termination is, as I have already remarked to you, of rare occurrence. If, upon examining the urethra, no pain be given, although the characteristic discharge of gleet be present, you should dispense with the remedies I have just spoken of, and have recourse to those which are calculated to check the discharge. Of these, I have no hesitation in saying injections hold the foremost place, but they are to be very differently applied

from the manner in which I have recommended you to use them in the treatment of the first stage of gonorrhœa, and they are also to be used of a very different strength. You are, of course, aware—at least I presume most of you are—that a great outcry has existed, and still exists, against the use of injections in any stage of gonorrhœa. I believe the fears entertained of the use of these valuable remedies have arisen, in some instances, from their abuse, and in others, from erroneously attributing to them the evil effects induced by other causes; but it is right to remark to you that the prejudice against their use is one which exists rather in the minds of the non-medical public than of the profession, although I am aware that some of its members entertain very strong—I will not say prejudiced—objections against them. It is common to hear a patient, who applies to us for stricture of the urethra—permanent stricture—attribute all his distressing symptoms to injections which he had used, perhaps some years ago, for a gonorrhœa, forgetful altogether of the neglected discharge from the urethra, under which he had suffered for months, or perhaps for years, and in which, no doubt, the contracted state of the canal had its origin; when, had this man used injections of a proper strength, and at a proper time, and in a proper manner, his discharge, in all human probability, would have been speedily arrested, and the stricture have been prevented. It is the subacute inflammatory action, which we know accompanies the third stage of gonorrhœa, being allowed to continue a length of time unheeded, that lays the foundation of all the evil train of consequences which are absurdly attributed to the use of injections; and if I remind you that the symptoms in this stage of gonorrhœa occasion but little inconvenience to a patient, and that, in consequence, he pursues his customary indulgences, with no friendly monitors, in the shape of chordee, to arrest his sexual desires, or severe scalding, to check his after dinner libations, the true reasons why the discharge continues, in spite of all our remedies, become apparent, and why interstitial deposits take place in the urethral submucous tissue, giving rise to the variety of shapes which permanent strictures assume.

At the onset of gonorrhœa, you may recollect I advised you to have recourse to a strong solution of nitrate of silver for an injection; but, in doing so, to be first satisfied that scalding to any great degree did not exist, nor any other symptom of very active inflammation of the parts. Now, in the stage of gleet the mildest injections should be used in the first instance, and, if necessary, should be gradually increased in strength, observing the utmost caution that their use should not give pain to the patient. Injections in these very opposite stages of gonorrhœa should be employed with very opposite intentions. In the first we use them to cut short the specific inflam-

mation, by substituting another, and perhaps a severer one, but of a description which Nature herself either gets rid of in a few days, or which, if she does not, is most amenable to the ordinary rules of treatment. In the second, we employ them to restore the healthy action of a mucous membrane, which has for months, or longer, as the case may be, suffered from a chronic or subacute inflammation, yielding as its product a specific secretion, which, although it gives but little inconvenience beyond its filth, is capable of infecting a healthy urethra with gonorrhœa in its severest form. In the first, moreover, it is admitted upon all hands that at most but the two anterior inches of the urethra are affected, although I myself believe that the inflammation in the first instance is confined to the lacuna magna and its vicinity; but in the second there is no saying to what extent the urethra is affected. Hence, in the former we use a powerful injection; we limit its effects to a certain part of the urethra, and employ it but twice, or at most three times; but in the latter we use a mild one; we inject it frequently in the twenty-four hours, and are anxious that all the affected urethra should feel its influence.

You may have observed me to recommend various kinds of injections in this institution in the treatment of the third stage of gonorrhœa, and in some cases to have merely prescribed the *injection aquea pura*. In the use of this latter simple and often valuable injection it is necessary to keep the patient ignorant of what it is, for most likely, did he know that it was only water, he would empty the contents of his bottle in the street as soon as he had left us, unless he had been initiated into the mysteries of Priessnitz and his followers, when his faith in the remedy would doubtlessly be very great. The injection I have certainly found most beneficial in the treatment of this stage of gonorrhœa is one composed simply of alum and distilled water, in the proportion of from two grains to five grains to the ounce of water. The acetate of zinc I have found the next in efficacy in the proportion of two or three grains to the ounce of water. The acetate of lead is a common injection, and has long been a favorite with many practitioners; but I must say we have been disappointed with its use in this hospital. In the female it is most efficacious, but in the male it is very inefficient. The nitrate of silver injection in this stage of the disease I have given up altogether, and also mercurial injections. Indeed, these latter have had their advocates only in those who support the identity of gonorrhœa and syphilis, a theory which is now all but exploded. There are a variety of other injections which have been recommended from time to time, but they all appear to depend for their efficacy upon their astringency; amongst others, port wine and water and claret and water I have used with the best success in private practice. I can also add my testimony to the value of the following injection, recommended by M. Ricord:

Port wine, six ounces;
Tannin, eighteen grains. Mix.

And which I find is also strongly recommended by Mr. Langston Parker in his valuable little work on Syphilitic Diseases.

You must not suppose, when I say I look upon the alum injection as the best, that I recommend it to the exclusion of the others; on the contrary, if it does

not answer its intention in a week or so, you should change it for another, say the acetate of zinc, and if this does not answer, it is best to have recourse to a third, or to fall back again upon the solution of alum. It is a curious fact that one local application will succeed when another will not in apparently similar cases, and it is equally interesting that when an injection does not stop the discharge, by suspending its use for a few days, and having recourse to it again, it will answer our intentions. The same thing, however, occurs with all our local applications in surgery—in strumous ophthalmia, for instance—in ulcers of various kinds, in the healing of sinuses, and so on.

I have hitherto, Gentlemen, been speaking of the treatment of gonorrhœa in its third stage, after all pain and irritation have been subdued; the discharges from the urethra, which depend upon irritation of certain parts of the canal unconnected with the gonorrhœal virus, which depend upon diseases of the bladder, prostate gland, and testicle; or which are found sometimes to accompany gout or rheumatism in the adult; or which are incidental to some of the diseases of children of both sexes, are very different matters. And I mention this to you at present, lest it might be supposed that I have carelessly passed over other local remedies which are of high value in some of these cases, but which are here inadmissible.

It remains for me now, Gentlemen, to make a few observations on some other remedies given internally in the treatment of this disease, and which are of the utmost value; of these, the preparations of iron hold the foremost place. The muriated tincture of iron is a favorite remedy with us here, in the proportions of from twenty to forty or fifty drops three times a-day in a glass of cold water. You have had daily opportunities of witnessing its good effects in this institution. The following pill, taken three times a-day, is a good formula for administering the sulphate of zinc, which is also a good medicine:—

Sulphate of zinc, two grains;
Extract of gentian, three grains. Mix.

You had an opportunity of witnessing its good effects in the case of H. M., which I quoted in my last lecture.

You will find the sulphate of quinine a good medicine in cases where the discharge is not constantly present, where it is absent for a day or two, and then returns.

The preparations of iodine are constantly administered here, either alone or in combination, particularly if the patient be of a scrofulous habit. Your own judgment will determine the propriety of administering these medicines in the respective cases and the regulation of their doses.

I must not pass over the great benefit to be derived in the treatment of gleet from the use of the cold bath. I have known cases, which have resisted every kind of treatment for many months, be almost instantaneously cured by the use of the plunge bath. In employing this remedy, the object should be to give a shock to the system; not to allow the patient to creep into the bath, to dabble in it for ten or fifteen minutes, and then to creep out again, and take fifteen minutes more to dry himself; but to plunge in, to get out again as quickly as possible, and to dry the body well with a coarse towel. Change of air,

Gentlemen, when your patient can afford it, is no considerable adjuvant to all our remedial agents.

SUCCESSFUL CASE OF TRANSFUSION OF BLOOD.

TO THE EDITORS OF THE PROVINCIAL MEDICAL JOURNAL.

GENTLEMEN,—A desire having been expressed by a correspondent, in the last Number of your Journal, that the particulars of a case mentioned at the Medical Association for Bath and Bristol, in which transfusion of blood was practised, should be published, I think it right to comply with that wish, and shall do it in as few words as possible.

The patient alluded to is a mercantile man, who had been in the habit of travelling in pursuit of his commercial business. He had been a strong, active man, and had lived as persons of his vocation generally do, but more temperately than many. When I saw him first, on the 11th of January, 1834, he had been upwards of two years an invalid, and had been under the care of Mr. Henry Clark, one of the surgeons of the Bristol Infirmary—a medical practitioner of great skill in this city. For some short time he had also been visited by Dr. Riley, and by these gentlemen I was called to see the patient. His early complaints had been of dyspeptic symptoms. These were followed by emaciation and loss of strength. His actual state was that of extreme inanition; his pulse was feeble, jerking, very compressible, the calibre of the artery apparently not filled; he had palpitation of the heart, increased by the slightest exertion, while any effort brought on an approach to syncope. No disease was discoverable by the sounds of the heart or respiratory organs, though some suspicion was entertained of slight dilatation.

The state of the urine alone threw some light on the nature of the disease. It had long deposited a very copious sediment, of a whitish color, slightly tinged with purple, which was redissolved by dilution, with the addition of an alkali. It appeared to consist of lithates, with some chyle; and it was thought probable by the medical gentleman who had previously attended the patient, and by myself at our first meeting, that an exhausted and exsanguineous state, brought on by a long continued draining of the system, constituted the principal disease, or at least that which first required attention. Under these circumstances, it was determined to try a restorative and repletive plan. A nourishing diet was ordered, with malt liquor and some other stimulants. His stomach would receive but little, and at length rejected food. His exhaustion increased to that degree that immediate fatal syncope was threatened. We determined, the patient being obviously *in extremis*, to try the effect of transfusion, and as patients under cholera had borne the injection of large quantities of fluid, there seemed to be no danger in injecting a considerable quantity of blood. Sixteen ounces were taken from the veins of a hale young man, a servant of the patient, and were injected most skilfully by Mr. Clark. The patient was immediately revived and roused. On the following day he appeared much stronger, but complained of some sense of fullness

about his head, and a few drops of blood escaped several times from his nostrils. This subsided; his appetite became good, and he ate plentiful meals of meat, and drank porter, &c. He gradually recovered his strength. The urine improved under the use of alkalies with lime-water, and these were nearly all the remedies used except a few bottles of an effervescing solution of citrate of iron. After two or three months he left his chamber, and then his house, and I understand he is now travelling on commercial business.

I think it may be inferred from the facts of this case that many lives may be saved by injecting good blood into the veins of exhausted patients, and, perhaps, this measure may become hereafter as important a remedial measure as the detraction of blood has been since the time of Hippocrates.

I am, Gentlemen,

Your obedient servant,

J. C. PRICHARD, M.D., F.R.S., M.R.I.A.,

Corresponding Member of the Institute of France, and of the Royal Academy of Medicine, &c.

Bristol, July 18, 1843.

A CASE OF GOUT,

IN WHICH

THE WATER-CURE WAS FOLLOWED BY DISEASED HEART, DROPSY, AND DEATH.

By CHARLES HASTINGS, M.D., F. G. S.

There exists no doubt in my mind that one great cause of the spurious celebrity of the Silesian peasant, Priessnitz, is that he has cunningly succeeded in turning the love of the marvellous, which is known to be a principle inherent in the human breast, to his own selfish ends. Every unprejudiced account which we receive from Graefenburgh confirms this opinion, by showing that there is little faith to be placed in the reputed cures there performed.

The cures said to be effected by the hydropathists are as wonderful as those of Prince Hohenlohe, and somewhat of the same faith which distinguished his disciples animates the hydro-maniacs.

Dr. Pfeufer, directing physician of the Universal Hospital at Bamberg, thus describes the scene of operation of this princely priest, and the rush of the infatuated sufferers from disease to his presence:—"Let us (he says) then imagine thousands and thousands of human beings under such circumstances, to each of whom every day brings new sufferings, new despondence, and who have but one wish remaining—release by death from their nameless pangs. Electrified by the news that at Bamberg and Würzburg there is help and salvation for them through the grace of God, a new life glows within them, and already they feel the term of their misery. With the same longing as that of the leper for the pool of Bethesda, they hurry to the place where all grief vanishes, and where the stream of health and deliverance flows. In ever increasing expectation, and with breasts full of hope, wholly occupied in anticipating the enjoyment of re-established health, dead to every other thought, they approach the scene of healing grace. At every step are announced to them the great wonders that take place there every minute. Scarcely arrived at the place of deliverance, they throw themselves at the foot of the altar, confess their sins, and, reconciled