Published in final edited form as:

Org Lett. 2006 January 5; 8(1): 75–78.

Cis-Dihydroxylation of Unsaturated Potassium Alkyl- and Aryltrifluoroborates

Gary A. Molander* and Ruth Figueroa

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323

Abstract

The *cis*-dihydroxylation of olefin-containing potassium alkyl- and aryltrifluoroborates proceeds readily in moderate to excellent yields. The resulting diols are efficient coupling partners in Suzuki-Miyaura type reactions with both alkenyl- and aryl bromides.

Organoboron compounds have been widely used in organic synthesis.¹ The utility of these organometallic reagents as synthetic tools is often based on the ability of the carbon-boron bond to be oxidized.² It is this characteristic that also limits the opportunity to perform oxidative transformations on other functional groups present in the molecule while still preserving the valuable carbon-boron bond. Common oxidizing reagents are often too powerful and are not well tolerated by many organoboron compounds. Only a few examples of this type of transformation have been reported.^{3,4}

Organotrifluoroborate salts are unique organoboron compounds, notable for their stability to moisture and air.⁵ They are powders or crystalline solids that are easy to access and handle, and these properties have made them attractive synthetic intermediates. The chemistry of these compounds has been expanded to various applications, including Suzuki-Miyaura coupling reactions,^{5a} rhodium-catalyzed 1,4-additions,⁶ as well as allylation of aldehydes⁷ and *N*-toluenesulfonylimines.⁸

Our group has focused on expanding the chemistry of organotrifluoroborates to processes in which the carbon-boron bond remains intact. The epoxidation of olefin-containing potassium organotrifluoroborates demonstrated the tolerance of such organoboron compounds to oxidative conditions such as MCPBA and dimethyldioxirane.⁹ These results led us to consider other oxidative reactions. This communication describes our initial results toward the *cis*-dihydroxylation of olefin-containing potassium organotrifluoroborates. We also demonstrate that the diols produced through this method are valuable synthetic intermediates.

The *cis*-dihydroxylation of olefins is one of the most widely used reactions for the synthesis of vicinal diols.¹⁰ Catalytic OsO₄ and *N*-methylmorpholine *N*-oxide (the Upjohn process) have

been widely applied after the initial publication of this method.¹¹ This process provides the advantage of minimizing overoxidation.¹⁰ With these factors in mind we explored the use of these reagents to accomplish the dihydroxylation of unsaturated organotrifluoroborates (eq 1). Acetone, *t*-BuOH, and water are solvents commonly used in this reaction,¹⁰ and based on our prior experience with the epoxidation reaction we knew that the appropriate solvent ratio would facilitate isolation of the diol by precipitation.⁹ The required potassium organotrifluoroborate salts were prepared following literature protocols¹² involving transmetalation of Grignard reagents with trimethyl borate¹³ or hydroboration of alkenes with the Snieckus reagent¹⁴ (diisopropylprenylborane). This was followed by hydrolysis and subsequent treatment with KHF₂. The products were powders or crystalline white solids that were stored on the benchtop without detectable degradation.

As outlined in Table 1, initial studies were directed toward sterically unhindered organotrifluoroborates containing a remote monosubstituted olefin. The dihydroxylation of potassium 9-decenyltrifluoroborate **1a** proceeded in very good yield using 1.3 mol % of OsO₄ and a 4.5:1:1 acetone:*t*-BuOH:water solvent mixture at room temperature (entry 1). The product was easily isolated as an analytically pure white solid in 75% yield by a simple filtration. The use of ¹H-, ¹³C-, ¹⁹F-, and ¹¹B-NMR as well as elemental analysis corroborated formation of the diol and retention of the valuable trifluoroborate moiety.

We subsequently investigated the dihydroxylation of potassium 4-butenyltrifluoroborate **1b** (entry 2), in which the olefin is closer to the electron rich boron atom. The reaction proceeded smoothly as expected to give the diol in excellent yield. Optimization was required only of the solvent ratio to a mixture of 18:1:1 acetone/*t*-BuOH/water. This new solvent mixture became our standard condition because it facilitated isolation of the diols. In addition, the dihydroxylation of potassium 4-vinylphenyltrifluoroborate **1c** (entry 3) demonstrated that the same reaction conditions are applicable to aryltrifluoroborates.

Our next aim was to carry out the dihydroxylation on more highly substituted olefins. Interestingly, 1,1-disubstituted alkyltrifluoroborate **1d** (entry 4) proved to be sensitive to the reaction conditions. The apparent low yield reflects the inability of all of the product to precipitate in pure form from the reaction mixture. Thus, the majority of the product coprecipitated with an impurity that could not be separated from the desired dihydroxytrifluoroborate under several different isolation protocols. Based on ¹H-, ¹¹B- and ¹⁹F-NMR, we suspect cyclized product **3** as the impurity (Scheme 1), which would be generated by displacement of a fluoride by one of the hydroxy groups of the diol. Similar behavior was also observed in the dihydroxylation of organotrifluoroborate **4**.

Nevertheless, 1,1-disubstituted olefins can be successfully dihydroxylated and isolated as shown by aryltrifluoroborate **1e** (entry 5). Diols from 1,2-disubstituted olefin-containing organotrifluoroborates were also obtained in good yield (entries 6–7). Trisubstituted olefins are usually challenging substrates in *cis*-dihydroxylation.^{10,15} This problem has been attributed to slow hydrolysis of the intermediate osmate(VI) ester, which results in low turnover in the catalytic cycle.^{10,16} The dihydroxylation of potassium 7-methyl-6-octenyl trifluoroborate **1h**, containing a trisubstituted olefin, was achieved in good yield without the need to facilitate the hydrolysis process (entry 8).

The lower yields for diols **2e**, **2g** and **2h** can be attributed to difficulties associated with isolation. These products did not precipitate readily.

The dihydroxylation of potassium allyltrifluoroborate was of particular interest, because the diol produced contains a heteroatom β to the boron. Normally, organoboron compounds with an electronegative atom such as oxygen at the β-position eliminate rapidly via tetracoordination to the boron (eq 2).⁴ This elimination process has been attributed to the thermodynamic stability

of the boron-oxygen bond.¹⁵ To the best of our knowledge, only limited examples of this type of organoboron compound have been reported.^{15,17}

In the event, dihydroxylation of potassium allyltrifluoroborate, **5**, proceeded smoothly under the reaction conditions described previously (Scheme 2).

Isolation of diol **6** was easily achieved using the cation exchange protocol reported by Batey and Quach.¹⁸ Tetracoordination of the boron atom by fluoride as well as the strength of the B-F bond prevented oxidative cleavage and protonolysis of the trifluoroborate moiety in this and our earlier examples;⁶ these characteristics have now been employed to prevent β -elimination as well.

The Suzuki-Miyaura type coupling of potassium organotrifluoroborates has been the subject of extensive study in our group.^{5a} Dihydroxylated alkyl and aryltrifluoroborates were anticipated to be good coupling partners. The presence of free hydroxy groups was deemed unlikely to present a problem because the cross-coupling reactions are often run using alcohols as solvents. Therefore, the diols were subjected to the cross-coupling reactions without the use of protecting groups. The optimized reaction conditions reported by our group for the Suzuki-Miyaura type coupling reaction between alkyltrifluoroborates and aryl bromides employ $\text{PdCl}_2(\text{dppf}) \cdot \text{CH}_2\text{Cl}_2$ (9 mol %) and CsCO_3 (3 equiv) as a base, heated to reflux in a 10:1 THF:H₂O mixture.¹⁹ Thus, with the potassium dihydroxytrifluoroborate **2b** in hand, we conducted a cross-coupling reaction with 4-cyanophenyl bromide under these conditions to afford 4-cyanophenyl-1,2-butanediol **7** in 65% yield after column chromatography (eq 3).

The cross-coupling of alkyltrifluoroborates with alkenyl bromides is part of ongoing research.²⁰ The ideal conditions to date [$\text{PdCl}_2(\text{dppf}) \cdot \text{CH}_2\text{Cl}_2$ (10 mol %), Cs_2CO_3 (3 equiv), toluene/pH 9 buffer (3:1), 90 °C] were used to couple potassium 3,4-dihydroxybutyltrifluoroborate (**2b**) with 3-bromo-2-methyl-2-butene affording the product in good yield (eq 4). This reaction is particularly relevant because it provides an efficient alternative to selective dihydroxylation at a single olefinic site within a diene or polyene system.

The Suzuki coupling of aryltrifluoroborates with aryl bromides has also been successfully developed by our group.²¹ Therefore, potassium 4-(1',2'-dihydroxyethyl) phenyltrifluoroborate, **2c**, was reacted with 4-cyanophenyl bromide under the previously optimized conditions [$\text{Pd}(\text{OAc})_2$ (0.5 mol %), K_2CO_3 (3 equiv), methanol, reflux] to afford the biaryl **9** in good yield (eq 5).

In summary, the first *cis*-dihydroxylation of potassium alkyl- and aryltrifluoroborates has been achieved. The reaction is general for both sterically unhindered and hindered olefins. The olefin-containing organotrifluoroborate salts are readily available by the Grignard approach or selective hydroboration of dienes. The diols were isolated as highly pure solids. Their versatility and value as synthetic units was also demonstrated. We successfully accomplished Suzuki-Miyaura type coupling reactions of the dihydroxylated trifluoroborates. The cross-coupling reactions were carried out without the need for protecting groups. Studies directed toward the asymmetric version of this reaction are underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank the National Institutes of Health (GM 352419), Merck Research Laboratories and Amgen for their generous support. Also, Johnson Matthey is acknowledged for their generous donation of catalysts. Agilent Technologies and Rakesh K. Kohli at the University of Pennsylvania are acknowledged for the determination of high resolution mass spectra of some of the organotrifluoroborates.

References

1. Cragg, GM. Organoboranes in Organic Synthesis. Marcel Decker; New York: 1973. Brown, HC. Organic Syntheses via Boranes. Wiley-Interscience; New York: 1975.
2. Burke, SD.; Danheiser, RL., editors. Oxidizing and Reducing Agents. Handbook of Reagents for Organic Synthesis. Wiley; Chichester: 1999.
3. de Filippis A, Morin C, Thimon C. Synth Commun 2002;32:2669.
4. Matteson DS. J Organomet Chem 1999;581:51.
5. For reviews of organotrifluoroborate salts see: (a) Molander GA, Figueroa R. Aldrichimica Acta 2005;38:49. (b) Darses S, Genêt JP. Eur J Org Chem 2003:4313.
6. (a) Navarre L, Pucheault M, Darses S, Genêt J-P. Tetrahedron Lett 2005;46:4247. (b) Pucheault M, Darses S, Genêt JP. Eur J Org Chem 2002:3552. (c) Pucheault M, Darses S, Genêt JP. Tetrahedron Lett 2002;43:6155. (d) Batey RA, Thadani AN. Org Lett 1999;1:1683.
7. (a) Thadani AN, Batey RA. Tetrahedron Lett 2003;44:8051. (b) Thadani AN, Batey RA. Org Lett 2002;4:3827. [PubMed: 12599469] (c) Batey RA, Thadani AN, Smil DV. Synthesis 2000:990. (d) Batey RA, Thadani AN, Smil DV. Tetrahedron Lett 1999;40:4289.
8. Li SW, Batey RA. Chem Commun 2004;12:1382.
9. Molander GA, Ribagorda M. J Am Chem Soc 2003;125:11148. [PubMed: 16220907]
10. For a review in dihydroxylation see: Schröder M. Chem Rev 1980;80:187.
11. (a) Schneider, WP.; McIntosh, AV. Upjohn. US 2.769.824. (b) VanRheennen V, Kelly RC, Cha DY. Tetrahedron Lett 1976;17:1973.
12. (a) Vedejs E, Fields SC, Hayashi R, Hitchcock SR, Powell DR, Schrimpf MR. J Am Chem Soc 1999;121:2460. (b) Garrett CE, Fu GC. J Org Chem 1996;61:3224. (c) Pereira S, Srebnik M. J Am Chem Soc 1996;118:909. (d) Vedejs E, Chapman RW, Fields SC, Lin S, Schrimpf MR. J Org Chem 1995;60:3020. (e) Kabalka GW, Narayana C, Reddy NK. Synth Commun 1994;24:1019. (f) Burgess K, Ohlmeyer MJ. Chem Rev 1991;91:1179. (g) Manning D, Nöth H. Angew Chem, Int Ed Engl 1985;24:878. (h) Brown HC, Bhat NG, Somayaji V. Organometallics 1983;2:1311.
13. Matteson DS. Tetrahedron 1989;45:1859.
14. Kalinin AV, Scherer S, Snieckus V. Angew Chem, Int Ed Engl 2003;42:3399. [PubMed: 12888969]
15. Ray R, Matteson DS. Tetrahedron Lett 1980;21:449.
16. (a) Akashi K, Palermo RE, Sharpless BK. J Org Chem 1978;43:2063. (b) Sharpless BK, Akashi K. J Am Chem Soc 1976;98:1986.
17. Pereira S, Srebnik M. Tetrahedron Lett 1996;37:3283.
18. Batey RA, Quach TD. Tetrahedron Lett 2001;42:9099.
19. (a) Molander GA, Ito T. Org Lett 2001;3:393. [PubMed: 11428022] (b) Molander GA, Yun CS, Ribagorda M, Biolatto B. J Org Chem 2003;68:5534. [PubMed: 12839444]
20. Dave Seapy, unpublished results.
21. Molander GA, Biolatto B. J Org Chem 2003;68:4302. [PubMed: 12762730]

Scheme 1.

Scheme 2.

Table 1
Dihydroxylation of Potassium Alkyl- and Aryltrifluoroborates^a

entry	substrate	time (h)	product	yield (%) ^e
1 ^b	(1a)	2	(2a)	75
2	(1b)	3	(2b)	88
3	(1c)	2	(2c)	84
4	(1d)	4	(2d)	30
5	(1e)	6	(2e)	56
6 ^c	(1f)	2	(2f)	53 ^f
7	(1g)	1	(2g)	71
8, ^{cd}	(1h)	6	(2h)	69

^a The reactions were conducted using OsO₄ (1.3 mol %), *N*-methylmorpholine *N*-oxide (1.0–1.5 equiv) and an acetone/*t*-BuOH/water mixture (18:1:1) at room temperature.

^b Solvent ratio was 4.5:1:1.

^c Solvent ratio was 60:1:1.

^d 0.62 mol % OsO₄.

^e Isolated yields.

^f Mixture of diastereomers in a 2:1 ratio.