Published in final edited form as:

J Org Chem. 2006 September 15; 71(19): 7491–7493.

Linchpin Synthons: Metalation of Aryl Bromides Bearing a Potassium Trifluoroborate Moiety

Gary A. Molander* and Noel M. Ellis

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323

Abstract

Aryl bromides bearing a potassium trifluoroborate moiety were subjected to lithium-halogen exchange at low temperature using a variety of alkyllithium reagents. A number of different electrophiles were evaluated in their reactions with the aryllithiums produced therein. Under carefully optimized conditions, potassium bromophenyl trifluoroborates afforded good to excellent yields of the corresponding alcohols (64–94% isolated yield) when aldehydes or ketones were used as the electrophilic partner. Esters were unfortunately found to be unreactive.

Metal-halogen exchange reactions have had a rich history in synthetic organic chemistry, specifically with respect to the formation of C-C bonds.¹ However, the tendency of boronate esters and boranes to act as a Lewis acid toward alkyllithiums and Grignard reagents has prevented the functionalization of boron-containing compounds via a metal-halogen exchange route. To date, there exists only one report in which a metal-halogen exchange, specifically a magnesium-halogen exchange, was used in the presence of a boron-containing functional group.² That method employed expensive iodoaryl pinacol boronates as substrates. Concurrent to that development, our continuing interest in the chemistry of potassium organotrifluoroborates led us to speculate on their use in metal-halogen exchange reactions. In recent years, organotrifluoroborates have proven to be valuable synthetic intermediates, participating in a range of palladium-catalyzed cross-couplings,^{3a,4} rhodium-catalyzed conjugate additions,⁵ and allylations of aldehydes⁶ as well as *N*-toluenesulfonylimines.⁷ Moreover, potassium organotrifluoroborate salts are easily prepared by the addition of inexpensive KHF₂ to various organoboron intermediates,⁸ and they are air- and moisture-stable crystalline solids that are readily isolated.^{3,8} Metalation of organotrifluoroborates would give rise to useful bimetallic linchpins capable of sequential functionalization at each of the two metal centers (Scheme 1), thus increasing molecular complexity in a convergent manner. In this Note we report the realization of these bimetallic linchpin reagents using the lithium-halogen exchange of potassium bromophenyl trifluoroborates.

Initial studies focused on the lithium-halogen exchange of potassium *p*-bromophenyl trifluoroborate (**1**). While methanol, acetonitrile, and acetone are known to be excellent solvents for a wide array of potassium organotrifluoroborates,³ all of these commonly employed solvents are far too acidic to remain inert under the strongly basic conditions of the metal-halogen exchange reaction. Out of necessity, it was discovered that THF also serves as an excellent solvent for a number of potassium organotrifluoroborates. Because of its ease of workup and known spectral properties, potassium phenyl trifluoroborate (**2**)⁹ made a logical target to probe the lithium-halogen exchange reaction of **1**. Unfortunately, early attempts at the reaction of **1** with 1 equiv of *n*-BuLi in THF at –78 °C, followed by a variety of aqueous

gmolandr@sas.upenn.edu.

quenches (Scheme 2), met with little success. Although the oils isolated from these early reactions were shown to contain “ate” complexes of boron by ^{11}B NMR, standard precipitation in diethyl ether did not yield a white crystalline solid, even though both ^1H and ^{13}C spectra of the crude oils were also consistent with **2**. Changing the workup protocol improved the situation to some degree. Thus, an aqueous quench with satd $\text{KHF}_2(\text{aq})$ led to a modest yield of KPhBF_3 (~30%). Undaunted by these disappointing initial results, we proceeded to investigate the quenching of the presumed organolithium intermediate with carbon electrophiles. In fact, utilization of cyclohexanecarboxaldehyde (**3**) as the electrophile led to the isolation of a white crystalline solid whose spectra were entirely consistent with the desired product (**4**) in 79% yield (Scheme 2).

Careful optimization studies were carried out on the preparation of alcohol **6** (Table 1). Variation of solvents (DME or THF) and molarity revealed that the best results were consistently obtained at 0.1 M in THF (85% yield). Though additives that decrease the aggregation state of *n*-BuLi have been shown to increase the rates of lithium-halogen exchange reactions,¹⁰ addition of 1 equiv of TMEDA gave **6** in only 30% isolated yield. Variation of lithium sources showed that *n*-BuLi was preferable to *s*-BuLi (40% yield) and *t*-BuLi (22% yield). Experiments to determine the time required for lithium-halogen exchange are summarized in Table 1. Remarkably, it was found that the best yields were obtained when the introduction of the electrophile was delayed as much as 1 h following the addition of *n*-BuLi at -78°C (87% yield). Longer delays led to only small decreases in isolated yield. Last, but not least, it was found while optimizing the reaction of **23** to **24** (eq 1) that the $\text{KHF}_2(\text{aq})$ quench proceeded very rapidly, with times longer than 10 min leading to diminished yields of the product.

In general, lithium-halogen exchanges followed by addition to aliphatic or aromatic aldehydes gave good to excellent yields of the secondary alcohols (Table 2, entries 1–4). As for α,β -unsaturated aldehydes, reaction of acrolein (**11**) provided solely the 1,2-addition product (**12**). Ketones **13** and **15** underwent smooth conversion to tertiary alcohols **14** and **16**, respectively. Esters, however, proved unreactive under the standard reaction conditions. Other classes of electrophiles also proved suitable to the method. Silylation with TMSCl (**17**) provided silyl trifluoroborate **18** in good yield. Iodination using I_2 (**19**) proceeded as expected to provide *p*-iodophenyl trifluoroborate (**20**). Finally, phenylisocyanate (**21**) reacted to give the amide-containing trifluoroborate (**22**).

Less general were the reactions of *meta*-substituted potassium bromoaryl trifluoroborates and various heteroaryl bromides. TMSCl (**17**) was found to react with the aryllithium arising from **23** to give **24** in good yield when *t*-BuLi was used as the metallating agent (Scheme 3). As a rule, however, the use of other electrophiles was hindered by the low crystallinity of the products, which frequently gave oils that could not be further purified. In these cases, direct conversion of the crude potassium trifluoroborate to the tetrabutylammonium trifluoroborate was possible, but did not effect a satisfactory purification of the products.¹¹ Various heteroaryl bromides were explored but were found to give results analogous to those using **23**.

In summary, a new entry into functionalized boron compounds has been developed via the lithium-halogen exchange reaction of potassium organotrifluoroborates. The method is less expensive and simpler than existing protocols.² Moreover, the trifluoroborates produced according to the method presented here may be cross-coupled following previously developed protocols for the Suzuki-Miyaura cross-coupling of aryl trifluoroborates.¹² Further studies aimed at widening the scope of this method will be the subject of future communications.

Experimental Section

General Experimental Procedure for the Lithium-Halogen Exchange of Potassium Bromoaryl Trifluoroborates. Preparation of Potassium 4-(1-Hydroxypropyl)phenyl Trifluoroborate (10)

To an oven-dried round bottom flask (50 mL) was added *p*-bromophenyl trifluoroborate (262 mg, 1.0 mmol). The flask was evacuated and back-filled with N₂ three times before adding freshly distilled THF (10 mL). A Dry Ice/acetone bath was then used to cool the reaction to -78 °C and *n*-BuLi (1.0 mmol) was added dropwise. The reaction was then left to stir at -78 °C for 1 h, during which time an insoluble salt precipitated. Next, neat propionaldehyde (58 mg, 1.0 mmol) was added dropwise, and stirring was continued for an additional 10 min at -78 °C. After warming to rt, the reaction was stirred at rt for 30 min, then satd KHF₂ (aq) (390 mg, 5.0 mmol) was added, and the reaction was stirred at rt for 10 min. Evaporation of the solvents *in vacuo* was followed by drying at 0.05 torr for a minimum of 6 h. Acetone extraction (3 × 20 mL) and filtration of the solids gave a solution of the product trifluoroborate in acetone. Reduction of the solvent followed by dropwise addition of Et₂O led to precipitation of the product. The product was then filtered, collected, and dried overnight at 0.05 torr to afford potassium 4-(1-hydroxypropyl)phenyl trifluoroborate (212 mg, 88%). mp > 250 °C. ¹H-NMR (500 MHz, DMSO-*d*₆): 7.26 (d, *J* = 7.8 Hz, 2H), 7.0 (d, *J* = 7.8 Hz, 2H), 4.82 (m, 1H), 4.31 (m, 1H), 1.58 (m, 2H), 0.80 (t, *J* = 7 Hz, 3H); ¹³C-NMR (125.8 MHz, DMSO-*d*₆): 142.3, 130.9, 123.9, 74.2, 32.1, 10.3; ¹⁹F-NMR (471 MHz, DMSO-*d*₆): -139.2; ¹¹B-NMR (128.37 MHz, DMSO-*d*₆): 3.37; IR (KBr) = 3403, 2964 cm⁻¹; HRMS (ESI) calcd. for C₉H₁₁OBF₃ (M⁻) 203.0855, found 203.0845.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was generously supported by the National Institutes of Health (GM35249), Amgen, and Merck Research Laboratories. N.M.E acknowledges the generous financial support of the University of Pennsylvania through a Douty Fellowship. We would also like to extend thanks to Frontier Scientific for their generous donation of boronic acids and Dr. Rakesh Kohli for high-resolution mass spectrometry.

References

- (a) Bailey WF, Patricia JJ. *J Organomet Chem* 1988;352:1–46. (b) Knochel P, Dohle W, Gommerman N, Kneisel FF, Kopp F, Korn T, Sapountzis I, Vu VA. *Angew Chem, Int Ed* 2003;42:4302–4320. (c) Parham WE, Bradsher CK. *Acc Chem Res* 1982;15:300–305. (d) Rogers HR, Houk J. *J Am Chem Soc* 1982;104:522–525. (e) Rieke RD, Lee J, Velarde-Ortiz R, Guijarro A, Wurst JR. *J Org Chem* 2000;65:5428–5430. [PubMed: 10993378] (f) Slocum DW, Carroll A, Dietzel P, Eilerman S, Culver JP, McClure B, Brown S, Holman RW. *Tetrahedron Lett* 2006;47:865–868. (g) Oshima K, Inoue A, Kitagawa K, Shinokubo H. *J Org Chem* 2001;66:4333–4339. [PubMed: 11397172]
- Knochel P, Baron O. *Angew Chem Int, Ed* 2005;44:3133–3135.
- For reviews of organotrifluoroborate salts, see: (a) Molander GA, Figueroa R. *Aldrichimica Acta* 2005;38:49–56. (b) Darses S, Genêt J-P. *Eur J Org Chem* 2003:4313–4327.
- (a) Molander GA, Felix LA. *J Org Chem* 2005;70:3950–3956. [PubMed: 15876083] (b) Molander GA, Figueroa R. *Org Lett* 2006;8:75–78. [PubMed: 16381571] (c) Cella R, Cunha RLOR, Reis AES,

- Pimenta DC, Klitzke CF, Stefani HA. *J Org Chem* 2006;71:244–250. [PubMed: 16388643] (d)
Molander GA, Sommers EM, Baker SR. *J Org Chem* 2006;71:1563–1568. [PubMed: 16468806] (e)
Molander GA, Yokoyama Y. *J Org Chem* 2006;71:2493–2498. [PubMed: 16526802]
5. (a) Batey RA, Thadani AN. *Org Lett* 1999;1:1683–1686. (b) Pucheault M, Darses S, Genêt J-P. *Tetrahedron Lett* 2002;43:6155–6157. (c) Pucheault M, Darses S, Genêt J-P. *Eur J Org Chem* 2002:3552–3557. (d) Navarre L, Pucheault M, Darses S, Genêt J-P. *Tetrahedron Lett* 2005;46:4247–4250.
6. (a) Batey RA, Thadani AN, Smil DV. *Tetrahedron Lett* 1999;40:4289–4292. (b) Batey RA, Thadani AN, Smil DV. *Synthesis* 2000:990–998. (c) Thadani AN, Batey RA. *Org Lett* 2002;4:3827–3830. [PubMed: 12599469] (d) Thadani AN, Batey RA. *Tetrahedron Lett* 2003;44:8051–8055.
7. Li S-W, Batey RA. *Chem Commun* 2004;12:1382–1383.
8. (a) Vedejs E, Chapman RW, Fields SC, Lin S, Schrimpf MR. *J Org Chem* 1995;60:3020–3027. (b) Vedejs E, Fields SC, Hayashi R, Hitchcock SR, Powell DR, Schrimpf MR. *J Am Chem Soc* 1999;121:2460–2470.
9. Genêt J-P, Darses S, Brayer J-L, Demoute J-P. *Tetrahedron Lett* 1997;38:4393–4396.
10. Korneev SM, Kaufmann DE. *Synthesis* 2002:491–496.
11. Flash chromatography using CHCl_3 with MeOH or MeCN was tried in an attempt to purify the tetrabutylammonium organotrifluoroborates. With the exception of one or two compounds, attempts at SiO_2 chromatography using this solvent system resulted in decomposition of the trifluoroborates to the corresponding boronic acids as indicated by ^{11}B NMR.
12. Molander GA, Biolatto B. *Org Lett* 2002;4:1867–1870. [PubMed: 12027634]

Scheme 1.

Scheme 2.

Table 1
Variation of the Duration of Lithium-Halogen Exchange

time with RLi prior to RCHO addition (min)	% yield
5	70
30	78
60	87
120	76
240	78

Table 2Optimized Lithium-Halogen Exchange Reactions of *p*-Bromophenyl Trifluoroborate (1)

<div> </div>			
entry	E ⁺	Product	% yield (isolated)
1			84
2			87
3			62
4			93
5			94
6			83

entry	E^+	Product	% yield (isolated)
7	15	16	65
9	TMSCl 17	18	67
10	I_2 19	20	64
11	21	22	68