Published in final edited form as:

J Org Chem. 2008 August 1; 73(15): 6000–6002.

Facile Synthesis of Highly Functionalized Ethyltrifluoroborates

Gary A. Molander^{*,†}, Wilma Febo-Ayala[†], and Montserrat Ortega-Guerra^{†,‡}

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania, 19104-6323

Abstract

Organotrifluoroborates are generating increased interest because of their ease of preparation and purification, and indefinite shelf-life. Herein we report the preparation of Organotrifluoroborates bearing functional groups that can be manipulated at different stages of the synthetic route, exploiting the inertness of their carbon-boron bonds. The alkylation of 2,2-dicyanoethyltrifluoroborate with a variety of electrophiles and of (EWG)₂CH₂ with potassium iodomethyltrifluoroborate resulted in di- and trisubstituted ethyltrifluoroborates in good to excellent yields.

Tricoordinate boronic acids are sensitive to bases, nucleophiles, and oxidants owing to their acidic protons and vacant p-orbital. Consequently, they cannot be carried through synthetic steps involving these reagents. This has limited the preparation of functionalized boronic acids under basic or nucleophilic conditions. The increased stability of boronate esters allows them to survive functionalization under some, but not all, of these conditions.

Organotrifluoroborates have emerged as a versatile alternative to boronic acids and boronate esters. They are tetracoordinate boron salts prepared from the addition of inexpensive KHF₂ to organoboron intermediates.¹ The absence of an empty p-orbital makes them mechanistically inert to many reagents utilized in organic synthesis, allowing their installation early on in a synthetic sequence. Potassium Organotrifluoroborates are crystalline solids that are air- and moisture stable, easy to handle, and can be stored indefinitely without special precautions. While aryl-, vinyl- and allyltrifluoroborates find use in a variety of applications,¹ alkyltrifluoroborates remain relatively unexplored, with a few notable applications, including their use in the Suzuki-Miyaura cross-coupling² as well as in the preparation of secondary amines³ and alkyl iodides.⁴

Despite the advantages and increasing applicability of alkyltrifluoroborates, to date they are normally prepared by three general methods: i) from commercially available boronic acids or boronate esters,⁵ ii) transmetalation of organolithium or Grignard reagents,⁶ and iii) hydroboration of alkenes.⁷ For that reason, access to more highly-elaborated or unique Organotrifluoroborates is an area of synthetic interest. Recently, we described a new method for the preparation of a wide range of Organotrifluoroborates, including potassium 2,2-dicyanoethyltrifluoroborate (**2**), by direct nucleophilic substitution of potassium iodomethyltrifluoroborate (**1**).⁸ Few alkylborons have been elaborated by nucleophilic reactions at ancillary electrophilic functional groups. Herein, we expand the functional diversity of Organotrifluoroborates by preparing both di- and trisubstituted ethyltrifluoroborates using just such a strategy.

gmolandr@sas.upenn.edu.

[†]Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19104-6323.

[‡]Departamento de Química Orgánica, Universidad Autónoma de Madrid. Ciudad Universitaria de Cantoblanco, 28049, Madrid.

The gram scale synthesis of the starting potassium 2,2-dicyanoethyltrifluoroborate (**2**) was achieved in 92% yield (eq 1). Malononitrile (3 equiv) was deprotonated with sodium hydride (3 equiv), followed by the addition of potassium iodomethyltrifluoroborate (1 equiv).⁸ After completion, the reaction was quenched with potassium hydrogen fluoride to insure counterion homogeneity, and the product was purified by precipitation from acetone-ether.⁹

(1)

With **2** in hand we investigated suitable reaction conditions for its subsequent alkylation. Initially, sodium hydride was used as the base, and the highly reactive iodomethane was utilized as the electrophile. The first solvent tested was THF, which either resulted in no reaction or complex reaction mixtures at longer reaction times. Hexamethylphosphoramide (HMPA) was added to facilitate solubilization of the anion, resulting in a 64% yield of the desired product. To avoid the use of toxic HMPA, more polar solvents (4:1 THF:DMF, DMSO, and DMF) were tested. After reacting for 4 h, all of these protocols resulted in higher yields of the desired product, with the best yields obtained using DMF. Three equivalents of the electrophile were needed - use of 1 or 2 equivalents gave lower yields. This set of conditions gave product **4a** in 82% yield after purification (Table 1), and the method was extended to other electrophiles with the results outlined in Table 1.

All of the products were obtained in good to excellent yields. Of note, we find no evidence for formation of the allenyl product upon using the propargyl bromide (**3g**) as an electrophile in the reaction (entry 7). Some electrophiles were ineffective in the substitution reactions. For example, only **2** was recovered when benzyl bromoacetate, 2-bromoacetophenone, and bromomethyl benzyl ether were used as the electrophiles.

Unfortunately, attempts to substitute **1** directly with nucleophiles derived from substrates such as **5a** under the conditions described in eq 1 were unsuccessful. The use of different bases (KH, NaOH, KOH, K₂CO₃) and solvents (DMSO, CH₃CN), or phase transfer catalysts did not give any product. The reaction conditions described in eq 1 along with a higher temperature (80 °C) and longer reaction time (3 d) in DMF yielded some product along with decomposition of materials as indicated by ¹H NMR. The attenuated electrophilic character of the iodomethyltrifluoroborate represents a challenge when highly enolizable or bulkier nucleophiles were employed.

An alternative route was explored starting from the pinacol iodomethylboronate (**6**),¹⁰ which reacts by an entirely different mechanism than the iodomethyltrifluoroborate. Thus, whereas the organotrifluoroborates appear to react by a direct S_N2 reaction on the halide, the pinacolboronates react by initial complexation of the nucleophile at the boron, generating an intermediate “ate” complex. This is followed by an α-transfer process in which the nucleophile migrates, displacing the halide intramolecularly. Reaction of **6** with various stabilized anions followed by the addition of KHF₂ thus afforded the desired organotrifluoroborates in good yields (Table 2). Only an equimolar amount of base, electrophile, and nucleophile was required to afford the product. A significant advantage of the organotrifluoroborate products over that of the corresponding boronates is the easy purification of the former products *via* acetone-ether precipitation. In contrast, the boronate esters are usually purified by distillation at high temperatures.¹¹

In summary, the high yielding synthesis of functionalized ethyl trifluoroborates has been described, including the 1 g synthesis of **2**. The scope and limitations in the alkylation of **2** were further described, demonstrating the manipulation and stability advantages of organotrifluoroborates over boronic acids and boronate esters. Studies are underway to optimize the Suzuki-Miyaura cross-coupling reaction conditions for these materials.

Experimental Section

Potassium 2,2-Dicyanoethyltrifluoroborate (**2**)

The title compound was prepared according to the literature procedure⁸ from potassium iodomethyltrifluoroborate (**1**) (1.00 g, 4.0 mmol) and the malononitrile anion generated by treatment of malononitrile (800.6 mg of a 99% purity reagent, 12.0 mmol) with NaH (484.5 mg of a 60% dispersion in oil, 12.0 mmol), affording 765 mg (92% yield) of the title compound as a pink solid whose spectroscopic data agree with those reported in the literature.⁸

General Experimental Procedure for the Alkylation of Potassium 2,2-Dicyanoethyltrifluoroborate. Preparation of Potassium 2,2-Dicyanobutyltrifluoroborate (**4b**)

To a solution of potassium 2,2-dicyanoethyltrifluoroborate (**2**) (100 mg, 0.53 mmol) in 500 μ L of DMF was sequentially added NaH (25 mg of a 60% dispersion in mineral oil, 0.64 mmol) and iodoethane (127 μ L, 1.59 mmol). The resulting suspension was stirred at 25 °C until the reaction was complete (4.5 h) as indicated by ¹⁹F NMR. The reaction was then quenched with 1 N KHF₂ (0.5 mL). The solvents were removed under high vacuum, the solid was re-dissolved in HPLC grade acetone (5 mL) and the insoluble salts were filtered off. After concentration, the resulting solid was re-dissolved in the minimum quantity of HPLC grade acetone (1 mL) and precipitated by adding Et₂O (3 mL). After decantation, the resulting solid was pulverized using a mortar and pestle and washed with Et₂O (3 mL) to give a white solid (111 mg, 98% yield), mp > 250 °C. ¹H NMR (500 MHz, DMSO-*d*₆) δ : 1.94 (q, *J* = 7.3 Hz, 2H), 1.07 (t, *J* = 7.3 Hz, 3H), 0.76 (br s, 2H). ¹³C NMR (125.8 MHz, DMSO-*d*₆) δ : 118.5 (2C), 31.6, 9.9. ¹⁹F NMR (470.8 MHz, DMSO-*d*₆) δ : -135.2. ¹¹B NMR (128.4 MHz, DMSO-*d*₆) δ : 1.91 (q, *J* = 52.5 Hz). IR (neat): 2937, 2250, 1683, 1463, 1269, 1024, 926, 770, 483 cm⁻¹. HRMS (*m/z*): calcd. for C₆H₇BF₃N₂, 175.0654 [M-K]⁺, found 175.0654.

General Experimental Procedure for the Alkylation with Pinacol Iodomethylboronate. Preparation of Potassium Methyl 2-Cyano-3-(trifluoroborato) Propanoate (**7b**)

A flame dried round bottom flask with a stir bar was charged with 95% NaH dispersion in mineral oil (14 mg, 0.54 mmol) followed by anhydrous THF (1.7 mL). The flask was cooled to 0 °C and methyl cyanoacetate (53.4 mg, 0.54 mmol) was added dropwise via syringe. The reaction was slowly warmed to 25 °C and stirred under N₂ for 2 h. Next, the pinacol iodomethylboronate (141.8 mg, 0.53 mmol) was added dropwise and stirred at 25 °C until the reaction was complete as indicated by ¹¹B NMR (12 h). The reaction was quenched with ~1 mL of H₂O and extracted using Et₂O (~3 mL). The organic layer was washed with H₂O (2×2 mL) and brine (1×2 mL). The product was dried (MgSO₄), filtered, and concentrated. (Note: In one instance the aqueous work up was not performed and the yield of the product was unaffected.) The crude product was re-dissolved in 5.25:1 MeOH:H₂O (1.35 mL) followed by the addition of KHF₂ (124.2 mg, 1.59 mmol). The reaction was stirred at 25 °C until completion as indicated by ¹¹B NMR (4 h). The solvent was then removed under high vacuum and the material re-dissolved in HPLC grade acetone (10 mL). The insoluble salts were filtered off and the filtrate was concentrated. The crude product was re-dissolved in the minimum amount of HPLC grade acetone (2 mL) necessary and precipitated by adding Et₂O (6 mL). The solid was collected *via* decantation to give a light yellow solid (88.4 mg, 76% yield), mp = 100–105 °C. ¹H NMR (500 MHz, acetone-*d*₆) δ : 3.69 (s, 3H), 3.44 (t, *J* = 7.5 Hz, 1H), 0.77 (d, *J* = 34.5 Hz, 2H). ¹³C NMR (125.8 MHz, acetone-*d*₆) δ : 171.1, 120.7, 53.1, 34.9. ¹⁹F NMR (470.8

MHz, acetone- d_6) 5: -142.32 (s). ^{11}B NMR (128.4 MHz, acetone- d_6) δ : 4.35 (s). IR (neat): 2958, 2254, 1740, 1439, 1259, 1036 cm^{-1} . HRMS (m/z): calcd. for $\text{C}_5\text{H}_6\text{BF}_3\text{NO}_2$, 180.0444 [$\text{M}-\text{K}^+$], found 180.0422.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors wish to thank Mr. Adam Brown (University of Pennsylvania) for his contribution to the project. Dr. Rakesh Kohli (University of Pennsylvania) is acknowledged for obtaining high-resolution mass spectra of the new compounds. This work was supported by the NIH (GM35249) and the Spanish Ministerio de Educación y Ciencia FPU grant.

References

1. a) Darses S, Genêt JP. Chem Rev 2008;108:288–325. [PubMed: 18095714] b) Molander GA, Ellis N. Acc Chem Res 2007;40:275–286. [PubMed: 17256882] c) Stefani HA, Cella R, Vieira AS. Tetrahedron 2007;63:3623–3658. d) Molander GA, Figueroa R. Aldrichim Acta 2005;38:49–56.
2. a) Molander GA, Ito T. Org Lett 2001;3:393–396. [PubMed: 11428022] b) Molander GA, Yun CS, Ribagorda M, Biolatto B. J Org Chem 2003;68:5534–5539. [PubMed: 12839444] c) Molander GA, Ham J, Seapy DG. Tetrahedron 2007;63:768–775. [PubMed: 18196177] d) Molander GA, Ribagorda M. J Am Chem Soc 2003;125:11148–11149. [PubMed: 16220907]
3. Matteson DS, Kim GY. Org Lett 2002;4:2153–2155. [PubMed: 12074655]
4. Kabalka GW, Mereddy AR. Tetrahedron Lett 2004;45:343–345.
5. a) Vedejs E, Chapman RW, Fields SC, Lin S, Schrimpf MR. J Org Chem 1995;60:3020–3027. b) Vedejs E, Fields SC, Hayashi R, Hitchcock SR, Powell DR, Schrimpf MR. J Am Chem Soc 1999;121:2460–2470.
6. Matteson DS. Tetrahedron 1989;45:1859–1885.
7. a) Brown HC, Bhat NG, Somayaji V. Organometallics 1983;2:1311–1316. b) Burgess K, Ohlmeyer MJ. Chem Rev 1991;91:1179–1191. c) Pereira S, Srebnik M. J Am Chem Soc 1996;118:909–910. d) Kabalka GW, Narayana C, Reddy NK. Synth Commun 1994;24:1019–1023. e) Männig D, Nöth H. Angew Chem, Int Ed Engl 1985;24:878–879. f) Garrett CE, Fu GC. J Org Chem 1996;61:3224–3225.
8. Molander GA, Ham J. Org Lett 2006;8:2031–2034. [PubMed: 16671774]
9. Alternatively, 2 can be prepared from pinacol iodomethylboronate (6), see the Supporting Information section for experimental details.
10. Smoum R, Rubinstein A, Srebnik M. Bioorg Chem 2003;31:464–474. [PubMed: 14613767]
11. Kinder DH, Ames MM. J Org Chem 1987;52:2452–2454.

TABLE 1

Alkylation of Potassium 1,2-Dicyanoethyltrifluoroborate (2)

<div style="text-align: center;"> <p> $\text{NC}-\text{CH}(\text{CN})-\text{CH}_2\text{BF}_3\text{K} \xrightarrow[\text{DMF, 25 } ^\circ\text{C, 4 h}]{\begin{matrix} 1) \text{ NaH} \\ 2) \text{ R-X (3a-h)} \\ 3) \text{ 1 N KHF}_2 \end{matrix}} \text{NC}-\text{C}(\text{R})(\text{CN})-\text{CH}_2\text{BF}_3\text{K}$ </p> <p>2 4a-h</p> </div>			
entry	R-X	product	isolated yield (%)
1	ICH ₃ 3a		82
2	ICH ₂ CH ₃ 3b		98
3	ICH(CH ₃) ₂ 3c		96
4	ICH(CH ₃)CH ₂ CH ₃ 3d		95
5	I(CH ₂) ₅ CH ₃ 3e		97
6 ^a	ICH ₂ CH=CH ₂ 3f		43
7	BrCH ₂ C≡CH 3g		83
8	BrCH ₂ Ph 3h		90

Reaction conditions: substrate (1 equiv), NaH (1.2 equiv), electrophile (3 equiv)

^a = substrate (1 equiv), NaH (1.1 equiv), electrophile (1.1 equiv)

TABLE 2

Preparation of Di- and Trisubstituted Ethyltrifluoroborates

<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center; margin-right: 20px;"> </div> </div>			
entry	nucleophile	product	isolated yield (%)
1	5a	7a	62
2	5b	7b	76
3	5c	7c	70
4	5d	7d	75
5	5e	7e	64
6 ^a	5f	7f	72

Reaction conditions: nucleophile (1 equiv), NaH (1 equiv), electrophile (1 equiv)