Published in final edited form as:

Neurosci Res. 2008 May ; 61(1): 56–69.

Ocular following responses of monkeys to the competing motions of two sinusoidal gratings

K. Matsuura¹, K. Miura², M. Taki³, H. Tabata¹, N. Inaba¹, K. Kawano¹, and F. A. Miles⁴

¹ Department of Integrative Brain Science Graduate School of Medicine Kyoto University, Japan

² Nano-Medicine Merger Education Unit Kyoto University, Japan

³ Department of Otolaryngology, Head and Neck Surgery Kyoto Prefectural University of Medicine Kyoto, Japan

⁴ Laboratory of Sensorimotor Research National Eye Institute National Institutes of Health Bethesda, MD 20892

Abstract

Ocular following responses (OFRs) were elicited in monkeys at short latencies (~50ms) by applying motion in the form of successive ¼-wavelength steps to each of two overlapping vertical sine-wave gratings that had different spatial frequencies. In the first experiment, the two sine waves had spatial frequencies in the ratio 3:5 and moved in opposite directions. The initial OFRs showed a highly nonlinear dependence on the relative contrasts of the competing sine waves. On average, when the contrast of one was less than a third of that of the other then the one with the lower contrast became ineffective—as though suppressed—and the OFR was entirely determined by the sine wave of higher contrast: winner-take-all. In a second experiment, the two sine waves had spatial frequencies in the ratio 3:7 and moved in the same direction (though at different speeds). The initial OFRs again showed a highly nonlinear dependence on the relative contrasts of the competing sine waves, with a winner-take-all outcome when the contrasts of the two sine waves were sufficiently different. In both experiments, the nonlinear dependence on the relative contrasts of the competing sine waves was well described by a contrast-weighted-average model with just two free parameters. These findings were very similar to those of Sheliga et al. (2006c) on the human OFR, indicating that the monkey is a good animal model for studying the nonlinear interactions that emerge when competing motions are used.

Keywords

visual motion; ocular following responses; winner-take-all; nonlinear interactions; mutual inhibition

1. Introduction

In primates, sudden movements of the visual scene elicit ocular following responses with ultra-short latencies: <75 ms in humans (Gellman et al., 1990; Sheliga et al., 2005a, b) and <60 ms in monkeys (Miles et al., 1986; Miura et al., 2006). It is generally believed that there are at

Address correspondence to: Kenichiro Miura, PhD, Nano-Medicine Merger Education Unit, Kyoto University, Konoe-cho, Yoshida, Kyoto-shi, Kyoto 606-8501, Japan, e-mail: kmiura@brain.med.kyoto-u.ac.jp, phone: +81-75-753-4485, fax: +81-75-753-4486.

⁴Unfortunately, the methodology of Sheliga et al. (2006c) was not an option in the present study because the monkey's OFR shows "anomalous contrast dependence" (Miles et al., 1986), involving effects on (and interactions between) latency and response amplitude that render the usual response measures inappropriate for examining contrast dependence closely (Miura et al., 2006).

least two different neural mechanisms by which primates analyze visual motion: one utilizes low-level motion detectors that are sensitive to first-order luminance modulation and the other (s) utilizes higher level mechanisms that are sensitive to features (Cavanagh and Mather, 1989). Recent findings strongly suggest that the initial OFRs of humans (Chen et al., 2005; Sheliga et al., 2005a, 2006a; Sheliga et al., 2006c) and monkeys (Miura et al., 2006) are mediated by the low-level mechanism. Much of the initial evidence for this came from studies using a special broadband stimulus, the so-called missing fundamental (*mf*), which indicated that the initial OFRs depended critically on the Fourier composition of the motion stimulus (Sheliga et al., 2005a; Miura et al., 2006; Sheliga et al., 2006c). The *mf* stimulus is a square wave that lacks the fundamental, so that in the frequency domain it consists of summed odd harmonics, the largest being the 3rd and the *i*th harmonic having an amplitude proportional to $1/i$. A special property of this stimulus, first recognized by Adelson (1982), is that when shifted $\frac{1}{4}$ of its wavelength, all of its harmonics each shift $\frac{1}{4}$ of their respective wavelengths, the $4n+1$ harmonics (where *n* is an integer) shifting in the direction of the actual image motion (i.e., forwards, along with the entire pattern and its features), whereas the $4n-1$ harmonics (which include the principal Fourier component, the 3rd harmonic) shift in the opposite direction (i.e., backwards).¹ The initial OFRs that were elicited when motion was applied to the *mf* stimulus in the form of successive $\frac{1}{4}$ -wavelength shifts of the pattern were always in the backward direction in both humans and monkeys, and this was attributed to the $\frac{1}{4}$ -wavelength (backward) shifts of the principal Fourier component (i.e., the 3rd harmonic).

This apparent dominance by the principal Fourier component suggested to Sheliga et al. (2006c) that there might be nonlinear interactions between the neural mechanisms sensing the conflicting motions of the various harmonics of this special broadband stimulus. In one test of this idea Sheliga et al. used moving stimuli with just two competing sine waves equivalent to the 3rd and 5th harmonics of the *mf* stimulus (the *3f5f* stimulus) that move in *opposite* directions, and found that the critical factor was the ratio of their two contrasts: when of similar contrast both were effective (vector sum/averaging), but when the contrast of one was less than about $\frac{1}{2}$ that of the other then the one with the higher contrast became dominant and the one with the lower contrast was rendered ineffective: winner-take-all. In another experiment, Sheliga et al. (2006c) restricted the moving stimuli to two competing sine waves equivalent to the 3rd and 7th harmonics of the *mf* stimulus (the *3f7f* stimulus) that move in the *same* direction (but at different speeds). Again, they found that when the contrast of one was less than about $\frac{1}{2}$ that of the other then the one with the higher contrast was dominant. Other studies on humans involving another short-latency oculomotor reflex, the disparity vergence response (DVR), indicated that it too depends on 1st-order energy mechanisms (Sheliga et al., 2006b) and shows nonlinear interactions, including winner-take-all behavior, when confronted with sine-wave gratings with competing binocular disparities that could have the same or opposite sign. (Sheliga et al., 2007) These nonlinear interactions in the OFR and DVR were attributed to mutual inhibition between the neural channels sensing the motion (or the disparity) of the two competing sine waves. Thus, the nonlinear interactions here were assumed to result entirely from the intrinsic properties of the sensory pathways: *bottom-up processes*.

Previous studies on the initial OFR (for review, see Miles, 1998) have suggested that the monkey provides a good animal model of the human, and the present study was undertaken to determine if the monkey's initial OFR shows evidence of nonlinear interactions when confronted with competing motion stimuli like those used in the study of Sheliga et al. (2006c) on humans. In Experiment 1 we used the *3f5f* stimulus (see Fig. 1A–C), whose two component sine waves moved in opposite directions when the entire pattern shifted in $\frac{1}{4}$ -

¹Of course, a $\frac{3}{4}$ -wavelength *forward* shift of a pure sine wave is exactly equivalent to a $\frac{1}{4}$ -wavelength *backward* shift (spatial aliasing), and it is the latter that determines the direction of both the perceived motion and any associated OFRs because the brain gives greatest weight to the nearest-neighbor matches (Sheliga et al., 2005a; Miura et al., 2006).

wavelength steps, and in Experiment 2 we used the $3f7f$ stimulus (see Fig. 1D–F), whose two component sine waves moved in the same direction when the pattern shifted in $\frac{1}{4}$ -wavelength steps. We report that the monkey's initial OFRs to these competing stimuli showed evidence of nonlinearities similar to those reported for humans.

2. Experiment 1: The initial OFR to the $3f5f$ stimulus and its dependence on the relative contrasts of the two components

In this experiment we used two superimposed vertical sine-wave gratings whose spatial frequencies were in the ratio 3:5 and recorded the initial OFRs that were elicited when the pattern underwent successive $\frac{1}{4}$ -wavelength shifts of the fundamental frequency. As shown in Fig. 1A–C, this meant that each of the components underwent successive $\frac{1}{4}$ -wavelength shifts of opposite sign. The dependent variable was the relative contrast of the two gratings and we report that when the contrast of one exceeded that of the other, on average, by a factor of 3.3 then the responses to the grating of lower contrast were almost totally suppressed.

2.1. Methods

Data were collected from three rhesus monkeys (*Macaca mulata*). All procedures were approved by the Animal Care and Use Committee of Kyoto University. Many of the general procedures were the same as those used in previous studies of ocular tracking in monkeys (Kawano et al., 1994; Kodaka et al., 2004; Miura et al., 2006) and humans (Sheliga et al., 2005a; Sheliga et al., 2006c).

2.1.1. Animal preparations—The monkeys were previously trained to fixate a small spot. Under pentobarbital sodium anesthesia and aseptic conditions, each monkey was implanted with a head holder, to allow the head to be fixed in the standard stereotaxic position during the experiments, and a scleral search coil to allow measurement of the position of the right eye (Judge et al., 1980).

2.1.2. Visual display and stimuli—The animals faced a 19 inch CRT monitor (Eizo T766, driven by a PC Radeon 9800 Pro video card), which was 50 cm in front of the eyes, in a dark room. Visual stimuli were presented on the monitor (resolution, 1280×1024 pixels; vertical refresh rate, 100 Hz). The RGB signals from the video card were converted to black and white images with 11-bit grayscale resolution through an attenuator (Pelli, 1997), exactly as described by Sheliga et al. (2005a). Briefly, a luminance look-up table with 256 equally-spaced luminance levels ranging from 0.3 cd/m^2 to 77.1 cd/m^2 was created by direct luminance measurements (LS-100 photometer; Konica-Minolta, Japan) under software control. This table was then expanded to 2048 equally-spaced levels by interpolation.

The visual stimuli used in our experiments were almost identical to those used by Sheliga et al. (2006c) and so will be described only in brief. The visual images consisted of one-dimensional vertical grating patterns that could have one of three horizontal luminance profiles in any given trial: 1) a sum of two sine waves with spatial frequencies in the ratio 3:5, creating a pattern with fundamental frequency, f (termed the “ $3f5f$ stimulus”), 2) a pure sine wave with a spatial frequency identical to that of the $3f$ component of the $3f5f$ stimulus (the “ $3f$ stimulus”), 3) a pure sine wave with a spatial frequency identical to that of the $5f$ component of the $3f5f$ stimulus (the “ $5f$ stimulus”). Each image extended 360 mm horizontally (39.6° ; 1280 pixels) and 270 mm vertically (30.2° ; 1024 pixels) and had a mean luminance of 38.7 cd/m^2 . The initial phase of a given grating was randomized from trial to trial at intervals of $\frac{1}{4}$ -wavelength. Motion was created by substituting a new image every frame (i.e., every 10 ms) for a total of 15 frames (i.e., stimulus duration, 150 ms), each new image being identical to the previous one except phase shifted horizontally by $\frac{1}{4}$ of the wavelength of the fundamental (whose spatial

frequency was f cycles/°). In any given trial, the successive steps were all in the same direction (rightward or leftward, randomly selected), so that the $3f$ stimulus/component moved *backwards* in steps that were each $\frac{1}{4}$ of its wavelength and the $5f$ stimulus/component moved *forwards* in steps that were each $\frac{1}{4}$ of its wavelength (see Fig. 1A–C). The dependent variable was the Michelson contrast, randomly sampled each trial from a lookup table with the following entries: the $5f$ stimulus had 3 entries (4, 8, 16%); the $3f$ stimulus had 21 entries (range, 1–64%); the $3f5f$ stimulus had 39 entries in which the $5f$ component could have 3 levels (4, 8, 16%) and the $3f$ component could have 13 levels (range, $\frac{1}{4}$ to 4 times the contrast of the $5f$ component). The spatial frequencies of the $3f$ and $5f$ stimuli were selected to be of roughly equal efficacy and optimal for each monkey: 0.30 and 0.50 cycles/°, respectively, for monkey A; 0.36 and 0.60 cycles/°, respectively, for monkey B; 0.45 and 0.75 cycles/°, respectively, for monkey C. The apparent speed of the $3f5f$ stimulus was 250 deg/s for monkey A, 208 deg/s for monkey B and 167 deg/s for monkey C, respectively.

2.1.3. Procedures—At the beginning of each trial, a grating pattern appeared together with a central target spot (diameter, 0.4°) that the animal had been trained to fixate. After the monkey's right eye had been positioned within 2° of the fixation target for a randomized period of 750 to 1000 ms (and no saccades had been detected for the last 250 ms in this period), the fixation target disappeared and the apparent-motion stimulus began. The motion lasted for 150 ms, after which time the screen became a uniform gray with the same mean luminance—signaling the end of the trial—and the animal was rewarded with a drop of juice. If there were any saccadic intrusions, the reward was withheld, the data were discarded and the trial was subsequently repeated. After an inter-trial interval of 1000–1500 ms, a new grating pattern appeared together with a fixation point, commencing a new trial. Data were collected in several sessions on consecutive days until each condition had been repeated an adequate number of times to permit good resolution of the responses through averaging.

2.1.4. Data collection and analyses—All aspects of the experimental paradigms were controlled by two PCs, which communicated via Ethernet using the TCP/IP protocol. One of the PCs was running a Real-time EXperimentation software package (REX) developed by Hays, Richmond and Optican (1982), and provided the overall control of the experimental protocol as well as acquiring, displaying, and storing the eye-movement data. The other PC was running MATLAB subroutines, utilizing the Psychophysics Toolbox extensions (Brainard, 1997; Pelli, 1997), and generated the visual stimuli upon receiving a start signal from the REX machine.

Eye movements were measured using the electromagnetic search coil technique (Fuchs and Robinson, 1966). The voltage signals encoding the horizontal and vertical components of the eye position were passed through an analog low-pass filter (-3dB at 200 Hz) and were digitized to a resolution of 12 bits, sampling at 1 kHz. All data were stored and transferred to another PC for analysis using computer programs based on MATLAB (The MathWorks Inc.). The eye-position data were smoothed with a 4-pole digital Butterworth filter (3 dB at 30 Hz), and eye velocity traces were derived from the two-point backward difference. Eye acceleration profiles were derived from the two-point backward difference of the eye velocity traces, and were used to detect small saccades that went undetected during the experiment. Trials with residual saccadic intrusions were then discarded.

Mean temporal eye velocity profiles (and standard deviations) were computed for each of the stimulus conditions. To obtain low-noise estimates of eye velocity, the responses to each stimulus were averaged over at least 36 trials. The eye velocity profiles over the 50-ms time periods starting 50 ms after the onset of the motion stimuli were used in the analyses. The minimum latency of the eye movements was ~ 50 ms so that these sections of the response profiles were restricted to the period prior to the closure of the visual feedback loop (i.e., twice

the reaction time): initial open-loop responses. The mean response profiles to rightward and leftward stimuli were pooled by subtracting the mean response profile to a given leftward motion stimulus from the mean response profile to the corresponding rightward motion stimulus, and these will be referred to the “R-L eye velocity profiles”. Because rightward eye movements were positive in our sign convention, these pooled R-L profiles were positive when the ocular following response was in the direction of the applied image shift (also referred to as the “forward” direction, in contradistinction to the “backward” or “reverse” direction).

2.2. Results

Figure 2 shows the R-L eye velocity profiles obtained with each of the three stimuli when the contrast of the $5f$ stimuli/components was 8% and the contrast of the $3f$ stimuli/components was 2, 4, 8, 16, or 32%; each column shows the data for a given monkey and each row the data for a given ($3f$) contrast. As expected, all responses to the pure $5f$ stimuli (blue dashed-dotted traces) were in the forward (positive) direction whereas all responses to the pure $3f$ stimuli (red dashed traces) were in the backward (negative) direction (cf., Miura et al., 2006). The responses to the $3f$ stimuli show dependence on the contrast, which can influence the latency, amplitude and waveform (cf., Miles et al., 1986). The responses to the $3f5f$ stimuli (grey traces) depended on the relative contrasts of the two components, but showed obvious nonlinearities. Thus, when the contrast of the $3f$ components was only 2% (i.e., one-fourth of the $5f$ component), the response profiles with the $3f5f$ stimuli were almost identical to those obtained with the pure $5f$ stimuli, even though pure $3f$ stimuli with a contrast of 2% always generated robust responses: see the top row of traces in Fig. 2. Clearly, the $5f$ components completely dominated the responses to the $3f5f$ stimuli here and the $3f$ components had no influence as though suppressed. The opposite situation prevailed when the $3f$ components of the $3f5f$ stimuli had a contrast of 32%, which was four times greater than the contrast of the $5f$ components: the response profiles obtained with the $3f5f$ stimuli were now almost identical to those obtained with the pure $3f$ stimuli, even though the pure $5f$ stimuli always generated robust responses: see the bottom row of traces in Fig. 2. Clearly, the $3f$ components now completely dominated the responses to the $3f5f$ stimuli and the $5f$ components had no influence as though suppressed. When the two components of the $3f5f$ stimuli were of the same (8%) contrast then responses were invariably very weak (see the middle row of traces in Fig. 2). When the $3f$ components of the $3f5f$ stimuli were of intermediate contrast, responses were more variable: in some cases, the component of higher contrast dominated (e.g., in monkeys A and B when the $3f$ component had a contrast of 4%) and in others both components contributed (e.g., in monkeys B and C when the $3f$ component had a contrast of 16%).

We attempted to reconstruct the R-L eye velocity profiles obtained with the $3f5f$ stimuli from the R-L eye velocity profiles obtained with the pure sine-wave stimuli using a *Contrast-Weighted-Average* (CWA) model, defined as follows:

$$R_{3f5f}(t) = w_{3f} \cdot R_{3f}(t) + w_{5f} \cdot R_{5f}(t) \quad (1)$$

in which the weights of the $3f$ and $5f$ components are given by:

$$w_{3f} = \frac{C_{3f}^{n_{3f}}}{C_{3f}^{n_{3f}} + C_{5f}^{n_{5f}}} \quad (2)$$

$$w_{5f} = 1 - w_{3f} \quad (3)$$

where $R_{3f5f}(t)$, $R_{3f}(t)$ and $R_{5f}(t)$ are the responses to the $3f5f$ stimuli, the pure $3f$ stimuli and the pure $5f$ stimuli, respectively. C_{3f} and C_{5f} are the contrast levels of the $3f$ and $5f$ components, respectively; the exponents, n_{3f} and n_{5f} , are the only free parameters and were determined by the optimization procedure. The least square fitting was performed for each level of $5f$ contrast independently and the optimal values of these two parameters were obtained (see Table 1).

This CWA model was the same as that used by Sheliga et al. (2006c) in their analyses of human OFRs. Figure 3, which has the same general layout as Fig. 2, shows the actual response profiles obtained with the $3f5f$ stimuli (gray traces, exactly as in Fig. 2) together with the profiles predicted from the CWA model (red dashed traces). We further show predictions from two other models: 1) a *sum* model (blue dashed-dotted traces), which assumed that the ocular responses to a given $3f5f$ stimulus were given by the sum of the responses to the component stimuli, and 2) a *simple average* model (green traces), which assumed that the responses to a given $3f5f$ stimulus were given by the average of the responses to the component stimuli.

Figure 3 suggests that, for all monkeys, the CWA model generally provided an excellent fit to the actual profiles over the full range of contrasts of the $3f$ component, whereas the sum and simple average models often did not. This general impression was confirmed by plotting the mean squared errors over the 50-ms time periods starting 50 ms after the onset of the motion stimuli for each of the model fits against the Contrast Ratio (C_{3f}/C_{5f}): see Fig. 4, which shows not only the errors for the data in Fig. 3, for which the $5f$ component had a contrast of 8% (dotted lines), but also the errors when the $5f$ component had a contrast of 4% (continuous lines) and 16% (dashed lines). It is now clear that the errors were lowest for all models when the Contrast Ratio was near 1 (actually, they were minimal when the contrast of the $3f$ component exceeded that of the $5f$ component very slightly) but only the CWA model consistently showed low error values over the full range of Contrast Ratios. The values of the n_{3f} and n_{5f} parameters for the least squares best-fit CWA model are listed in Table 1, together with the r^2 values indicating the goodness of fit. In fact, these r^2 values were always ≥ 0.924 (mean, 0.962), indicating that the CWA model always provided a very good description of the dependence of the $3f5f$ data on the Contrast Ratio. It is clear from Table 1 that, for a given stimulus with a given monkey, the values of n_{3f} and n_{5f} were generally quite similar, with overall means of 2.7 and 2.8, respectively, hence it is not surprising that when these two parameters were given the same value, n , there was little impact on the r^2 values (mean, 0.956 ± 0.022).

2.3. Discussion of Experiment 1

The data in Fig. 2 clearly indicate that the profiles obtained with the $3f5f$ stimuli depended critically on the relative contrasts of the two components and strongly suggest that there were major nonlinearities. For example, when there was a four-fold difference in the contrast of the two sine waves the OFR was completely determined by the one with the higher contrast and the one with the lower contrast lost all of its influence even though when the latter was presented alone it generated a robust response. In this situation, the sine wave with the higher contrast effectively suppressed the influence of the sine wave with the lower contrast in a winner-take-all fashion. Not surprisingly, the sum and simple average models failed to capture these features of the data but the CWA model provided an excellent fit, in line with the findings of Sheliga et al. (2006c) on the human OFR and of others working on other systems displaying similar nonlinear behavior (Krommenhoek and Wiegerinck, 1998; McGowan et al., 1998; Port and Wurtz, 2003; Walton et al., 2005; Sheliga et al., 2007). The sum and simple average are special cases of a generalized linear model given by, $R_{3f5f}(t) = k_1 \cdot R_{3f}(t) + k_2 \cdot R_{5f}(t)$, where k_1 and k_2 both have values of 1 and 0.5, respectively. Even when k_1 and k_2 were both free parameters, this linear model gave least square errors that were on average 3.04 (range, 1.50–5.87) times greater than for the CWA model. Additionally we attempted to reconstruct the R-L eye velocity profiles using a linear version of the CWA model, in which the weights of the $3f$ and $5f$ components are given by: $w_{3f} = A \cdot (C_{3f}/C_{5f}) + B$, $w_{5f} = 1 - w_{3f}$. A and B are both free parameters. The best-fit linear model gave least square errors that were on average 1.66 (range, 1.15–2.60) times greater than for the CWA model. The r^2 values for this linear model (mean: 0.913, SD: 0.022) were lower than the r^2 values for the CWA model in all cases. This supports the

conclusion that the profiles obtained with the $3f5f$ stimuli showed a nonlinear dependence on the relative contrast of the two components.

An important issue is the transition from dominance by one component to dominance by the other component as the Contrast Ratio is shifted from one extreme to the other. In particular, we were interested in the sharpness of this transition and how different the two contrasts had to be for one component to prevail completely over the other, factors important for assessing the likely biological significance of these nonlinear interactions. The CWA model allowed us to address these issues quantitatively. For this we used Equations 1–3 and the best-fit values of the two free parameters, n_{3f} and n_{5f} , to calculate the effective weights of the $5f$ components, given by w_{5f} , which we then plotted as a function of the Contrast Ratio (C_{3f}/C_{5f}) for each of the three data sets obtained from each of the three monkeys: see Fig. 5. The plots in Fig. 5 each have a sigmoidal shape with values ranging from zero (indicating that the $5f$ component of the $3f5f$ stimulus makes no contribution and OFRs are totally determined by the $3f$ component) to one (indicating that the $3f$ component of the $3f5f$ stimulus makes no contribution and OFRs are totally determined by the $5f$ component).² Note that the steepness of these sigmoids, which indicates the extent of the nonlinear interactions and, hence, the sharpness of the transition from dominance by one component to dominance by the other, is determined by the exponents, n_{3f} and n_{5f} , higher values indicating more powerful interactions and sharper transitions. Like Sheliga et al. (2006c), we examined these transitions further by calculating a Transition Zone, defined as the range of Contrast Ratios over which the values of w_{5f} ranged from 0.05 to 0.95: see the “5%” and “95%” listings in Table 1. On average, this Transition Zone extended from 2.76 to 0.48 in monkey A, 3.68 to 0.38 in monkey B, and 5.30 to 0.23 in monkey C. The means of the “5%” values and the reciprocals of the “95%” values for each of the three monkeys indicate that, on average, one component of the $3f5f$ stimulus effectively dominated the OFR when its contrast exceeded that of the other component by a factor of at least 2.4, 3.2, and 4.8, respectively.³ This indicates that there were substantial differences in the strengths of the nonlinear interactions in the different monkeys, those in monkey A being, on average, twice as powerful as those in monkey C.

When the two components of the $3f5f$ stimuli were of similar contrast then the OFRs were generally weak, perhaps suggesting that now neither component prevailed over the other and responses represented some sort of vector sum or average. However, in their study on the human OFR, Sheliga et al. (2006c) pointed out that the winner-take-all situation might also prevail inside the Transition Zone (as well as outside), in which case the response distributions with the $3f5f$ stimuli would be determined probabilistically. For example, a value of 0.6 for w_{5f} in Fig. 5 could mean that the OFR was effectively driven exclusively by the $5f$ component in some proportion of the trials and exclusively by the $3f$ component in the remainder of the trials. In this event, we might expect the distributions of the OFRs to the $3f5f$ stimuli on individual trials to be much broader—perhaps even bimodal in some cases—inside the Transition Zone than outside (where only one component was effective in most trials). We examined this issue quantitatively by simulating the response distributions predicted by the winner-take-all/probabilistic model when the Contrast Ratios were 0.5, using the distributions of the measured responses to the pure $3f$ and $5f$ stimuli. An example based on data from monkey A is shown in Fig. 6. The histograms in Fig. 6A show the distributions of the OFR measures on individual trials using the following stimuli: 1) a leftward pure $3f$ stimulus of contrast 8% (orange plot), 2) a rightward pure $5f$ stimulus of contrast 8% (green plot), and 3) a $3f5f$ stimulus whose $3f$ and $5f$ components had directions and contrasts matching those of the corresponding sine-wave

²The curves in Fig. 5 are equivalent to the Response Ratios plotted in Fig. 4 of Sheliga et al (2006c), which had very similar sigmoidal forms.

³Note that we examined only the special case in which the two sine waves were of similar efficacy when of equal contrast and applied singly.

stimuli (blue plot). The best-fit Gaussians for the three distributions are also shown in continuous line and provide a good description of the data (r^2 values were 0.973, 0.931, and 0.986, respectively). We then calculated the Response Ratio of Sheliga et al. (2006c) from the mean OFR measures obtained with each of the 3 stimuli using the following expression:

$$\frac{R_{3f5f} - R_{3f}}{R_{5f} - R_{3f}} \quad (4)$$

where R_{3f5f} is the mean response to the $3f5f$ stimulus when the $3f$ and $5f$ components have particular contrast values, and R_{3f} and R_{5f} are the mean responses to the pure $3f$ and $5f$ stimuli with contrasts matching those values. The Response Ratio has a value of one when the $3f5f$ responses are determined exclusively by the $5f$ component and a value of zero when those responses are determined exclusively by the $3f$ component. For the data in Fig. 6A, the Response Ratio was 0.61. We then simulated the distribution of the responses to the $3f5f$ stimuli predicted by the winner-take-all/probabilistic model by summing the response distributions obtained with the pure $3f$ and $5f$ stimuli, weighted in accordance with this Response Ratio: see the black/grey histogram in Fig. 6B labeled, “simulated $3f5f$ ”. The simulated response distribution in Fig. 6B is clearly bimodal and much broader than the actual response distribution, which is repeated in Fig. 6B (in blue) to facilitate the comparison. This indicates that the responses were *not* probabilistically determined. Similar findings were obtained from the other two monkeys. In fact, based on the best-fit Gaussian functions, the real $3f5f$ data (mean $r^2 = 0.949$) were always much more narrowly distributed than the simulated $3f5f$ data (mean $r^2 = 0.862$) with mean SDs ($n = 18$) of 0.018 and 0.033, respectively. Note that the CWA model also predicts a normal distribution for the $3f5f$ responses with a mean given by $w_{3f} \cdot R_{3f} + w_{5f} \cdot R_{5f}$ and a variance given by $w_{3f}^2 \cdot V_{3f} + w_{5f}^2 \cdot V_{5f}$, where w_{3f} , R_{3f} and V_{3f} are the weight, mean and variance of the $3f$ responses, respectively, and w_{5f} , R_{5f} and V_{5f} are the same entities for the $5f$ responses. For the data in Fig. 6, the $3f5f$ responses predicted by this CWA model had a mean $\pm$ SD of -0.0031 ± 0.0122 (not shown in Fig. 6). The differences between the distributions of the real and the predicted responses were not significant at the 0.05 level on the Kolmogorov-Smirnov two-sample test.

Previous authors who reported winner-take-all oculomotor responses (Ferrera and Lisberger, 1995, 1997; Recanzone and Wurtz, 1999; Ferrera, 2000; Sheliga et al., 2006c; Sheliga et al., 2007) invoked nonlinear interactions in the form of mutual inhibition between motion-sensitive channels that are selectively sensitive to opposite directions of motion. This is often referred to as, “motion opponency”, and has supporting evidence from psychophysical studies (Levinson and Sekuler, 1975; Mather and Moulden, 1983; Stromeyer et al., 1984; van Santen and Sperling, 1984; Qian et al., 1994; Zemany et al., 1998), functional magnetic resonance imaging (Heeger et al., 1999), and single unit recordings in areas V1 and MT (Mikami et al., 1986; Rodman and Albright, 1987; Snowden et al., 1991; Qian and Andersen, 1994; Bradley et al., 1995; Rust, 2004; Rust et al., 2005). In our data, the exponents, n_{3f} and n_{5f} , provide an estimate of the strengths of the mutual inhibition between the neural channels mediating the competing motions. In 7 of 9 cases, $n_{3f} < n_{5f}$, indicating that the channels mediating the $5f$ component of the $3f5f$ stimulus usually exerted a slightly greater inhibitory effect on the channels mediating the $3f$ component than vice versa. This result is consistent with the result of Sheliga et al (2006c).

3. Experiment 2: The initial OFR to the $3f7f$ stimulus and its dependence on the relative contrasts of the two components

In Experiment 1 we used two superimposed sine-wave gratings whose spatial frequencies were in the ratio 3:5 and we recorded the initial OFRs that were elicited when the two gratings were subject to successive $\frac{1}{4}$ -wavelength shifts of *opposite* sign. In the present Experiment we combined two gratings whose spatial frequencies were in the ratio 3:7 and whose $\frac{1}{4}$ -wavelength

shifts had the *same* sign. We again report that when the contrast of one component exceeded that of the other by a certain amount then the component with the lesser contrast lost its influence on the initial OFR as though suppressed: winner-take-all.

3.1. Methods

Most of the methods and procedures were identical to those used in Experiment 1, and only those that were different will be described here.

3.1.1. Visual display and stimuli—The visual images consisted of one-dimensional vertical grating patterns that could have one of three horizontal luminance profiles in any given trial: 1) a sum of two sine waves with spatial frequencies in the ratio 3:7, creating a pattern with a fundamental spatial frequency, f (the “ $3f7f$ stimulus”); 2) a pure sine wave with a spatial frequency identical to that of the $3f$ component of the $3f7f$ stimulus (the “ $3f$ stimulus”), 3) a pure sine wave with a spatial frequency identical to that of the $7f$ component of the $3f7f$ stimulus (the “ $7f$ stimulus”). Motion was created by substituting a new image every frame, each new image being identical to the previous one except phase shifted horizontally by $\frac{1}{4}$ of the wavelength of the fundamental (whose spatial frequency was f cycle/°). In any given trial, the successive steps were all in the same direction (rightward or leftward, randomly selected) and both the $3f$ and the $7f$ stimuli/components moved *backwards* in steps that were each $\frac{1}{4}$ of their respective wavelengths (see Fig. 1D–F). The dependent variable was the Michelson contrast, randomly sampled each trial from a lookup table with the following entries: the $7f$ stimulus had 3 entries (4, 8, 16%); the $3f$ stimulus had 21 entries (range, 1–64%); the $3f7f$ stimulus had 39 entries in which the $7f$ component could have 3 levels (4, 8, 16%) and the $3f$ component could have 13 levels (range, $\frac{1}{4}$ to 4 times the contrast of the $7f$ component). The spatial frequencies of the $3f$ and $7f$ stimuli were selected to be of roughly equal efficacy and optimal for each monkey: 0.30 and 0.70 cycles/°, respectively, for monkey A; 0.36 and 0.84 cycles/°, respectively, for monkey B; 0.45 and 1.05 cycles/°, respectively, for monkey C. Then, the apparent speed of the $3f7f$ stimulus was 250 deg/s for monkey A, 208 deg/s for monkey B and 167 deg/s for monkey C, respectively.

3.2. Results

Figure 7, which has a general layout similar to that of Fig. 2, shows the R-L eye velocity profiles obtained with each of the three stimuli when the contrast of the $7f$ stimuli/components was always 16% and the contrast of the $3f$ stimuli/components was 4, 8, 16, 32, or 64%. As expected, both of the responses to the pure $7f$ stimuli and the $3f$ stimuli in Fig. 7 are in the backward (negative) direction. The responses to the $3f$ stimuli (red dashed traces) show a strong dependence on contrast, which can affect the latency, amplitude and waveform (cf., data in Fig. 2). The responses to the $3f7f$ stimuli (grey traces) clearly depended on the relative contrasts of the two components, and showed evidence of nonlinearities. Thus, when the contrast of the $3f$ components was only 4%, (i.e., one-fourth of the $7f$ component), the response profiles obtained with the $3f7f$ stimuli were almost identical to those obtained with the pure $7f$ stimuli, even though pure $3f$ stimuli with a contrast of 4% generated robust responses that always deviated from those to the pure $7f$ stimuli at some point in their profile: see the top row of traces in Fig. 7. Clearly, the $7f$ components completely dominated the responses to the $3f7f$ stimuli here and the $3f$ components had little or no influence as though suppressed. The opposite situation prevailed when the $3f$ components of the $3f7f$ stimuli had a contrast of 64%, which was four times the contrast of the $7f$ components: the response profiles obtained with the $3f7f$ stimuli were now almost identical to those obtained with the pure $3f$ stimuli, even though the pure $7f$ stimuli always generated responses with very different profiles: see the bottom row of traces in Fig. 7. Clearly, the $3f$ components now completely dominated the responses to the $3f7f$ stimuli and the $7f$ components had no influence as though suppressed. When the $3f$ components of the $3f7f$ stimuli were of intermediate contrast, responses were more variable:

in some cases, the component of higher contrast dominated (e.g., in all monkeys when the 3f components had a contrast of 32%) and in others both components contributed (e.g., in monkeys A and C when the 3f components had a contrast of 16%).

Figure 8, which has the same general layout as Fig. 7, shows the actual response profiles obtained with the 3f7f stimuli (grey traces, exactly as in Fig. 7) together with the best-fit simulations using the same three models as were used in Experiment 1: the CWA (red dashed traces), the sum (blue dashed-dotted traces), and the simple average (green traces). It is apparent from Fig. 8 that, for all monkeys, the CWA model generally provided a good fit to the actual profiles over the full range of contrasts of the 3f component, whereas the sum and simple average often deviated substantially. That the CWA model provided the best fit is evident from Fig. 9, which shows the mean squared errors over the 50-ms time periods starting 50 ms after the onset of the motion stimuli for each of the three model fits plotted as a function of the Contrast Ratio (C_{3f}/C_{7f}) for each of the three monkeys (cf., Fig. 4). Thus, the CWA model (red lines in Fig. 9) consistently showed lower error values than the sum (blue) and simple average (green) models over the full range of Contrast Ratios, regardless of whether the 7f component had a contrast of 4% (continuous lines), 8% (dotted lines), or 16% (dashed lines). The r^2 values for the least squares best-fit CWA model were always ≥ 0.896 (mean, 0.934), indicating the excellence of these fits, but many of the exponents were much higher than in Experiment 1 (mean $n_{3f} = 42.8$, mean $n_{7f} = 33.8$) and reflect a severe limitation of Experiment 2: the differences in the responses to the pure 3f and 7f stimuli were often very subtle, sometimes even when the two sine waves had very different contrasts. The net result was that the free parameters in the CWA fits (n_{3f} and n_{7f}) were very sensitive to noise when the difference between the responses to the 3f and 7f stimuli was small. When we excluded data for which the difference in the measured responses to the pure 3f and 7f stimuli was less than 0.035° (cf., Sheliga et al., 2006c), only 3/9 CWA fits remained, with mean n_{3f} and n_{7f} values of 3.13 and 2.93, respectively: see Table 2 for the parameters.

3.3. Discussion of Experiment 2

The data in Fig. 7 clearly indicate that the profiles obtained with the 3f7f stimuli depended critically on the relative contrasts of the two components and again suggest that there were major nonlinearities. For example, when the 3f components had four times the contrast of the 7f component (bottom row in Fig. 7) the OFR profiles obtained with the 3f7f stimuli closely followed those obtained with the 3f stimuli even though the 7f stimuli alone generated very different response profiles. In this situation, the sine waves with the higher contrast effectively suppressed the influence of the sine waves with the lower contrast in a winner-take-all fashion. As in Experiment 1, the sum and simple average models often failed to capture the main features of the data but the CWA model always provided an excellent fit, again in line with the findings of Sheliga et al. (2006c) using the 3f7f stimuli with humans. We again wanted to use the CWA model and the best-fit values of its two free parameters to characterize the shift in dominance from one component to the other as the Contrast Ratio was changed but, unfortunately, the response profiles obtained with the 3f and 7f stimuli were often too similar (see Results) so only 3/9 data sets could be used to examine this transition. The three resulting plots in Fig. 10 have sigmoidal shapes similar to those in Fig. 5 and we again defined a Transition Zone: see the “5%” and “95%” listings in Table 2. Based on these limited data, the average Transition Zone extended from 2.38 to 0.34 so that one component of the 3f7f stimulus effectively dominated the OFR when its contrast exceeded that of the other component by a factor of at least 2.7, which is within the range of values reported in Experiment 1.

Again we attempted to reconstruct in three cases above the R-L eye velocity profiles using the linear model, in which the weights of the 3f and 7f components are given by: $w_{3f} = A \cdot (C_{3f}/C_{7f}) + B$, $w_{7f} = 1 - w_{3f}$. A and B are both free parameters. The best-fit linear model gave least

square errors that were on average 1.08 (range, 1.02–1.16) times greater than for the CWA model. The r^2 values for this linear model (mean: 0.943, SD: 0.022) were a little lower than the r^2 values for the CWA model in all cases. This result shows that, within the Transition Zone, the linear model is comparable to the CWA model with the $3f7f$ stimulus.

We again invoke mutual inhibition between the neural mechanisms sensing the two motions to explain the nonlinear interactions between the two components of the $3f7f$ stimuli. However, here the two gratings are moving in the same direction so that the brain must rely on the differences in their spatial frequencies and/or speeds in order to distinguish between them. There is evidence for mutual inhibition between neurons in striate cortex tuned for different spatial frequencies (e.g., Movshon et al., 1978; Albrecht and De Valois, 1981; De Valois and Tootell, 1983), though these studies were not specifically concerned with motion-selective neurons. The exponents, n_{3f} and n_{7f} , provide an estimate of the strengths of the mutual inhibition between the neural channels mediating the competing motions. In 2 of 3 cases, $n_{3f} > n_{7f}$, indicating that the channels mediating the $3f$ component of the $3f7f$ stimulus in those cases exerted a slightly greater inhibitory effect on the channels mediating the $7f$ component than vice versa.

4. General Discussion

4.1. Is the monkey a good animal model?

Using competing motion stimuli, the present study on the OFR of the monkey uncovered nonlinear interactions that were generally similar to those recently reported for the human OFR using similar stimuli (Sheliga et al., 2006c). Perhaps not surprisingly there were some quantitative differences, especially in the apparent strength of the nonlinear interactions as indicated by the exponents of the best-fit CWA functions. In Experiment 1, the mean values of n_{3f} and n_{5f} were 2.7 and 2.8, respectively, values that are approximately half those obtained by Sheliga et al (2006c) in analogous experiments on humans (5.5 and 5.7, respectively). Thus, based on these estimates, the nonlinear interactions were less powerful in the monkey than in the human. However, the method of quantification was slightly different in the study of Sheliga et al. (2006c), which used the changes in horizontal eye position over the initial open-loop interval rather than the form of the response profiles as in our present study.⁴³ In Experiment 2, the database available for estimating the strength of the nonlinear interactions was small (33% of the total), as too was the database obtained with the $3f7f$ stimuli in the study of Sheliga et al. (2006c) and a direct quantitative comparison with the human data was not attempted. Although there might be some quantitative differences between the two species, to the extent that the monkey shows clear evidence of nonlinear interactions with both the $3f5f$ and $3f7f$ stimuli it provides a very promising animal model for studying the initial OFRs to competing motion stimuli. Furthermore, there is considerable evidence concerning the neural mediation of the OFR in monkeys: The MST region of cortex is known to play an important role (Kawano et al., 1994; Inoue et al., 1998; Kawano et al., 2000; Takemura et al., 2000, 2002; Takemura et al., 2007), together with its brainstem projections through the dorsolateral pontine nuclei (Kawano et al., 1992) to the ventral paraflocculus (Miles et al., 1986; Shidara and Kawano, 1993; Shidara et al., 1993; Kawano and Shidara, 1994; Kawano et al., 1996a; Kawano et al., 1996b; Gomi et al., 1998; Takemura et al., 2001a; Takemura and Kawano, 2002). In addition, Rust (2004) used competing motions similar to those in Experiment 1 and reported similar nonlinear interactions in the associated neuronal responses in cortical areas V1 and MT. This is particularly interesting because MT is a major source of inputs to MST (Maunsell and van Essen, 1983; Ungerleider and Desimone, 1986).

4.2. Functional significance of nonlinear interactions?

The previous studies on the human OFR (Sheliga et al., 2006c) and DVR (Sheliga et al., 2007) that used visual stimuli with competing harmonics and reported a highly nonlinear dependence on their relative contrasts argued that this feature would endow these ocular reflexes with a preference for objects in the plane of fixation: because of accommodation, the retinal images of objects in the plane of fixation tend to be better focused—and hence tend to have higher contrasts—than those of objects in other depth planes. It was pointed out that this would be in line with earlier studies on monkeys and humans, which showed that when broadband random-dot stimuli were used, the OFR was effectively disabled by—and the DVR was unresponsive to—binocular disparities of more than a few degrees (Busetini et al., 1996; Takemura et al., 2000; Busettini et al., 2001; Masson et al., 2001; Takemura et al., 2001b; Yang et al., 2003; Yang and Miles, 2003), indicating that, in everyday conditions, these reflexes will have a strong preference for objects in the immediate vicinity of the plane of fixation and will tend to ignore objects in other depth planes. Of course, the more powerful the nonlinear interactions then the sharper the distinctions between the competing stimuli. However, if the nonlinear interactions are too powerful then the system will be overly sensitive to noise (perhaps even unstable). Presumably, the actual strength of the interactions is a compromise between the need to discriminate between stimuli and the need for noise immunity.

Acknowledgements

This work was supported by Grant-in-Aid for JSPS Fellows (#18-3259), the JSPS.KAKENHI (#16GS0312), the MEXT.KAKENHI (#17700309), the Japan-US Brain Research Cooperative Program, and the Intramural Research Program of the National Eye Institute at the NIH.

References

- Adelson EH. Some new motion illusions, and some old ones, analysed in terms of their Fourier components. *Invest Ophthalmol Vis Sci* 1982;34(Suppl):144.(Abstract)
- Albrecht DG, De Valois RL. Striate cortex responses to periodic patterns with and without the fundamental harmonics. *J Physiol* 1981;319:497–514. [PubMed: 7320923]
- Bradley DC, Qian N, Andersen RA. Integration of motion and stereopsis in middle temporal cortical area of macaques. *Nature* 1995;373:609–611. [PubMed: 7854417]
- Brainard DH. The Psychophysics Toolbox. *Spat Vis* 1997;10:433–436. [PubMed: 9176952]
- Busettini C, Fitzgibbon EJ, Miles FA. Short-latency disparity vergence in humans. *J Neurophysiol* 2001;85:1129–1152. [PubMed: 11247983]
- Busettini C, Miles FA, Krauzlis RJ. Short-latency disparity vergence responses and their dependence on a prior saccadic eye movement. *J Neurophysiol* 1996;75:1392–1410. [PubMed: 8727386]
- Cavanagh P, Mather G. Motion: the long and short of it. *Spat Vis* 1989;4:103–129. [PubMed: 2487159]
- Chen KJ, Sheliga BM, Fitzgibbon EJ, Miles FA. Initial ocular following in humans depends critically on the fourier components of the motion stimulus. *Ann NY Acad Sci* 2005;1039:260–271. [PubMed: 15826980]
- De Valois KK, Tootell RB. Spatial-frequency-specific inhibition in cat striate cortex cells. *J Physiol* 1983;336:359–376. [PubMed: 6875912]
- Ferrera VP. Task-dependent modulation of the sensorimotor transformation for smooth pursuit eye movements. *J Neurophysiol* 2000;84:2725–2738. [PubMed: 11110803]
- Ferrera VP, Lisberger SG. Attention and target selection for smooth pursuit eye movements. *J Neurosci* 1995;15:7472–7484. [PubMed: 7472499]
- Ferrera VP, Lisberger SG. The effect of a moving distractor on the initiation of smooth-pursuit eye movements. *Vis Neurosci* 1997;14:323–338. [PubMed: 9147484]
- Fuchs AF, Robinson DA. A method for measuring horizontal and vertical eye movement chronically in the monkey. *J Appl Physiol* 1966;21:1068–1070. [PubMed: 4958032]
- Gellman RS, Carl JR, Miles FA. Short latency ocular-following responses in man. *Vis Neurosci* 1990;5:107–122. [PubMed: 2278939]

- Gomi H, Shidara M, Takemura A, Inoue Y, Kawano K, Kawato M. Temporal firing patterns of Purkinje cells in the cerebellar ventral paraflocculus during ocular following responses in monkeys I. Simple spikes *J Neurophysiol* 1998;80:818–831.
- Hays AV, Richmond BJ, Optican LM. A UNIX-based multiple process system for real-time data acquisition and control. *WESCON Conf Proc* 1982;2:1–10.
- Heeger DJ, Boynton GM, Demb JB, Seidemann E, Newsome WT. Motion opponency in visual cortex. *J Neurosci* 1999;19:7162–7174. [PubMed: 10436069]
- Inoue Y, Takemura A, Kawano K, Kitama T, Miles FA. Dependence of short-latency ocular following and associated activity in the medial superior temporal area (MST) on ocular vergence. *Exp Brain Res* 1998;121:135–144. [PubMed: 9696382]
- Judge SJ, Richmond BJ, Chu FC. Implantation of magnetic search coils for measurement of eye position: an improved method. *Vision Res* 1980;20:535–538. [PubMed: 6776685]
- Kawano K, Inoue Y, Takemura A, Kodaka Y, Miles FA. The role of MST neurons during ocular tracking in 3D space. *Int Rev Neurobiol* 2000;44:49–63. [PubMed: 10605641]
- Kawano K, Shidara M. Information representation by Purkinje cells in the cerebellum during ocular following responses. *Neurosci Res* 1994;21:13–17. [PubMed: 7708291]
- Kawano K, Shidara M, Takemura A, Inoue Y, Gomi H, Kawato M. Inverse-dynamics representation of eye movements by cerebellar Purkinje cell activity during short-latency ocular-following responses. *Ann NY Acad Sci* 1996a;781:314–321. [PubMed: 8694423]
- Kawano K, Shidara M, Watanabe Y, Yamane S. Neural activity in cortical area MST of alert monkey during ocular following responses. *J Neurophysiol* 1994;71:2305–2324. [PubMed: 7931519]
- Kawano K, Shidara M, Yamane S. Neural activity in dorsolateral pontine nucleus of alert monkey during ocular following responses. *J Neurophysiol* 1992;67:680–703. [PubMed: 1578251]
- Kawano K, Takemura A, Inoue Y, Kitama T, Kobayashi Y, Mustari MJ. Visual inputs to cerebellar ventral paraflocculus during ocular following responses. *Prog Brain Res* 1996b;112:415–422. [PubMed: 8979846]
- Kodaka Y, Miura K, Suehiro K, Takemura A, Kawano K. Ocular tracking of moving targets: effects of perturbing the background. *J Neurophysiol* 2004;91:2474–2483. [PubMed: 14762158]
- Krommenhoek KP, Wiegerinck WA. A neural network study of precollicular saccadic averaging. *Biol Cybern* 1998;78:465–477. [PubMed: 9711820]
- Levinson E, Sekuler R. Inhibition and disinhibition of direction-specific mechanisms in human vision. *Nature* 1975;254:692–694. [PubMed: 1124126]
- Masson GS, Busetini C, Yang DS, Miles FA. Short-latency ocular following in humans: sensitivity to binocular disparity. *Vision Res* 2001;41:3371–3387. [PubMed: 11718780]
- Mather G, Moulden B. Thresholds for movement direction: two directions are less detectable than one. *Q J Exp Psychol A* 1983;35:513–518. [PubMed: 6571322]
- Maunsell JH, van Essen DC. The connections of the middle temporal visual area (MT) and their relationship to a cortical hierarchy in the macaque monkey. *J Neurosci* 1983;3:2563–2586. [PubMed: 6655500]
- McGowan JW, Kowler E, Sharma A, Chubb C. Saccadic localization of random dot targets. *Vision Res* 1998;38:895–909. [PubMed: 9624439]
- Mikami A, Newsome WT, Wurtz RH. Motion selectivity in macaque visual cortex. I Mechanisms of direction and speed selectivity in extrastriate area MT. *J Neurophysiol* 1986;55:1308–1327. [PubMed: 3016210]
- Miles FA. The neural processing of 3-D visual information: evidence from eye movements. *Eur J Neurosci* 1998;10:811–822. [PubMed: 9753150]
- Miles FA, Kawano K, Optican LM. Short-latency ocular following responses of monkey. I Dependence on temporospatial properties of visual input. *J Neurophysiol* 1986;56:1321–1354. [PubMed: 3794772]
- Miura K, Matsuura K, Taki M, Tabata H, Inaba N, Kawano K, Miles FA. The visual motion detectors underlying ocular following responses in monkeys. *Vision Res* 2006;46:869–878. [PubMed: 16356529]

- Movshon JA, Thompson ID, Tolhurst DJ. Spatial and temporal contrast sensitivity of neurones in areas 17 and 18 of the cat's visual cortex. *J Physiol* 1978;283:101–120. [PubMed: 722570]
- Pelli DG. The VideoToolbox software for visual psychophysics: transforming numbers into movies. *Spat Vis* 1997;10:437–442. [PubMed: 9176953]
- Port NL, Wurtz RH. Sequential activity of simultaneously recorded neurons in the superior colliculus during curved saccades. *J Neurophysiol* 2003;90:1887–1903. [PubMed: 12966180]
- Qian N, Andersen RA. Transparent motion perception as detection of unbalanced motion signals. II. Physiology. *J Neurosci* 1994;14:7367–7380. [PubMed: 7996182]
- Qian N, Andersen RA, Adelson EH. Transparent motion perception as detection of unbalanced motion signals. I. Psychophysics. *J Neurosci* 1994;14:7357–7366. [PubMed: 7996181]
- Recanzone GH, Wurtz RH. Shift in smooth pursuit initiation and MT and MST neuronal activity under different stimulus conditions. *J Neurophysiol* 1999;82:1710–1727. [PubMed: 10515961]
- Rodman HR, Albright TD. Coding of visual stimulus velocity in area MT of the macaque. *Vision Res* 1987;27:2035–2048. [PubMed: 3447355]
- Rust, NC. Center for Neural Science. New York: New York University; 2004. Signal transmission, feature representation and computation in areas V1 and MT of the macaque monkey; p. 165
- Rust NC, Schwartz O, Movshon JA, Simoncelli EP. Spatiotemporal elements of macaque v1 receptive fields. *Neuron* 2005;46:945–956. [PubMed: 15953422]
- Sheliga BM, Chen KJ, Fitzgibbon EJ, Miles FA. Initial ocular following in humans: a response to first-order motion energy. *Vision Res* 2005a;45:3307–3321. [PubMed: 15894346]
- Sheliga BM, Chen KJ, Fitzgibbon EJ, Miles FA. Short-latency disparity vergence in humans: evidence for early spatial filtering. *Ann NY Acad Sci* 2005b;1039:252–259. [PubMed: 15826979]
- Sheliga BM, Chen KJ, FitzGibbon EJ, Miles FA. The initial ocular following responses elicited by apparent-motion stimuli: reversal by inter-stimulus intervals. *Vision Res* 2006a;46:979–992. [PubMed: 16242168]
- Sheliga BM, FitzGibbon EJ, Miles FA. Short-latency disparity vergence eye movements: a response to disparity energy. *Vision Res* 2006b;46:3723–3740. [PubMed: 16765403]
- Sheliga BM, FitzGibbon EJ, Miles FA. Human vergence eye movements initiated by competing disparities: evidence for a winner-take-all mechanism. *Vision Res* 2007;47:479–500. [PubMed: 17118422]
- Sheliga BM, Kodaka Y, FitzGibbon EJ, Miles FA. Human ocular following initiated by competing image motions: evidence for a winner-take-all mechanism. *Vision Res* 2006c;46:2041–2060. [PubMed: 16487988]
- Shidara M, Kawano K. Role of Purkinje cells in the ventral paraflocculus in short-latency ocular following responses. *Exp Brain Res* 1993;93:185–195. [PubMed: 8491260]
- Shidara M, Kawano K, Gomi H, Kawato M. Inverse-dynamics model eye movement control by Purkinje cells in the cerebellum. *Nature* 1993;365:50–52. [PubMed: 8361536]
- Snowden RJ, Treue S, Erickson RG, Andersen RA. The response of area MT and V1 neurons to transparent motion. *J Neurosci* 1991;11:2768–2785. [PubMed: 1880548]
- Stromeyer CF 3rd, Kronauer RE, Madsen JC, Klein SA. Opponent-movement mechanisms in human vision. *J Opt Soc Am A* 1984;1:876–884. [PubMed: 6470838]
- Takemura A, Inoue Y, Gomi H, Kawato M, Kawano K. Change in neuronal firing patterns in the process of motor command generation for the ocular following response. *J Neurophysiol* 2001a;86:1750–1763. [PubMed: 11600636]
- Takemura A, Inoue Y, Kawano K. The effect of disparity on the very earliest ocular following responses and the initial neuronal activity in monkey cortical area MST. *Neurosci Res* 2000;38:93–101. [PubMed: 10997582]
- Takemura A, Inoue Y, Kawano K. Visually driven eye movements elicited at ultra-short latency are severely impaired by MST lesions. *Ann NY Acad Sci* 2002;956:456–459. [PubMed: 11960839]
- Takemura A, Inoue Y, Kawano K, Quaia C, Miles FA. Single-unit activity in cortical area MST associated with disparity-vergence eye movements: evidence for population coding. *J Neurophysiol* 2001b;85:2245–2266. [PubMed: 11353039]

- Takemura A, Kawano K. Sensory-to-motor processing of the ocular-following response. *Neurosci Res* 2002;43:201–206. [PubMed: 12103438]
- Takemura A, Murata Y, Kawano K, Miles FA. Deficits in short-latency tracking eye movements after chemical lesions in monkey cortical areas MT and MST. *J Neurosci* 2007;27:529–541. [PubMed: 17234585]
- Ungerleider LG, Desimone R. Cortical connections of visual area MT in the macaque. *J Comp Neurol* 1986;248:190–222. [PubMed: 3722458]
- van Santen JP, Sperling G. Temporal covariance model of human motion perception. *J Opt Soc Am A* 1984;1:451–473. [PubMed: 6726493]
- Walton MM, Sparks DL, Gandhi NJ. Simulations of saccade curvature by models that place superior colliculus upstream from the local feedback loop. *J Neurophysiol* 2005;93:2354–2358. [PubMed: 15615826]
- Yang DS, FitzGibbon EJ, Miles FA. Short-latency disparity-vergence eye movements in humans: sensitivity to simulated orthogonal tropias. *Vision Res* 2003;43:431–443. [PubMed: 12536000]
- Yang DS, Miles FA. Short-latency ocular following in humans is dependent on absolute (rather than relative) binocular disparity. *Vision Res* 2003;43:1387–1396. [PubMed: 12742108]
- Zemany L, Stromeyer CF 3rd, Chaparro A, Kronauer RE. Motion detection on flashed, stationary pedestal gratings: evidence for an opponent-motion mechanism. *Vision Res* 1998;38:795–812. [PubMed: 9624431]

Fig. 1.

The $3f5f$ and $3f7f$ stimuli. Traces show luminance as a function of horizontal spatial position when the stimuli undergo successive $1/4$ -wavelength rightward shifts. (A–C) The $3f5f$ stimulus; the phase of the $3f$ and $5f$ components is the same; black open circles and associated arrows in A (grey circles and arrows in B, C) indicate the rightward shifts of one particular trough in the $3f5f$ stimulus profile; black open circles and associated arrows in B and C indicate the $1/4$ -wavelength leftward shifts of the $3f$ component and the $1/4$ -wavelength rightward shifts of the $5f$ component, respectively; the black horizontal lines terminated by arrow heads in A indicate one wavelength of the $3f5f$ stimulus (i.e., the fundamental) and these are repeated in grey in B and C; the black horizontal lines terminated by arrow heads in B and C indicate one wavelength of the $3f$ and $5f$ components, respectively. (D–F) The $3f7f$ stimulus; the phase of the $3f$ and $7f$ components is the same; black open circles and associated arrows in D (grey circles and arrows in E, F) indicate the rightward shifts of one particular peak in the $3f7f$ stimulus profile; black open circles and associated arrows in E and F indicate the $1/4$ -wavelength leftward shifts of the $3f$ and $7f$ components, respectively; the black horizontal lines terminated by arrow heads in D indicate one wavelength of the $3f7f$ stimulus (i.e., the fundamental) and these are repeated in grey in E and F; the black horizontal lines terminated by arrow heads in E and F indicate one wavelength of the $3f$ and $7f$ components, respectively.

Fig. 2.

The initial OFRs to the $3f5f$ stimuli: dependence on the contrast of the $3f$ component when the $5f$ component always had a contrast of 8% (R-L eye velocity profiles over time). Each column shows data for one monkey and each row shows data for $3f$ stimuli/components of one contrast (see labels near traces in left column). Data were obtained with $3f$ stimuli (red dashed traces), $5f$ stimuli (blue dashed-dotted traces), and $3f5f$ stimuli (grey traces). The responses of a given monkey to the pure $5f$ stimuli are all identical. Eye movements in the direction of pattern shift are positive and result in upward deflections. Horizontal dashed lines, zero eye velocity. Vertical calibration bars, 4°/s. Time zero denotes the time of the first $1/4$ -wavelength image shift.

Fig. 3.

The initial OFRs to the $3f5f$ stimuli: dependence on the contrast of the $3f$ component when the $5f$ component always had a contrast of 8% (R-L eye velocity profiles over time, real and simulated). The layout of columns and rows and the actual data obtained with the $3f5f$ stimuli (grey traces) are all as in Fig. 2. Least-squares best fit responses based on the CWA (red dashed traces), the sum (blue dashed-dotted traces), and the simple average (green traces). Other conventions as in Fig. 2.

Fig. 4.

The initial OFRs to the $3f5f$ stimuli: RMS errors for the least-squares best fit simulations based on the CWA (red circle), the sum (blue square), and the simple average (green diamond) of the response profiles to the corresponding $3f$ and $5f$ stimuli. Each graph shows data for one monkey. Individual plots show the errors when the $5f$ components had contrasts of 4% (continuous lines), 8% (dotted lines), and 16% (dashed lines). Abscissas, Contrast Ratio, C_{3f}/C_{5f} (log scale).

Fig. 5.

The initial OFRs to the $3f5f$ stimuli: dependence of the weight functions (w_{5f}) for the best-fit CWAs on the Contrast Ratio, C_{3f}/C_{5f} (log scale). Each graph shows data for one monkey. Individual plots show the simulations when the $5f$ components had contrasts of 4% (red dashed), 8% (green), and 16% (blue dashed-dotted). Individual data points (red, green, or blue circles) show the best-fit weight values (w_{5f}) computed by the weighted average model for every stimulus condition, that is, for each contrast of the $5f$ component and each contrast of the $3f$ component.

Fig. 6.

The initial OFRs to the $3f/5f$ stimuli: simulation of the response distributions to particular stimuli based on the winner-take-all model. A: Histograms of the distributions of the response measures obtained from monkey A with leftward pure $3f$ stimuli of contrast 8% (orange plot), rightward pure $5f$ stimuli of contrast 8% (green plot), and the $3f/5f$ stimulus whose $3f$ and $5f$ components had matching directions and contrasts (blue plot); smooth curves are best-fit Gaussian functions. B: Histograms of the simulated $3f+5f$ distributions for monkey A obtained by summing the measured distributions for the leftward pure $3f$ stimuli and the rightward pure $5f$ stimuli but weighted in accordance with the measured Response Ratio of 0.61 (black/grey plot). Histograms were generally binned using custom Matlab subroutines in which the optimal bin width for each individual distribution was given by $2(IQR) N^{-1/3}$, where IQR is the interquartile range (the 75th percentile minus the 25th percentile) and N is the number of samples; the only exception to this was the histogram of the simulated $3f/5f$ data, for which we used the same bin widths as for the real $3f/5f$ data.

Fig. 7.

The initial OFRs to the $3f7f$ stimuli: dependence on the contrast of the $3f$ component when the $7f$ component always had a contrast of 16% (R-L eye velocity profiles over time). Each column shows data for one monkey and each row shows data for $3f$ stimuli/components of one contrast (see labels near traces in left column). Data were obtained with $3f$ stimuli (red dashed traces), $7f$ stimuli (blue dashed-dotted traces), and $3f7f$ stimuli (grey traces). The responses of a given monkey to the pure $7f$ stimuli are all identical. Other conventions as in Fig. 2.

Fig. 8.

The initial OFRs to the $3f7f$ stimuli: dependence on the contrast of the $3f$ component when the $7f$ component always had a contrast of 16% (R-L eye velocity profiles over time, real and simulated). The layout of columns and rows and the actual data obtained with the $3f7f$ stimuli (grey traces) are all as in Fig. 7. Least-squares best fit responses based on the CWA (red dashed traces), the sum (blue dashed-dotted traces), and the simple average (green traces). Other conventions as in Fig. 2.

Fig. 9.

The initial OFRs to the $3f7f$ stimuli: RMS errors for the least-squares best fit simulations based on the CWA (red circle), the sum (blue square), and the simple average (green diamond) of the response profiles to the corresponding $3f$ and $7f$ stimuli. Each graph shows data for one monkey. Individual plots show the errors when the $7f$ components had contrasts of 4% (continuous lines), 8% (dotted lines), and 16% (dashed lines). Abscissas, Contrast Ratio, C_{3f}/C_{7f} (log scale).

Fig. 10.

The initial OFRs to the $3f7f$ stimuli: dependence of the weight functions (w_{7f}) for the best-fit CWAs on the Contrast Ratio, C_{3f}/C_{7f} (log scale). Selected data from monkey A when the contrast of the $7f$ component was 8% (green line) and 16% (blue dashed-dotted line), and from monkey C when the contrast of the $7f$ component was 16% (blue dashed-dotted line). Individual data points (green or blue circles) show the best-fit weight values (w_{7f}) computed by the weighted average model for every stimulus condition, that is, for each contrast of the $7f$ component and each contrast of the $3f$ component. The data points shown are for those cases in which the ratio of maximum error values to minimum error values was more than 1.5 and the weight value of the $7f$ component was between zero and one.

Table 1

The CWA model: least squares best fit parameters for the 3f5f data

Monkey	5f contrast	n_{3f}	n_{5f}	r^2	5%	95%
A	4%	3.7	3.8	0.988	2.30	0.47
	8%	3.2	3.4	0.989	2.86	0.45
	16%	3.3	3.6	0.988	3.14	0.53
B	4%	1.9	1.9	0.930	4.71	0.21
	8%	3.4	3.7	0.947	2.86	0.51
	16%	2.8	3.0	0.967	3.49	0.43
C	4%	1.7	1.7	0.924	5.65	0.18
	8%	1.9	2.0	0.967	5.25	0.24
	16%	2.0	2.1	0.955	5.01	0.26
Mean $\pm$ SD		2.7 $\pm$ 0.8	2.8 $\pm$ 0.9	0.962 $\pm$ 0.025	3.92 $\pm$ 1.24	0.36 $\pm$ 0.14

5% and 95% refer to the Contrast Ratios when w_{5f} has values of 0.05 and 0.95, respectively.

Table 2

The CWA model: least squares best fit parameters for the $3ff$ data

Monkey	$7f$ contrast	n_{3f}	n_{7f}	r^2	5%	95%
A	8%	3.8	3.3	0.962	1.65	0.35
	16%	3.1	3.1	0.964	2.59	0.39
C	16%	2.5	2.4	0.924	2.91	0.28
	Mean $\pm$ SD	3.1 ± 0.7	2.9 ± 0.5	0.950 ± 0.022	2.38 ± 0.65	0.34 ± 0.06

5% and 95% refer to the Contrast Ratios when w_{7f} has values of 0.05 and 0.95, respectively.