50 Years Ago in CORR

Postspinal Anesthesia Osteomyelitis of the Lumbar Spine

P. L. Day MD and J. J. Hinchey MD CORR 1958;11:185–193

Infections following spinal anesthesia are rare. In a meta-analysis of 1.37 million women having epidural anesthesia for childbirth, Ruppen et al. [10], estimated a rate of deep infection of 1 in 145,000. Infections following spinal anesthesia have been well recognized for a long time. Day and Hinchey, in *Clinical Orthopaedics and Related Research*, reported five patients with vertebral osteomyelitis following spinal anesthesia [3]. Four of the five patients had spinal anesthesia and one had multiple diagnostic lumbar punctures. All developed back symptoms within 1–8 weeks. The person performing the punctures did not use gloves in four of the five patients; presumably the extremely low incidence of infections led some to believe rigorous sterile procedures were unnecessary. Also in four of the five cases the symptoms seemingly resolved with nonoperative treatment (followup times were not given in several patients).

Importantly, Day and Hinchey commented, “In each instance the diagnosis was delayed by failure to recognize the condition.” They also commented, “It is not our intention to discourage lumbar puncture for any purpose, whether diagnostic or therapeutic, but rather to call attention to the possibility of a serious major complication” [3].

Spinal and epidural anesthesia are widely used today owing to their

remarkable safety and efficacy. These two forms of anesthesia are particularly considered and used for total hip and knee arthroplasty not only because they provide effective anesthesia, but also because of evidence they reduce the risk of pulmonary embolus and deep vein thrombosis [12, 13] and evidence some anticoagulants increase the overall death rate [8, 9, 11] even if they reduce the rates of deep vein thrombosis [1, 2, 4–6]. Given the frequency of such anesthetics, it is remarkable we see so few complications, particularly infection. However, isolated cases still do occur [7] and the orthopaedist should be aware of the problem in patients with back pain after surgery in which spinal anesthesia was used. As noted by Day and Hinchey [3], “A diagnosis of postspinal anesthesia osteomyelitis should be considered in any case of persistent pain at the site of a lumbar puncture soon after the puncture.”

Richard A. Brand MD
Editor-in-Chief
*Clinical Orthopaedics
and Related Research*

References

1. Colwell CW Jr. Consortium data: comparative efficacy of low molecular weight heparin and warfarin following total hip replacement. *Orthopedics*. 1995;18 Suppl:21–23.
2. Colwell CW Jr, Collis DK, Paulson R, McCutchen JW, Bigler GT, Lutz S, Hardwick ME. Comparison of enoxaparin and warfarin for the prevention of venous thromboembolic disease after total hip arthroplasty. Evaluation during hospitalization and three months after discharge. *J Bone Joint Surg Am*. 1999;81:932–940.
3. Day PL, Hinchey JJ. Postspinal anesthesia osteomyelitis of the lumbar spine. *Clin Orthop Relat Res*. 1958;11:185–193.
4. Francis CW, Pellegrini VD Jr, Marder VJ, Totterman S, Harris CM, Gabriel KR, Azodo MV, Leibert KM. Comparison of warfarin and external pneumatic compression in prevention of venous thrombosis after total hip replacement. *JAMA*. 1992;267:2911–2915.
5. Francis CW, Pellegrini VD Jr, Totterman S, Boyd AD Jr, Marder VJ, Liebert KM, Stulberg BN, Ayers DC, Rosenberg A, Kessler C, Johanson NA. Prevention of deep-vein thrombosis after total hip arthroplasty. Comparison of warfarin and dalteparin. *J Bone Joint Surg Am*. 1997;79:1365–1372.
6. Heit JA, Colwell CW, Francis CW, Ginsberg JS, Berkowitz SD, Whipple J, Peters G. Comparison of the oral direct thrombin inhibitor ximelagatran with enoxaparin as prophylaxis against venous thromboembolism after total knee replacement: a phase 2 dose-finding study. *Arch Intern Med*. 2001;161:2215–2221.
7. Kranke W, Angster R, Forst H. Epidural abscess following a lumbar epidural catheter for mobilization of

50 Years Ago in CORR

- the knee [in German]. *Anaesthesist*. 1999;48:169–172.
8. Murray DW, Britton AR, Bulstrode CJ. Thromboprophylaxis and death after total hip replacement. *J Bone Joint Surg Br*. 1996;78:863–870.
 9. Murray DW, Carr AJ, Bulstrode CJ. Pharmacological thromboprophylaxis and total hip replacement. *J Bone Joint Surg Br*. 1995;77:3–5.
 10. Ruppen W, Derry S, McQuay H, Moore RA. Incidence of epidural hematoma, infection, and neurologic injury in obstetric patients with epidural analgesia/anesthesia. *Anesthesiology*. 2006;105:394–399.
 11. Sharrock NE, Gonzalez Della Valle A, Go G, Lyman S, Salvati EA. Potent anticoagulants are associated with a higher all-cause mortality rate after hip and knee arthroplasty. *Clin Orthop Relat Res*. 2008;466:714–721.
 12. Sharrock NE, Haas SB, Hargett MJ, Urquhart B, Insall JN, Scuderi G. Effects of epidural anesthesia on the incidence of deep-vein thrombosis after total knee arthroplasty. *J Bone Joint Surg Am*. 1991;73:502–506.
 13. Sharrock NE, Salvati EA. Hypotensive epidural anesthesia for total hip arthroplasty: a review. *Acta Orthop Scand*. 1996;67:91–107.

50 Years Ago in CORR:

Postspinal Anesthesia Osteomyelitis of the Lumbar Spine

P. L. Day MD and J. J. Hinchey MD

Richard A. Brand MD