any doubt between a solid lung and a pleural effusion. On looking at the skiagraph Fig. 1 (the chest is viewed from the

Fig. 1.

back) we see on the left side a moderately dense uniform shadow, with a well-defined curved margin extending downwards and inwards from near the apex to the articulation of the seventh rib, where the shadow becomes continuous from spine to ribs until it merges into the shadow of the diaphragm. The shadow is not sufficiently dense to obliterate the outline of the ribs, but is sufficiently dense to obscure the outline of the heart, which is not displaced to the right. On looking at the ribs on the affected side an alteration in their normal curve is seen as compared with the opposite side. The ribs have suffered a rotation upwards and outwards. A clear oval space is observed between the well-defined edge of the shadow and the spine, corresponding, no doubt, to the compressed but not altogether airless lung. The diagnosis of fluid in the chest was made. The chest was tapped and 30 ounces of serum withdrawn. Another point worthy of note is that the skiagraph showed no evidence of tubercle. The after-history of this case has quite justified the opinion expressed at the time the chest was tapped, that the effusion was a simple one, and not complicated with any tuberculous mischief of the lung or pleura.

CASE II.—A male, aged 19, was admitted to the City of London Hospital for Diseases of the Chest, with the diagnosis of an empyæma of the right side. On taking a skiagraph of this patient's chest it was at once seen (Fig. 2) that the whole of

Fig. 2.—Chest viewed from the back.

the right chest was occupied by a very dense uniform shadow, completely obliterating the outline of the ribs on that side, and blending with the heart shadow, which was displaced to the left. The chest was tapped and the pus withdrawn. The skiagraph (Fig. 3) was taken after the patient's convalescence. The difference between the two skiagraphs is

Fig. 3.—Chest viewed from the back.

very striking. The heart has returned to its normal position. The right side of the chest is nearly as clear as the left, but not quite, some pleural thickening remaining as the shadow shows; a curved line may be noticed to the right of the X, this corresponding to the track of the sinus, and extending down to the scar left in the chest wall. Again, in this case the skiagraph shows no evidence of tubercle. The patient still remains quite well.

As the above cases illustrate, in doubtful cases of pleural effusion the Roentgen rays will enable us to say if fluid be present or not, and from the density of the shadow to give an opinion as to whether such fluid be serum or pus.

FOUR CASES OF PRIMARY THROMBOSIS OF CEREBRAL VEINS AND SINUSES IN CHILDREN.*

By THEODORE FISHER, M.D., M.R.C.P.,

Physician to Out-patients, Bristol Hospital for Sick Children;
Pathologist, Bristol Royal Infirmary.

THE morbid anatomy of hemiplegia of acute onset occurring in children still remains in great measure a mystery. Encephalitis is possibly responsible for some cases, while thrombosis of veins or arteries may be present in others; but *post-mortem* evidence of the existence of either inflammation of the brain substance or of acute affections of the cerebral vessels is small. Even in those instances where the hemiplegia has complicated acute disease of serious nature necropsies are rare. Thus in 30 cases of hemiplegia occurring in diphtheria collected by Thomas,¹ only two necropsies are mentioned. In one there was thrombosis of the middle cerebral artery, and in the other a small hæmorrhage in the lenticular nucleus. In another case of diphtheria recorded by Dr. Edgeworth,² in which I made the necropsy the thrombosis of the basilar artery was present. Sir William Gowers³ has suggested thrombosis of the cerebral veins as a cause of infantile hemiplegia, and quotes a case recorded by Angel Money of such thrombosis occurring after scarlet fever. A case mentioned by Abercrombie⁴ is also sometimes referred to, but that is by no means a clear example of hemiplegia secondary to thrombosis of a cerebral vein. A more suggestive case is one recorded by Handford,⁵ which proves that primary thrombosis

* Read before the Society for the Study of Disease in Children.

may occur in the venous circulation of the brain of an apparently healthy child. But apart from the question of the possibility of infantile hemiplegia being caused by thrombosis of the cerebral veins, the occurrence of such thrombosis is interesting on account of its rarity. Osler, for example, has only met with two instances and they were associated with meningitis.⁶ In the following cases there was no disease of the cerebral meninges that could have been responsible for the thrombosis.

CASE I. History and State on Examination.—A girl, aged 4, was admitted November 23rd, 1896, into the Bristol Royal Infirmary, under the care of Dr. Shingleton Smith, suffering from ascites of short duration. Swelling of the abdomen had been noticed for a week only. The abdomen measured 25 inches at the level of the umbilicus. A week later the distension of the abdomen was 1 inch less. On December 1st, at 10.15 P.M., the child had an attack of convulsions which lasted 3 minutes. The convulsions were followed by twitchings of the left side which persisted for 8 hours. On December 2nd and 3rd twitchings occurred again at intervals, and on the 4th her mouth was noticed to be drawn to the right. On December 5th the left upper and lower extremities were found to be paralysed. It may be worthy of mention that on the night of the convulsions the temperature rose to 100° F., and on the following day it reached 103°, but in the course of the next two days returned to the normal. In spite of the hemiplegic attack the child in many ways seemed to be improving. The abdomen steadily decreased in size, and on December 16th measured 4½ inches less in circumference than at the time of admission. On December 21st, however, nearly three weeks after the onset of the convulsions, the child complained of sore throat; membranous tonsillitis proved to be present, but no diphtheria bacilli were found. The larynx became implicated, and the child died.

Necropsy.—The superior longitudinal sinus was found to contain partially decolorised and softening thrombus throughout its length, which, however, was not sufficient to obstruct more than half the diameter of the lumen. On removal of the dura mater all the veins running up to the superior longitudinal sinuses on the right side of the brain were seen to be full of pale thrombus, and a large number of veins over the left cerebral hemisphere were also thrombosed. The appearance and consistency of the clot in the veins on the two sides of the brain did not materially differ, but it was evident from the difference in aspect of the convolutions of the two hemispheres that thrombosis had commenced on the right side. While the convolutions of the left side preserved a natural appearance except for congestion of the meninges covering them, the convolutions of the right hemisphere were greatly swollen, and had become flattened against the dura mater by the pressure to which their increased size had subjected them. The upper end of the right ascending parietal convolution measured three times as much across as the same convolution on the opposite side. The convolutions most affected were extremely soft, and dotted over with punctiform hæmorrhages. The frontal, parietal, and occipital lobes had all suffered, but the upper parts of the frontal and parietal lobes were most affected. The lateral sinuses on both sides also contained patches of *ante-mortem* thrombus, and the right internal jugular was blocked throughout its course by adherent but softening clot. Examination of the tympanic cavities showed that the right contained mucus only, but the left contained some mucus. This, however, was probably of recent date, and secondary to the membranous tonsillitis which was present. The peritoneal cavity contained only a few ounces of fluid, but well-marked cirrhosis of the liver was present. Several divisions of the portal vein contained pale thrombus. The blocking of these vessels had led to some congestion of the small intestine. There was nothing noteworthy in any of the other organs except the larynx and bronchi. Membranous exudation extended from the larynx to the small bronchial tubes. Some broncho-pneumonia of the lobes of the lower lungs was present.

It is worthy of note that the onset of the cerebral symptoms occurred nearly a month before death. This suggests that the cerebral veins were first affected. Thrombosis of the right jugular vein probably followed considerably later, and the clotting found in the branches of the portal vein must have been of quite recent origin, since the ascites had been steadily disappearing, and at the time of death no longer existed. The presence of diphtheria, a disease which sometimes gives rise to *ante-mortem* clotting, may have led to rapid increase of the thrombosis. The disappearance of the ascites has already been referred to. Peritoneal fluid was being rapidly absorbed when the hemiplegia occurred, and it seems possible that some organic body, absorbed from the peritoneal cavity, may have started the thrombosis. Twice I have seen extensive thrombosis of pulmonary veins in chronic peritonitis with effusion. No cultures, however, were taken from the clot in this case, so that the presence of micro-organisms cannot be excluded, but the discovery of such organisms would not have proved the primary cause of the thrombosis to have been infection, since microbic invasion possibly followed the membranous tonsillitis. Whatever may have been the cause of the thrombosis of the cerebral veins, we may with some reason conclude that the thrombosis of the veins over the right cerebral hemisphere occurred at the time of the onset of the convulsions; and it is also probable that, had it not been for an attack of membranous tonsillitis and laryngitis, the child would have left the infirmary temporarily free from ascites, but with left hemiplegia.

CASE II. History and State on Examination.—A child, aged 2, was admitted into the Bristol Royal Infirmary, under the care of Mr. Munro Smith, in September, 1897, for noma of the left cheek. A few days after admission cerebral symptoms, which were thought to indicate meningitis, set in. Twenty-four hours after their onset the child died.

Necropsy.—The superior longitudinal sinus was found free from *ante-mortem* clot, but the meningeal veins over both hemispheres were thrombosed, the meninges intensely congested, and the convolutions studded with petechiæ. Both lateral sinuses contained a little *ante-mortem* clot. The right tympanum contained a little mucus, but the left was quite free. Dr. Symes found a bacillus, both in the discharge from the ulcer on the cheek and in a culture taken from one of the thrombosed cerebral vessels, which he considered corresponded to the bacillus described by Schimmelbusch as occurring in cases of noma.

The next case is of greater interest, but instead of the superficial veins being affected, the choroid plexuses and veins of Galen were thrombosed. The thrombosis was secondary to summer diarrhœa.

CASE III. History and State on Examination.—A female infant, aged 5 months, was admitted into the Bristol Royal Infirmary under the care of Dr. Prowse for diarrhœa and vomiting on August 5th, 1899. Four days later spasmodic opisthotonos set in and continued at frequent intervals for forty-eight hours. Some rigidity of all the limbs was present. There was convergent squint of both eyes, and some lateral and vertical nystagmus. After the opisthotonos passed off, rigidity of the left side remained. The knee-jerks were exaggerated on both sides. The diarrhœa ceased, but the child remained ill. The temperature, which was 100.4° F. on admission, continued high: it reached 106° on the second day of the opisthotonos, but it was usually at some point between 103° F. and 104° F. On the day of death, which took place on August 14th, seven days after the onset of the opisthotonos, the temperature rose to 106° F.

Necropsy.—The exterior of the brain presented a perfectly normal appearance. No tubercles were present, and there was no trace of meningitis. The arteries and veins were everywhere quite normal and free from *ante-mortem* thrombus. On resting the brain upon its base the cerebral hemispheres fell apart, owing to softening of the corpus callosum. The exposed ependyma of the lateral ventricles was much congested, and the brain substance in places dotted over with punctiform hæmorrhages. Further examination showed the veins of Galen to be distended with pale *ante-mortem* clot. The choroid plexuses presented an unusual appearance. The left choroid plexus was five or six times its normal size, and the right also was much enlarged. Both were of dirty-white colour, and the surface of each was smooth compared with that of a normal choroid plexus. The appearance was found to be due to their distension with hard thrombus. A few small thrombi stood out from the commencement of the straight sinus, but the sinus itself was free from *ante-mortem* clot. All the other sinuses were also free from *ante-mortem* thrombus. The optic thalami were softened, but there was no hæmorrhage into their substance. Both tympanic cavities contained a little mucus. Some broncho-pneumonia was present at the bases of both lungs, but none of the other organs presented anything abnormal. Unfortunately the intestines had been washed out before coverslip preparations had been obtained. Cultures from the thrombosed plexuses proved sterile. Microscopic sections of a thrombosed choroid plexus showed the venous spaces filled with granular disintegrating clot. In many of the arterioles similar disintegrating clot was present, but others showed normal blood cells. The villi were glued together by fibrinous exudation, which in some parts was still fibrinous, but in others granular. In many places the epithelium over the villi was proliferating, and desquamated cells were entangled in the fibrinous exudation. No micro-organisms could be discovered.

Although no micro-organisms were discovered in the thrombosed plexuses it seems probable that the thrombosis was produced by bacterial invasion from the intestine. In connection with this point it may be worthy of mention that the previous summer in a case of death from summer diarrhœa in a child in which thrombosis of the right renal vein was found, cultures from the thrombus showed streptococci, and coverslip preparations from the contents of the small intestine also showed streptococci. The presence of marked opisthotonos in this case is possibly of some interest. Dr. Herringham has recorded a case in which a small glioma in the centre of the iter gave rise to attacks of this rigidity, and has suggested that it is produced by irritation of the grey matter which is continued upwards from the spinal cord.⁷ It is of interest to note in this connection that Dr. Goodall has recorded a case of thrombosis of the veins of Galen and choroid plexuses in a girl aged 8 years recovering from scarlet fever, in which general rigidity followed convulsions,⁸ and in another case occurring in a woman aged 25, recorded by Gibson and Stewart⁹; opisthotonos was present.

CASE IV. History and State on Examination.—A boy, aged 11 months, was admitted into the Bristol Royal Infirmary, under the care of Dr. Shingleton Smith, on February 23rd, 1900, with broncho-pneumonia. He had been ill for a week. He was drowsy when admitted. His head was held somewhat retracted during the first week he was in the infirmary, but no symptoms manifested themselves which led to the suspicion of any cerebral affection. The drowsiness, however, continued. The temperature was raised, sometimes nearly reaching 105° F., but occasionally dropping to the normal, or even below. He died three weeks after admission.

Necropsy.—On opening the dura mater, an excessive amount of cerebro-spinal fluid flowed away. Several of the cerebral veins contained the long ribbon-like pale thrombi not infrequently seen, which, if not entirely *post-mortem*, must be formed very shortly before death. In addition to these flat thrombi, thrombosis of more definite *ante-mortem*

character was present. A vein over the centre of the right parietal lobe was distended with hard pale *ante-mortem* clot. The left middle cerebral vein was also fully distended with hard stratified pale thrombus, and equalled a quill pen in diameter. The superior longitudinal sinus was quite free from *ante-mortem* clot; the straight sinus was also free, but attached to the anterior end of the straight sinus were two thrombi apparently drawn from the veins of Galen. In the left lateral sinus, where it passes below the tentorium cerebelli, was a star-shaped thrombus drawn out of veins entering from the occipital lobe. The sinus was partially filled with *ante-mortem* clot at this point for about the space of half an inch, but elsewhere was free. There was no *ante-mortem* thrombus in the right lateral sinus, but the left cavernous sinus contained mixed *ante-mortem* and *post-mortem* clot. There was some congestion of the meninges in the neighbourhood of the thrombi. No distension of the ventricles was present. Both tympanic cavities contained pus, but this is so common an occurrence in acute diseases in children that any relation to the thrombosis of the cerebral veins must be considered doubtful. The upper lobe of the right lung was solid throughout, but of the mottled character which frequently distinguishes a confluent broncho-pneumonia from a fibrinous pneumonia, and in this case there were also three or four small areas of commencing suppuration. There was nothing else noteworthy except in the large intestine, where the solitary follicles were much swollen, and here and there small islets of swollen mucous membrane surrounded several enlarged follicles. Cultures from the thrombosed veins and from the lungs showed the presence of streptococci.

In this case the absence of localising symptoms was noteworthy. The middle cerebral vein was completely blocked by the thrombus, and, although the clot was not commencing to break down, it was stratified, and probably had been in process of deposition during several days before death. There was, however, little congestion of the meninges in the neighbourhood, and no evidence of softening of the brain substance.

Two at least of the above cases illustrate the fact that thrombosis of the cerebral veins and sinuses may be a cause of cerebral symptoms, when they arise as a complication of disease in children. Especially is this likely to be the case when the disease is one in which micro-organisms are likely to obtain access to the circulation. In two of the above cases micro-organisms were found in the thrombus, and in a third, although not discovered, they were probably present in the blood. In the remaining case micro-organisms were not looked for, and, although we have thought it probable that thrombosis occurred independently of infection, we have no proof that such infection was absent.

A more interesting question than the causation of thrombosis of cerebral veins in children is the possibility of this thrombosis being a lesion which may give rise to hemiplegia in a healthy child. In one of the above cases hemiplegia resulted from thrombosis, which in the character of its onset resembled some cases of infantile hemiplegia. Generally the paralysis develops more rapidly, but apparently thrombosis of cerebral veins may occasion hemiplegia, which sets in suddenly. A brief history referring to a specimen recently added to the St. Bartholomew's Hospital Museum illustrates this point. The thrombosis occurred, not in a child, but in a woman aged 24 years. The age of the patient does not, however, materially detract from the value of one feature of the case.

After suffering from headache and pain in the back for a week she was admitted to the Brook Fever Hospital, Shooter's Hill. Within five hours of admission she was seized with complete right hemiplegia and aphasia, without loss of consciousness. Epileptiform fits followed, but she did not lose consciousness until just before death, which occurred five days later. The superior longitudinal sinus and right lateral sinus were found thrombosed, together with many of the surface veins opening into them.¹⁰

Yet although the hemiplegia set in suddenly in this case it had been preceded by headache and thus differs from most cases of hemiplegia occurring in children who are old enough to complain of their ailments. A perusal of a few of the recorded cases of thrombosis of cerebral veins and sinuses occurring in adults seems to show that headache which lasts for some days almost invariably precedes any local or general convulsions. This appears to be the case even when the thrombosis is limited to cerebral veins.¹¹

In a case, however, occurring in a child, a girl aged 7 years, recorded by Dr. Phear, the first symptom appears to have been toothache, which preceded unconsciousness of sudden onset only by a few hours,¹² and in another case recorded by Sir Dyce Duckworth, in a girl, aged 14 years, localised convulsions of the left hand occurred as a symptom of thrombosis of the superior longitudinal sinus and the veins entering it from the right side of the brain on the day headache was first complained of. In this case, however, the focal symptoms set in with much greater severity the following day. They were ushered in by a scream, but were associated with no loss of

consciousness.¹³ In Phear's case the child also screamed at the onset of the cerebral attack. Twice I have known mothers describe the onset of unconsciousness preceding hemiplegia in a child in apparently good health as being ushered in with a scream. One does not lay any stress upon this. A fit due to any cause may announce its onset by a scream, but Phear's case is interesting as showing that even thrombosis of a cerebral vein or sinus may manifest its presence by an attack of apoplectic suddenness, resembling that seen in infantile hemiplegia.

Allowing, however, that thrombosis of a vein may give rise to cerebral symptoms of acute onset, the cause of the thrombosis has still to be considered. When the thrombosis occurs as a complication of some disease, although the immediate factor in the causation of the *ante-mortem* clotting may not be obvious, its presence is comparatively easy of explanation. Thrombosis of a cerebral brain or sinus in a healthy child is an occurrence more difficult to understand, especially if it manifest itself without being preceded by symptoms that suggest even indisposition. Infection of some kind would be the most probable cause of the thrombosis, and one would expect more or less indication of an abnormal state of the blood before clotting in a vein takes place.

Yet this point cannot be strongly urged against the view that thrombosis is the cause of infantile hemiplegia. Consideration of other morbid conditions of the blood due to infection—for example, some varieties of purpura—shows that they may display themselves suddenly without warning. If, however, thrombosis of cerebral veins of infective origin be the cause of infantile hemiplegia, one would expect in at least a small percentage of cases that the thrombosis would be progressive and lead to a fatal ending. It is scarcely necessary to remark that such an event is virtually unknown. Yet it must be remembered that even when hemiplegia occurs as a complication of some acute disease, and in such cases vascular thrombosis is known to exist, the great majority of the children attacked recover.

A possible explanation for the thrombosis might be that it is set up, not by a blood infection, but by a toxin absorbed from the stomach or intestines. Belief, however, in such an origin would not enable one more readily to understand some of the features of infantile hemiplegia. It would not be clear why the dose of toxin should always be sufficient only to cause clotting limited to a few cerebral veins, and why it should not be repeated more than once during childhood.

We have been speaking of *ante-mortem* clotting in the veins. It is more easy to believe that thrombosis may occur in a healthy cerebral vein than in a healthy cerebral artery, but the occasional occurrence of thrombosis in a cerebral artery as a complication of some febrile states in children shows that arterial thrombosis must be considered as one of the possible causes of infantile hemiplegia.

If, however, the theory of cerebral vascular thrombosis as a cause of infantile hemiplegia be rejected, it seems necessary to look upon the presence of encephalitis as the only reasonable alternative. Here again we meet with difficulty. Although, when viewed from the clinical standpoint, infantile hemiplegia may not very closely resemble most of the recorded cases of thrombosis of cerebral veins and sinuses, it has still less in common with the descriptions of cases of encephalitis. In inflammation of the brain substance, premonitory symptoms are well marked, unconsciousness or mental confusion is prolonged, while the focal symptoms are variable, and often long in making their appearance. If encephalitis be the cause of infantile hemiplegia, it must be encephalitis of specific character.

[Since writing the above a case has been under my care at the Bristol Children's Hospital that seems to lend support to the vascular thrombosis theory of infantile hemiplegia. A girl aged 10 years, who was brought to the out-patient department for some slight ailment, had a large tortuous vein running up from the centre of the right groin over the right half of the abdomen. This was said to have appeared when the child was 18 months of age, towards the end of an attack of repeated convulsions, associated with unconsciousness, of a week's duration. The right leg and thigh were much swollen at the time the enlarged vein was first noticed. The child is

now weak-minded. She cannot read or be trusted with the most simple errand. Before the convulsions, however, she is said to have been bright and intelligent, and could say several words. It may be of interest to add that the mother attributed the convulsions to the falling of the child into a dirty stream two or three weeks before their onset. The child was nearly drowned, and after being pulled out vomited much dirty water. In this case it is obvious that blockage of the right external or common iliac vein is present, which may with reason be attributed to thrombosis occurring at the time the enlarged abdominal vein was first noticed. The evidence of thrombosis in one vein suggests that the associated cerebral symptoms were due to thrombosis within the cranial cavity. This thrombosis may have been of infective nature consequent upon the entrance of micro-organisms contained in swallowed dirty water into the blood.]

REFERENCES.

- ¹ *American Journal of Med. Sciences*, 1896, vol. cxi, p. 384. ² *Lancet*, 1899, vol. ii, p. 1563. ³ *Diseases of Brain*, second edition, p. 462. ⁴ *BRITISH MEDICAL JOURNAL*, 1887, vol. ii, p. 1325. ⁵ *Transactions of the Pathological Society of London*, vol. xxxviii, p. 40. ⁶ *Textbook of Medicine*, Pepper, vol. i, p. 694. ⁷ *St. Bartholomew's Hospital Reports*, vol. xxxvi, p. 31. ⁸ *Clinical Society's Transactions*, vol. xxx, p. 139. ⁹ *Edinburgh Hospital Reports*, vol. iii, p. 454. ¹⁰ *St. Bartholomew's Hospital Reports*, vol. xxxvi, p. 308. ¹¹ *Vide* a case recorded by Richardson, *Journal of Mental and Nervous Diseases*, 1897, p. 403. ¹² *Clinical Society's Transactions*, vol. xxxi, p. 5. ¹³ *Ibid.*, vol. xxiii, p. 101.

SYPHILITIC MENINGO-MYELITIS : ERYSIPELAS : RECOVERY.

By E. F. TREVELYAN, M.D., M.R.C.P.,

Honorary Physician to the Leeds Public Dispensary; Assistant Physician to the Leeds General Infirmary.

THE following case presents several points of interest :

History.—A woman, aged 23, had a miscarriage a few months after her marriage three years before the onset of the paralysis. Her first and second child died 7 and 2 months respectively after birth. She herself had been under treatment at the Leeds General Infirmary for syphilis, and her husband was known to be syphilitic. Six weeks before the commencement of the paraplegia she complained of pain and cramp in the left, and later in the right leg. This was followed by girdle pain. On February 24th, 1900, her right leg became weak on her way up to the dispensary, and she could only continue her journey with difficulty and with assistance. When she woke up on the morning of February 26th she had lost power in the left leg, and the paraplegia soon became complete. There was loss of sensation in the legs, and all the reflexes were slightly exaggerated. There was incontinence of urine, and a large sacral bed-sore was noticed on March 17th.

Condition on Admission.—The patient was admitted into the Leeds Infirmary under my care on March 17th. There was at this time an almost complete paraplegia with considerable rigidity, but no wasting. The reflexes were all increased, but no ankle clonus could be obtained. There was incontinence of urine and to a less extent of feces. The urine was alkaline, and contained a small quantity of pus. Tactile sensation was impaired. There was the remains of a syphilitic rash on the chest, as well as scars about the mouth.

After-History.—On March 24th an attack of facial erysipelas supervened. Two injections of antistreptococcus serum were given. The temperature became normal on March 30th, but there was a further slight rise of short duration on April 6th. On April 3rd it was noticed that she could move her toes, and the power in the legs began to return rather rapidly. Unfortunately, owing to the cleaning of the wards she had to be sent home on April 14th, and she was again attended from the dispensary. On June 12th she walked up to the out-patient room with the help of a stick. The knee-jerks were exaggerated, especially on the left side. All forms of sensation appeared to be normal. The bladder had recovered completely. On June 28th she walked up without a stick, and on July 19th the recovery appeared to be quite complete. On April 11th, 1901, she was carefully examined. She was quite well, with the exception of a slight weakness of the anal sphincter; thus she said that she could not hold her motions for more than a minute. The sphincter did not grasp the finger at all strongly.

The diagnosis of an incomplete myelitis involving the lumbar enlargement and part of the lower dorsal cord was based on the spastic paralysis, loss of sensation, paralysis of the sphincters, and bed-sore. The presence of pains and cramps in the legs showed that the meninges were also involved. The meningo-myelitis was believed to be syphilitic for the following reasons: (1) There was not only undoubted evidence of past syphilis, but the patient still presented manifestations of active syphilis when the paralysis supervened. (2) The presence of paralysis combined with rigidity and exaggeration of the reflexes, is known to occur in spinal syphilis; and (3) the absolute absence of any other effective cause. There was apparently some variation in the severity of the symptoms during the early stage of the disease, and this also has been associated with spinal syphilis by some

writers. It was imagined that the lesions consisted of an inflammation of the pia-arachnoid, most probably of a genuinely syphilitic nature, and that the disease had spread into the cord which it affected in a patchy fashion, and not uniformly.

A point of peculiar interest in the case was the attack of erysipelas, which was followed, at least in point of time, by improvement in the paralytic symptoms. There is nothing preposterous in assuming a direct relationship between the erysipelas and the recovery, as one infection has been known both experimentally and clinically to antagonise another and independent infection. By some curious oversight the patient received no antisyphilitic treatment between the onset of the paralysis and June 28th. There is reason to believe, though unfortunately the out-patient notes cannot be found, that the patient was under antisyphilitic treatment during at least a part of the twelve months preceding the paralysis.

The importance of the recognition of cases of spinal syphilis is twofold, namely, on account of its relation to (a) prognosis and (b) treatment, although the above-related case hardly exemplifies the latter statement. Williamson (of Manchester) in his interesting paper on spinal syphilis¹ mentions 32 cases of various forms of the disease. Of the 32 cases 10 recovered and 9 died, the remaining 13 passing from observation without any great change in their condition. Five out of 6 cases of acute syphilitic paraplegia died. While insisting on the importance of antisyphilitic treatment Dr. Williamson makes the very guarded statement that the prognosis of spinal syphilis is on the whole better than that of other affections of the spinal cord. Obviously the extent to which the spinal cord is involved in these cases is of the first importance in the prognosis.

Unless the actual lesions present in the membranes and cord are syphilitic (gummatous) in character or perhaps due to syphilitic vascular disease, little hope of any good can be expected of antisyphilitic treatment. As, however, we are mostly not in a position to say whether these lesions are genuinely syphilitic or not, a course of antisyphilitic treatment consisting in mercurial inunction and potassium iodide should always be had recourse to unless it is certain that the patient has recently been efficiently treated in this way. If no result is obtained in a few weeks such treatment should be abandoned, at least for some time, as it may possibly do injury instead of good.

I have to thank Mr. H. R. Cross, one of the resident medical officers to the dispensary, and Mr. Hirst, the clinical clerk to the case, for additional notes of the case.

REFERENCE.

- ¹ *Edinburgh Medical Journal*, October, 1900.

CASE OF CEREBRAL TUMOUR AT THE PARIETO-OCCIPITAL FISSURE.

By R. T. WILLIAMSON, M.D. LOND., F.R.C.P.,

Physician to the Ancoats Hospital, Manchester, and Assistant Lecturer on Medicine, Owens College.

THE following record of a case of cerebral tumour presents several points of interest with respect to the localisation of cerebral lesions.

G. P., aged 43, was admitted to the Ancoats Hospital on December 31st, 1900, complaining of headache and failure of sight.

History.—His illness had commenced nineteen weeks previously with troublesome headache and sleeplessness. At that time vision was not affected. After the headache had continued for two or three weeks he began to be troubled with slight twitches in the left leg; these continued from time to time all the morning and afternoon, and prevented him from walking about. The same day, about 5 P.M., the twitches in the left leg became suddenly more severe and rapid, and developed into a true clonic convulsion. The twitches spread from the left leg to the left arm and left side of the face. The face was drawn to the left, and the tongue bitten. After a few minutes the patient lost consciousness, and finally the right side of the body was convulsed also, but not so markedly as the left. The attack was therefore a true unilateral convulsion (Jacksonian epilepsy) beginning in the left leg. The duration of the fit was about twenty minutes. The patient fell asleep afterwards, and he remained in a semi-conscious state for four days. About thirty hours after the first convulsion there was a second of a similar nature, and again about twenty hours later there was a third. In all three fits the spasm commenced in the left leg; it then spread to the left arm and left side of the face, and remained unilateral for some time before the right side was affected. The patient recovered consciousness completely at the end of four days. There was no paralysis. The sight began to fail