solids. The muscle was stained by Marchi method, and shows that quite one half of the fibres have undergone acute fatty degeneration. The dense black masses are fat cells; but the degenerated muscle fibres are indicated by the diffuse black staining. The musculi papillares of the heart showed even more marked fatty degeneration.

Figures 1, 2, 3, 4, 5, 7, 8, 9, 13, 14, 15, 16 are photomicrographs of preparations stained by the Nissl method. Figure 6 was stained by the Weigert iron method. Figures 10 and 11 were stained by the rapid Golgi method. Figures 12 and 17 were stained by the Marchi method.

ON LEAD ENCEPHALOPATHY AND THE USE OF DIACHYLON AS AN ABORTIFACIENT.

By W. B. RANSOM, M.A., M.D., F.R.C.P.,
Physician to the General Hospital, Nottingham.

In the following paper I propose to record three cases of cerebral disease due to lead poisoning, two of which were caused by diachylon taken to procure abortion, and some other cases in which diachylon taken with the same object produced a different set of symptoms:

CASE I.—A youth of 17 was sent into hospital for cerebral tumour.

History.—For three weeks he had suffered from headache, vomiting—chiefly after food—colicky pains in the abdomen and dimness of vision. The patient was a carriage painter who for 2 years had been engaged on fine work, but for the last three months had taken to coarse work in which much red lead was used.

Condition on Admission.—He was found to be well nourished but anæmic, lethargic, but complained of intense frontal headache. No paralysis, spasm, or anæsthesia, pupils and reflexes normal. Well marked double optic neuritis and a large retinal hemorrhage in the left eye. Heart somewhat dilated, blowing murmur at the apex; soft, regular pulse. Urine: specific gravity 1012 to 1016, free from albumen and sugar. Bowels open, tongue moist, furred and tremulous. Slight blue line on the gums.

Treatment.—The bowels were kept open and potassium iodide and bromide, with an occasional dose of antipyrin, ordered. A fortnight later he was no better, and then digitalis was added to his medicine. Within a week he was better, and all his symptoms left him five weeks after admission. The vision improved and the optic neuritis became perceptibly less, but some yellow striæ developed round each macula lutea. A trace of albumen was found four times in the urine, the daily output of which increased under the use of digitalis from 30 to 50 ounces, while its specific gravity decreased to 1006 to 1010, with urea only 1 per cent. The blue line was not quite gone when he left the hospital.

In this case the brunt of the disease fell on the brain, but clearly the kidneys, alimentary tract, and heart also suffered. I am sorry not to have been able to trace the further history of this patient, who was under my care eight years ago.

CASE II.—In April, 1899, I saw in consultation a married woman, aged 39, for brain disease.

History.—She was said to have had severe nervous symptoms and a mysterious abdominal pain a year ago, but these abated when it was proposed to send her into hospital. She had had two live children, the youngest 15 months old, and six miscarriages, but there was no history of a syphilitic rash. Apparently she had not been really well for a year, but had for four or five months suffered severely from headache and vomiting without relation to food. She had had three fits within the previous three months. A month ago she began to be unsteady on her feet, and had been confined to bed for three weeks. For a week she had been semi-delirious, having visual hallucinations, and for two days had had diplopia. No information as to her menstrual history could be obtained, either then or during her stay in hospital.

Anamnesis.—Only an imperfect examination was possible in the small dark room of her home, but we noted no paralysis, ataxy resembling that of cerebellar disease, and double optic neuritis. A week later she came into hospital, where in the well-lit ward I at once observed on her gums the most intense blue line I have ever seen. The notes say: "Patient is a stout, florid woman; intellect and memory poor, but she can answer some questions fairly; has some visual hallucinations at night; complains of severe headache on the vertex. She can sit up in bed, but can hardly stand or walk alone, tottering with the head craned forward and swaying from side to side. The ataxy is not worse when the eyes are shut. No muscular wasting or tremors, and no local paralysis of the limbs, though all are weak. No anæsthesia. Knee-jerks well marked. The pupils are of medium size, the left a shade the larger; they react normally. There is some convergent strabismus, as the left eye cannot be moved completely outward. There is occasional diplopia; no facial or lingual paralysis; speech normal. She can read ordinary type with ease; hears a watch at 5 inches from left ear, at 15 from right ear. The ophthalmoscope shows double optic neuritis, swelling of disc = 1.5 D. in each eye. There are hemorrhages on and around each disc; no choroiditis or sign of syphilis. The urine contained a trace of albumen for the first two days only. The uterus showed no distinct sign of a recent miscarriage and was not enlarged.

Treatment.—The patient was put upon potassium iodide (gr. xv daily) and magnesium sulphate, and improved with wonderful rapidity. Within three weeks the headache had gone, she was quite rational, and could walk well and steadily. She left the hospital a month after admission with the weakness of the left rectus externus muscle nearly gone and the optic neuritis diminishing. The blue line had also nearly gone.

This patient admitted having taken diachylon in pills for the purpose of checking the course of pregnancy, but we could not ascertain from her exactly when or what quantity she had taken, or what the effect had been on the uterus. I called,

however, to see Mrs. C. on March 20th, 1900, and found her in perfect health, and the proud mother of a three weeks old baby. There was neither blue line nor squint, she could read the local newspaper with ease, and the ophthalmoscope showed nothing abnormal but the faintest haze over the origins of the vessels in the optic discs. She must have conceived immediately on leaving the hospital, and did not repeat her experiments on ecobolics. The baby is, however, a puny creature, said to have "consumption of the bowels," but is not hydrocephalic, as the children of lead workers are at times. On interrogation, she now asserted that she had taken diachylon for a fortnight a year before I saw her, being then six weeks pregnant. Abortion took place at the end of the third month. She adhered to the statement that she had taken none since, although the symptoms of poisoning steadily increased up to her admission into hospital. I have no means of testing the truth of this statement.

CASE III.—H. W., an unmarried girl, aged 22, was admitted in June, 1899, for pains in the stomach and head, of three or four months' duration.

History.—She had been treated outside for gastric ulcer and kept on milk diet, as solid food caused acute pain. There had also been some vomiting and some severe colicky pains. For the last month she had suffered from intense headache chiefly in the frontal and right occipital regions. On interrogation, the patient admitted that in the March previous, having gone a week beyond her proper menstrual period, she had taken one day a half-pennyworth of diachylon, which she had made into two pills. She was positive that she had never taken any more. The period soon came on, and her menses had been regular since, but she began to feel weak and dizzy within a week of taking the pills.

Anamnesis.—After admission she was found to be a pale and rather thin girl with features drawn with constant pain in the head. A distinct blue line on the gums. She is drowsy, and occasionally a little delirious. She can sit up, but not stand alone, the knees doubling under her, and the feet being dragged. Knee-jerks absent; plantar reflexes well marked. Arms rather weak, but not paralysed. Pupils, eye and face movements normal; no anæsthesia; well-marked double optic neuritis. Her vision was unfortunately not properly tested at this time, but she could recognise the doctors and nurses, and I think read large type.

After-History.—During the first fortnight after admission she was frequently sick independently of food, and paresis of the left external rectus muscle was observed. For a month she suffered from intense headache, and remained drowsy and apathetic. On August 28th the optic neuritis had subsided, and the discs were atrophic. She was blind, and hardly able to point to the direction of the window. The paresis of the left external rectus had gone, and the general muscular tone and balancing power were much better. On September 10th she could walk well, and had the complete use of all her limbs, was free from headache, sickness, or abdominal pain, and was plump and rosy. The eyesight was, however, unaltered. I visited this patient at her home in March, 1900, and found her in excellent general health, but still blind. The medicinal and general treatment was the same as in the last case.

These three cases illustrate most of the usual symptoms of lead encephalopathy, and show how closely this affection resembles cerebral tumour. One symptom only of this disease was not prominent in these cases, namely, epileptiform attacks, which Dr. Oliver states are a frequent and early symptom. The woman had had three fits, but the other cases had none. One satisfactory feature of the cases was that all recovered, only the poor girl being left seriously injured. The fact that the married woman was said to have shown severe hysterical symptoms a year before is of interest, for a "toxic hysteria" has been found in other cases to sometimes precede the severe and even fatal form of the disease, and to have obscured the diagnosis. Possibly some cases diagnosed as hysteria which have ended fatally may have been due to undiscovered lead or other poisoning.

It is very remarkable that such acute, severe, and prolonged disease should have been produced in the girl by a single dose of lead—namely, one halfpennyworth of diachylon taken at once. To get some idea of the amount of lead taken by this girl I bought a pennyworth of diachylon at a chemist's, and had it analysed by Mr. Trotman, the city analyst. The diachylon, which was in the form of a ball about the size of a walnut, contained 86 per cent. of oleate of lead and 46.7 grs. of lead, so that we may suppose the girl to have taken about 23 grs. of lead.

This case proves that the excretion of the metal must be very slow, as does also that of Mrs. C., if her story be true.

The treatment of these cases in addition to rest, milk diet, and good hygienic conditions, consisted mainly of moderate doses of potassium iodide with Epsom salts. It is curious that the iodide should have obtained such a reputation in the treatment of lead poisoning, for Dr. Dixon Mann's researches seem to prove that it does not increase the elimination of the metal, the greater part of which escapes by the fæces and not by the urine. It should be noted, however, that in the first case improvement did not occur until the heart had been

strengthened and the flow of urine increased by digitalis. Dr. Oliver thinks that the iodide may occasionally even do harm in acute cases by redissolving lead which was lying inert in the tissues. There was, however, no evidence of this in these cases, nor of it causing albuminuria, as Dr. Oliver has observed in a few cases. I incline to think Epsom salts, with light diet and generally tonic conditions, the most useful elements of treatment. Hot or electric baths and massage are, however, praised by some.

Lead poisoning, as is well known, occurs in four forms. First, the acute form of brain disease which we have described; secondly, the chronic affection of peripheral nerves, which most frequently affects the posterior interosseous nerve and causes wrist-drop; thirdly, lead colic, in which we have acute paroxysmal attacks of pain, associated usually with chronic constipation and some anæmia and gastric disturbance; fourthly, chronic plumbic cachexia, in which the symptoms of chronic Bright's disease, and sometimes of gout, are predominant. Though these groups of symptoms may be more or less united in an individual, yet they may, as the cases I have reported show, be remarkably distinct.

I now pass on to some cases in which lead taken to procure abortion produced other symptoms.

CASE IV.—One evening last year a woman was sent into hospital from the country, with a doctor's letter, saying she had intestinal obstruction and fecal vomiting, and also that she was pregnant. I was asked to see her at 10 P.M. to decide whether immediate operation was necessary. We could not diagnose the cause of the obstruction, but as she was four months' pregnant, had had obstruction for three if not four days, and as the vomited matter was stated by a medical man to be fecal, we decided not to postpone operation till the morning. However, when the abdomen was opened no obstruction could be detected. There was a normal pregnant uterus, and the coils of intestine seemed all empty and tightly contracted. So the wound was sewn up and the patient put to bed. Next morning a distinct blue line was seen on her gums, and the "obstruction" yielded to hourly doses of Epsom salts. She admitted afterwards that she had taken diachylon to procure abortion. She was discharged some weeks later, with the wound well healed, and the pregnancy still progressing.

CASE V.—About two years ago a respectable young married lady who had had two children came to tell me she feared she was again pregnant, having gone three weeks over her time, and to implore me to help "put her right," as she and her husband did not want any more children. On my declining and lecturing her as to the iniquity and folly of such a proceeding, she started to do it herself by long and rough bicycle rides and other violent exercises. Ultimately she brought something away, and was ill with severe colicky pains at home without seeing a doctor. When she came to see me again, she had some metritis and a blue line on her gums. She admitted having in despair resorted to diachylon. With rest and nursing she made a good recovery. In this case there were no brain, nerve, or kidney symptoms, and apparently not much colic.

CASE VI.—Another case was one which I saw four years ago in consultation with Dr. Cornwall in Sneynton. A married woman of 34, when six weeks' pregnant took a halfpennyworth of diachylon and aborted in a fortnight. When I saw her a fortnight later she was very ill with colic, vomiting, dizziness, and some headache. She had very slight jaundice and a well-marked blue line. The right kidney was felt to be rather large and tender. The urine was dark coloured, but contained no bile or albumen, but a large precipitate of uric acid. There was no paralysis, the uterus was normal for its condition, but had been swabbed out by the doctor after the miscarriage.

From my own experience, therefore, I have reason to believe that the use of diachylon as an abortifacient is a fast-growing evil. There is now in hospital a woman with wrist-drop from this cause, and Dr. Handford tells me that he has had in his wards three cases of lead encephalopathy and two or three others of less serious lead poisoning, all from taking diachylon. Two of these brain cases had to be transferred to the City Asylum. Inquiries I have made among medical men in one quarter of the town only—Sneynton—lead to the same conclusion. Thus Dr. Cole has sent two cases into hospital in the last year, and has seen several others. Dr. Dabell writes:

I remember two cases of poisoning due to diachylon taken to procure abortion. One was mild, with no nerve symptoms; the other was more severe, and her peripheral nerves were affected. Both recovered. I am now visiting a patient in a fairly good position who, in addition to a large quantity of purgatives, took a pennyworth of diachylon in one day, but with no effect. It is not uncommon for women to ask me about this drug.

Possibly the purgatives taken by this lady saved her from the ill-effects of the lead. Dr. William Thomson says:

I have had a good many cases of abortion due to lead poisoning from pills taken. Abortion usually occurs before nerve symptoms set in, although I had one case with severe anæmia, jaundice, albuminuria, colic, and slight wrist-drop. I think the practice is very common and on the increase, as I can recall half a dozen cases within the last year or two.

Dr. Cornwall writes:

I have had several cases such as you mention, and found the brain

always more or less affected, also the peripheral nerves. Jaundice was common.

Thinking it would be of interest to ascertain whether persistent weakness of the brain often ensued, I wrote to Dr. Powell, of the City Asylum, who replied:

I have had two cases of insanity from the cause you name during the past year, both with melancholia, one having also active hallucinations of hearing. One recovered in three months, the other is now slowly recovering after a residence of five months, but there is much secondary mental enfeeblement.

These two cases were the two of Dr. Handford's already mentioned.

Dr. Moore Bennett, of Ruddington, lately told me the practice was common in his rural district, and kindly sent me brief notes of three cases in which the abortion was associated with lead colic due to diachylon. In one of these there was also wrist-drop and much tremor, in another coffee-coloured vomit, and in the other septic metritis. All recovered after severe illnesses. Dr. Bennett thinks that these cases usually suffer from septic mischief.

Dr. Cole also informs me that he has no doubt many women take the drug without ever coming under medical treatment for it, as he has been frequently told by patients that they have used it without the desired effect. He considers it a by no means certain abortifacient.

Turning to the medical journals of the last few years, I find several cases of poisoning by diachylon taken to procure abortion. Dr. Pope (Leicester)¹ records two cases which died with cerebral symptoms. The late Dr. Crooke (Birmingham)² described a similar fatal case, and expressed his belief in the common use of this drug. Dr. Bell Taylor³ gives a case in which the drug had no other effect than to produce optic atrophy and permanent blindness in the left eye. Dr. Branson⁴ relates the case of a multipara who took a pennyworth of diachylon in forty-eight hours when three months' pregnant. When seen a month later the uterus was empty, and she was suffering from severe colic. He adds that the Birmingham chemists told him diachylon is much used for this purpose.

There can be no doubt that diachylon is largely used by women of various classes to procure abortion. It is easily purchased. Anyone can go to a chemist's and buy a pennyworth of diachylon, as I have myself done, without being asked any question except as to whether the purchaser wants it spread or in the mass. Penny balls of the emplastrum plumbi are kept by the most respectable chemists ready wrapped in a handy drawer, and there is absolutely no restriction on its sale. For a penny a woman can buy enough lead not only to empty the uterus, but to cause grave disease of the bowels, the kidneys, and the brain, a disease which not rarely proves fatal. The drug appears to be an uncertain abortifacient, always endangers and often destroys life, or leaves permanent bodily and mental enfeeblement. It is a question whether something should not be done to restrict the indiscriminate sale of this drug and to class it with poisons. Of course, lead can be bought in other forms—as in paint or putty, but we might at least limit the evil, and reduce temptation by removing the present dangerous facility for manslaughter or suicide.

P.S.—The following case, seen as a hospital out-patient on May 11th, strikingly illustrates the subject of this paper:

CASE VII.—The patient, a married woman of 30, who had had three children, said she had been ill five weeks. For the first fortnight she had been in bed and actively delirious, seeing rats, devils, etc., in the room, and was treated by two medical men. She had now been out of bed for a week and was quite rational, but felt very weak and low, had some headache, and a good deal of abdominal pain (not colic); occasionally she saw double. She was thin and anæmic, showed some general tremors but no localised paresis. There was double optic neuritis and a well-marked blue line on the gums. When taxed with having taken diachylon, she said: "Yes, but that was only a pennorth, thirteen weeks ago." She obviously did not connect it with her illness. It appeared that, having gone a fortnight beyond her usual month without menstruating, she, at the advice of another woman, took the diachylon, and a week later passed some clots, which she thinks may have been a miscarriage. Since then the menses are said to have been regular. The serious brain symptoms do not appear to have come on till seven or eight weeks later, although she gradually lost colour and flesh during the whole time.

REFERENCES.

- ¹ BRITISH MEDICAL JOURNAL, July, 1893. ² *Lancet*, July, 1898. ³ *Ibid.*, September, 1898. ⁴ BRITISH MEDICAL JOURNAL, December, 1899.

A CONVERSAZIONE will be held at King's College, London, on the evening of July 3rd, when the prizes will be distributed by Professor A. V. Dicey.