the impairment caused by the adenoids as was thus immediately removed by the operation could not have been dependent upon diminished intratympanic air tension depression of the membrane and consequent depression of the ossicles; owing to the perforation of the membrane these factors would not be affected by the operation and it is evident that another cause is at work.

The foregoing is part of a paper which in addition contained a criticism of the stenosal theory and a statement of considerations which might be useful when seeking an explanation of the difficulties which had been raised.

Mr. JOBSON HORNE said it was still within his recollection, and no doubt within that of others present, that discussion on a similar paper had taken place last year in the section at Ipswich. On that occasion he, together with others, expressed the opinion that the immediate improvement in hearing was in all probability due in a great measure to the sudden local depletion following the operation. It seemed to him that the present discussion had come round to the point where they left off last year, and that no material advance had been made in the elucidation of the problem.

A SIMPLE AND PORTABLE SPRAY PUMP FOR DISINFECTION.

By W. LESLIE MACKENZIE, M.A., M.D., D.P.H.,
Medical Inspector to the Local Government Board for Scotland.

My problem was to find a portable, facile, and inexpensive spray pump. For such a pump the demand has been obvious ever since disinfection by spray became a common method in Britain. By portable, I mean light enough to be carried easily on a bicycle, which is a usual mode of transit for county sanitary inspectors or disinfecting officers. For the town areas portability is not of the same urgency, because several modes of conveyance are usually available; but even in towns, portability means a saving of labour, money, and time. By facile, I mean capable of being worked by a single man without inducing such rapid fatigue as might tempt him to scamp his work. By inexpensive, I mean cheap enough to be within the ordinarily permissible outlays of minor local authorities, or hospital committees, or School Boards, or model lodging-house keepers, or hall owners, or owners of private hospitals, or, generally, of all such as would readily use a simple disinfecting apparatus if they had it.

In the spray pumps shown in the photographs, the three conditions are, I think, reasonably fulfilled. As to portability, the small pump weighs 5 lbs., the larger (the bulb, by the way, is not essential) weighs under 6 lbs.; with crosspiece added the weight does not exceed $5\frac{1}{2}$ lbs. or $6\frac{1}{2}$ lbs. This load any boy can manage. As to facility, either pump can be worked with ease by one hand. To keep the spray nozzle playing requires, in the heavier pump, only some three up strokes and three down strokes per minute, or an equivalent of this in short strokes; in the lighter pump perhaps one or two more strokes are required. In either pump, short—say, 2-inch—strokes to the number of some 25 per minute are enough. Twenty-five short strokes or 6 full strokes per minute, that is, 1 every two or 1 every ten seconds approximately, cannot be regarded as hard work. As to inexpensiveness, the prices will be found to fulfil the indications I have mentioned.

How are these conditions secured? By the adaptation of a continuous current garden pump to the particular purposes of spray disinfection of surfaces. The essential pieces are three: a pump with an adjustable stirrup, a tube, partly flexible, partly rigid, and a spray-forming nozzle.

The Pump.

The pump is a strong brass pump, about $1\frac{1}{2}$ inch in diameter, about 2 feet in length, provided with a ball-valve inlet at the bottom, a ball-valve piston, a hollow piston-rod, and an outlet near the top. The pump is, in fact, a well-known force pump with an air chamber in the piston rod to keep the flow perfectly continuous over the "dead" points of up and down stroke. Continuity of flow with intermittent stroke could be secured by the elasticity of the flexible tube alone,

but the life of such a tube would be shorter, and the tube itself would necessarily be weaker. The continuous current in this pump is independent of the elasticity of the tube, which, consequently, may be made very strong and very inelastic. Further, an intermittent spray could be obtained from an ordinary garden syringe, but the value of the instrument for ordinary disinfecting work would be much less. Continuity of spray, therefore, is essential. In this pump, both upstroke and downstroke yield a current; both on upstroke and on downstroke the air locked in the piston is compressed; both on upstroke and on downstroke the recoil of the air cushion steadies the flow and maintains it in perfect continuity. As the pump inlet is large relatively to the pin-hole outlet of the spray nozzle, the strokes required to keep up the flow are very few, and the liquid can be delivered under very high pressure, the spray being finer when the pressure is high. The pressure, however, required for ordinary work is very little. The spray begins to play with the second stroke. No adjustment of stop-cocks, or indeed of anything, is necessary. The pump, once fitted with tube and nozzle, yields its spray. What, however, is essential is that the passages be kept clear, and the piston and piston-rod well oiled. The mechanical knowledge necessary to work the pump should be within the capacity of any man with two hands and a foot.

Fig. 1.—Spray pump; left-hand figure, with single nozzle; right hand, tube coiled for transit.

The Tube.

The tube is partly flexible, partly rigid. The flexible part is simply a piece of strong bicycle pump (or inflating pump) rubber tubing; rather, I should say it is three pieces jointed to make a length of 7 feet. Needless to add, this tube stands very high pressure; it is light, and it is as flexible as the case requires. Ordinary garden-hose tubing would serve the purpose, if lightness and convenience of packing were no object. The brass part is 3 feet long; it can be unscrewed from the rubber tube, and further unscrewed into two lengths of 18 inches each. This, as shown in Fig 1, is for convenience of packing; the two pieces of the rigid tube are seen inside

the coils of the rubber tube. In Fig. 2, the rubber tube is somewhat thicker, but admits of the same or an equivalent mode of coiling. A thin brass tube, however, is difficult to grasp firmly for any length of time. Accordingly, at the junction of brass and rubber tubes, I have had a thick wooden handpiece adjusted. With this handpiece, the brass tube can be directed with perfect nicety, and the weight of the whole tube—flexible and rigid—even when charged with liquid, is trifling. These points are all important in practice, the raised arm being very readily fatigued.

On the principle of the sweep's brush, the brass tube can be lengthened by the intercalation of extra 18-inch lengths. For high ceilings this is important. The tube, when thus lengthened to 6 feet—that is, by adding two extra lengths—is still easily manipulated. Few ceilings are more than 13 feet high. If we assume the average man's height to be $5\frac{1}{2}$ feet, and his reach with stretched arm $6\frac{1}{2}$ feet, then, with a 6-foot brass tube he will be able to come within 6 inches of a 13-foot ceiling; but if he come within a foot, that is enough. He can, therefore, without scaffolding of any kind, spray a high ceiling. The pump seems to work rather better with the longer tube than with the shorter. In buildings, such as halls, where ceilings are very high, stronger and longer tubes could be adapted to any reasonable height. The chief labour in spraying such places is due to the provision of scaffolding, shifting of steps, and the like. With the "sweep's-brush handle" tube, all this is avoidable.

Fig. 2.—Spray pump with 3-nozzle crosspiece.

The Nozzle.

The spray-forming nozzle shown in the photograph is one in common use for spraying delicate plants. The pump usually associated with it is a low-pressure air-pump. On attaching the nozzle to the pumps described above, I found that it yielded a copious, delicate, and constant spray. The nozzle itself consists of two parts, which are easily unscrewed for cleaning when necessary. As indicated already, the energy necessary to produce the spray is very small. The nozzle is designed to give a spray of a certain tenuity; but

one nozzle can, in a few seconds, be substituted for another and nozzles can be made in various grades—from coarse to fine. For all ordinary purposes the spray produced by the nozzle shown is perfectly satisfactory. The mistake commonly made is to use too fine a spray, in case wall papers or other surfaces may be destroyed. Elsewhere I have shown that this fear is due simply to *a priori* ideas about wetted surfaces. The vast majority of walls—papered, sized, white-washed, varnished, painted, etc.—can be thoroughly wetted without suffering any damage, provided the wetting proceeds from below upwards, not from above downwards.

Three-nozzle Crosspiece.

For ordinary house disinfection a single nozzle is enough; but where wall surfaces are very large, as in hospitals or halls or schools, the labour may be much reduced by the three-nozzle crosspiece shown in Fig. 2. For convenience of packing I have had the crosspiece made in detachable pieces of 18 inches each (like the pieces of the handling tube), and I have had attached three nozzles, one at each extremity and one in the centre. The capacity of the pump, I find, is more than sufficient to make all three nozzles play uniformly. The spray clouds produced just intermix at about 1 foot from the nozzles, and the combination yields a spray that may be made to wet an area of about 6 feet by 3 feet at a time. With a gentle waving motion the area can be much extended. As the amount of liquid discharged is at least three times that discharged by a single nozzle, a large area can be covered very rapidly. A ward wall 48 feet long could be taken in eight vertical sections. The spray could be moved upwards at the rate of, say, one foot a second. A section 13 feet high would thus take thirteen seconds, or say fifteen. Let this be doubled for contingencies, and quadrupled for shifting of plant. Each section would then take a minute. The eight sections would take eight minutes, or say ten. If the opposite wall take other ten minutes, the ends five each, and the ceiling (say 25 feet broad) twenty, the total time would still be only fifty minutes, a small allowance for a ward of 15,600 cubic feet, or approximately 8 beds at 2,000 cubic feet per bed. Where two persons are available a ward could easily be sprayed in this time or less; and where formalin, or other solution of formaldehyde is used, time is not unimportant to the comfort of the operators.

For these and similar reasons the three-nozzle crosspiece, which is still easily manipulated by one hand, should, I think, result in economy of time and labour. The extension of the number of nozzles is limited only by convenience and the capacity of the pump. For habitual use the crosspiece should be fitted with a stronger and heavier handling tube, but for the occasional use of the crosspiece the handling tube shown is sufficiently strong.

The Stirrup.

The vessel used for mixing the disinfectant may be an ordinary milkpail or pitcher or basin. A vessel of special capacity could easily be designed, but every house has something that will suit. To keep the pump rigid in the vessel a Z-shaped stirrup has been fitted on the cylinder. The upper arm of the Z is high enough to clear the edge of an ordinary milk pail, and if a higher vessel is used a block under the lower arm of the Z will make this equally suitable. On this lower arm the foot is placed. With the upstroke, foot and hand pull in opposite directions without moving the cylinder.

Preparation of Disinfectant.

The disinfecting officer should always be provided with a measure for the disinfectant to be used. A measure for the quantity of water, however, is likely to be clumsy. The pump itself may be used as a substitute—so many full strokes (with nozzle removed) yielding, say, a gallon. There need thus never be any chance work in making up the proper strength of disinfectant.

Packing.

The tubes are both detachable; the rubber tube can be coiled round the brass tube, or a case may be made for pump and tubes together. The pump should always be pumped clear before being packed.

Conclusion.

The spray pumps shown in the photographs have been made by Messrs. Benton and Stone, brassfounders, Bracebridge Street, Birmingham, who have carried out in a practical model all the suggestions I made to them for the adaptation of their garden sprays to the purposes of wet disinfection. In the *Sanitary Journal* for March, 1897, and in *Public Health* for March and May, 1900, I discussed in considerable detail the method of disinfection by spray. The pump now described will, I think, fulfil all that can be legitimately claimed for spray disinfection, and I say so in the light of five years' experience of the method in the ordinary routine work of a medical officer of health. There are, no doubt, many varieties of spray producer in the market, and experience will continue to suggest improvements. All I claim for the spray pump described is that it is light, easily worked, and cheap.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, ETC.

CASE OF STRANGULATED PERINEAL HERNIA.

B. S., aged 44, a native officer of Sikh cavalry, a very tall, stout heavy man, and a noted rider and cyclist, returned quite well to his headquarters (Deoli) on October 16th, 1897, having completed a ride of 109 miles on horseback.

On October 19th he noticed a hard, tender lump the size of a horse bean to the right of the centre of the perineum, and on October 26th he took to his bed and treated the swelling with fomentations.

On October 30th, when I saw him, the whole of the right ischio-rectal fossa was swollen and hard, the swelling implicating the cellular tissue of the perineum generally, and extending to the right buttock. On the same day an incision was made by the right side of the anus. Some dirty thin pus, together with liquid faeces, escaped from the wound, whilst a quantity of black, necrosed, stinking omentum prolapsed. Rectal examination revealed nothing wrong with the lower gut itself, but a hard mass was felt in the pelvis to the right. There was no abdominal swelling, pain, or tenderness. The bowels had been moved regularly every day, and there had been no symptoms indicating strangulation. The temperature was 100.6°, and the pulse 100.

On November 3rd the tongue was brown and furred, and the breath foul. The bowels continued to be opened naturally, though a little liquid faeces still escaped from the wound, indicating leakage from the intestine. The gangrenous omentum was removed with scissors. The parts were cleaner, but still emitted a horrible smell. The temperature was 101°, the pulse 100 and feeble. There was slight delirium at night. Brandy was administered at intervals.

On November 5th more necrosed omentum was removed from the wound, from which, after November 7th, no more liquid faeces escaped. The parts then gradually cleaned. Two deep sinuses remained, one passing backward and upwards between the rectum and the bulb on the right side, the other passing upward for 3 inches behind the rectum. The skin was also undermined for 3 inches forwards in the direction of the symphysis, and this part being laid open on November 27th soon filled up and healed over. The deep sinuses were reserved to be dealt with later, but the patient refused further operative treatment, and went off on leave to his home in the Punjab. I now (March, 1901) hear that he has since continued regularly to perform his duties, riding at the head of his troop, though he has still a small sinus (or fistula) behind the rectum, which he says gives him little inconvenience, and for which he declines any further surgical treatment.

Udaipur.

H. R. WOOLBERT, M.B., F.R.C.S.

THE STAINING OF THE TUBERCLE BACILLUS.

WITH reference to the note by Dr. Macleod in the *BRITISH MEDICAL JOURNAL* of July 6th, 1901, I should like to point

out that the method may be still further shortened by not using a coverglass at all.

So soon as the film has been stained in the ordinary method with the carbol-fuchsin and methylene blue and dried, a drop of cedar-wood oil or castor oil can be put on to the specimen without the intervention of the coverglass and examined with a $\frac{1}{2}$ -inch power, giving quite as good results, to say nothing of the time and expense saved. Personally I have used this method for two years.

E. MELLO SAUNDERS, M.R.C.S., L.R.C.P.

King's College Hospital.

FOREIGN BODY IN THE LEFT NASAL DUCT.

ON June 20th, 1901, a Singalese woman, aged 50, consulted me for a left fistula lachrymalis of fifteen years duration.

I slit the canaliculus and attempted to pass a probe down the nasal duct, but failed. I then thrust a knife down the duct, and kept the stricture open by daily probing until June 26th, when I introduced a style of lead.

On June 28th she again came up to see me as the pain and discharge continued, and the swelling had increased in size. I removed the style, and saw protruding through the canaliculus a fine thread, which looked like a shred of tissue. This I seized with a forceps, and to my surprise pulled out a roll of cloth $\frac{3}{4}$ in. in circumference, $1\frac{1}{2}$ in. long, and $\frac{3}{4}$ in. wide. On questioning my patient about this piece of rag, she said she had no knowledge of its existence there, but said that she had consulted a Bengalese "doctor" a year ago, who opened a swelling in the position of the present fistula, and he might have introduced the rag. No doubt her explanation is correct.

Recovery rapidly took place after this, and on July 1st, the last time I saw her, the fistula had healed and the passage down the nasal duct was free.

A. J. McClosky, M.B., C.M. Edin.

Selangor, Federated Malay States.

District Surgeon.

REPORTS

ON

MEDICAL AND SURGICAL PRACTICE IN THE
HOSPITALS AND ASYLUMS OF THE
BRITISH EMPIRE.

HORTON INFIRMARY, BANBURY.

SPONTANEOUS CURE OF AN UNUSUALLY LARGE
INTUSSUSCEPTION.

(Under the care of A. H. BOISSIER, L.R.C.P.)

MRS. R., aged 23 years, was admitted on February 20th, 1901, in a state of semi-collapse, having been sent from a village some seven miles distant.

History.—She stated that a month earlier she had suddenly been seized with severe pain over the abdomen generally. She did not remember vomiting at the outset of the pain, but the next day vomited several times a green, slimy, material, but no food or faecal matter. Attacks of pain and vomiting, each lasting ten minutes or a quarter of an hour, occurred several times daily during the month. She had passed motions nearly every day during the month, but was not clear as to the passage of blood on any occasion. She had been able to take light nourishment during the whole time.

On the morning of admission she had the worst and longest attack of pain, but it immediately ceased on the protrusion of a mass from the anus while straining at stool. She had been married two years, had had one child and no miscarriages. Four or five years earlier she had had an attack lasting three weeks of "gastric" fever for which she was treated in a hospital.

Condition on Admission.—She was quite conscious and free from acute pain. The pulse was feeble and thready; the tongue coated, but moist; the lips dry and cracked. The abdomen showed no marked distension, but was somewhat tympanitic generally. In the left ilio-lumbar region was a hard mass slightly longer than a kidney, immovable, painful on pressure, and dull on percussion. The uterus was enlarged and easily felt, suggesting a pregnancy of about three months (amenorrhoea had existed for some weeks, but no definite period could be given). Protruding from the anus was a twisted membranous-looking mass, thickly coated with faecal matter, which on gentle traction was seen to be continuous with a flat membranous cord suggesting collapsed intestine; further traction on this flat cord brought it away free from the anal orifice without pain. The rectum was digitally explored and found to be full of soft faecal matter. The uterus could also be felt enlarged and hard. Subsequent vaginal examination confirmed the impression of pregnancy.