be permanent. He tried also magnesium, which is absorbed by the tissues.¹

I myself should think an objection to the use of any absorbable metal is that it would not be everywhere removed simultaneously. If, therefore, movements were commenced too soon, a part of the magnesium would be still present, and would be a foreign body, possibly worked up into a lumpy one in the joint. If, on the other hand, movements were postponed, adhesions would form in those parts of the joint where the magnesium was first absorbed. Moreover, the precise time when absorption of the magnesium would be finished would be uncertain.

The slight crumpling of my gold plate was, of course, due to the cicatricial contraction in the size of the joint cavity during the two months the gold was buried and the elbow fixed in plaster-of-paris. I think one month's fixation would have sufficed.

In order to facilitate the removal of the plate I would recommend the use of a rather thicker sheet. The technical description of that used is "fine gold plate, gauge 0.50." I would recommend the use of gauge 0.75, and that the gold plate used should be carefully weighed before and after removal. I should add that Messrs. Wellby, to whose assistance and courtesy, given to me through Mr. Evans of that firm, I am greatly indebted, take back and allow for all the gold plate not required by the surgeon.

I am convinced that the procedure I here describe is practicable and useful. Unfortunately its value is limited by the fact that, in tuberculosis, which causes so large a proportion of stiff joints, it is often best, for weighty reasons, not even to try to get a movable joint.

REFERENCE.

- ¹ An admirable abstract of Chlumsky's paper is given in the *EPITOME* of the *BRITISH MEDICAL JOURNAL*, January 14th, 1901, par. 24.

AN OBLIQUE PELVIS ASSOCIATED WITH A CONGENITAL DISLOCATION OF THE HIP- JOINT.

By LLEWELLYN PHILLIPS, M.B., B.C., F.R.C.S.

THE specimen described and figured is of interest, as it shews the changes in the articular surfaces, the result of congenital dislocation of the hip, of a higher grade than are usually encountered, together with concomitant pelvic changes.

It was obtained from the body of a female of very short stature sent for dissection to St. Bartholomew's Hospital.

Her back showed well marked scoliosis and kyphosis. She was an elderly subject, and her hymen was intact.

Fig. 1.

The right hip joint is quite normal. The left acetabulum is

flattened (Fig. 1) and can be divided into three parts, a lower, abutting on the obturator foramen and corresponding to the non-articular portion of the normal bone, is rather depressed and extends upwards in a cave-like manner beneath the upper articular portion. Above this first part there is an area which articulated with the remains of the neck of the femur; it is eburnated in part. Below and behind is another area which was in contact with the remains of the small trochanter; it is neither eburnated nor covered with cartilage. These two latter surfaces have been formed by the flattening of the margin of the acetabulum. At the seat of the upper articular surface the bone is thicker than on the other side, but the non-articular is thinner.

The upper extremity of the femur (Fig. 2) has lost all trac

Fig. 2.

of the head, but there is a portion of the neck remaining, which slightly overhangs and is eburnated where it was in contact with the os innominatum. The small trochanter is almost unrecognisable, being flattened and elongated from above down.

The surfaces of the two bones shew that the thigh was flexed, adducted and slightly internally rotated, as well as shortened.

There was a joint capsule, and the lower and anterior acetabular surface was covered by fibrous tissue, but the other two portions were in direct contact with the femur.

Fig. 3.

The pelvis is markedly asymmetrical, the right half being better developed than the left. The asymmetry is especially

noticeable in the ilium, the left being smaller and more vertical than the right. This is well shown in Fig. 3, the asymmetry shown there being real and not due to foreshortening as might be supposed on first sight. The right sacro-iliac articulation is superficially ossified in front. The condition of the left is uncertain, as the bones had been separated during dissection. The brim is markedly oblique, the symphysis lying opposite the right anterior sacral foramina. The curve of the brim is greater on the right side than on the left. The coccyx is ankylosed to the last piece of the sacrum. The sacrum is almost flat.

That this is not an old case of tuberculous disease of the hip, the entire absence of adhesions and ankylosis is evidence, and the normal condition of the bones immediately around. That it is not due to osteo-arthritis, the total absence of osteo-arthritic changes, save in the eburnation of the neck and corresponding surface of the os innominatum, is witness. The remaining joints of the lower extremity were quite normal. It may have been due to traumatic dislocation early in life, but the changes, being those of congenital dislocation in an exaggerated degree, mark the specimen as such.

Hoffa figures several advanced specimens, but none so extreme.

The following are the principal measurements of the pelvis:—

Brim.—Conjugate diameter, $4\frac{1}{2}$ inches; transverse diameter, $4\frac{1}{2}$ inches; left sacro-iliac synchond. to right iliopect. emin., $4\frac{1}{2}$ inches; right sacro-iliac synchond. to left iliopect. emin., $4\frac{1}{2}$ inches; outlet, antero-posterior diam. (from lower end of sacrum), 4 inches; transverse (middle of tubera), 3 inches.

Bistiliac.—Outside measure, $8\frac{1}{2}$ inches, inside measure, $8\frac{1}{2}$ inches; ant. sup. spine to symphysis, right side, $4\frac{1}{2}$ inches; left side, $4\frac{1}{2}$ inches; ant. sup. to a spot opposite middle of sacrum, right side, $5\frac{1}{2}$ inches; left side, 4 inches.

Sacrum.—4 inches long by $3\frac{1}{2}$ inches broad at its widest part.

In the Museum of St. Bartholomew's Hospital there is a somewhat similar specimen. Such is described as "repair after hip disease," but that is merely conjectural, as there is no history attached to it.

AN OUTBREAK OF EPIDEMIC CATARRHAL JAUNDICE IN DERBYSHIRE.

By HERBERT PECK, M.D., D.P.H.,

Medical Officer of Health for the Rural District of Chesterfield and the Borough of Chesterfield.

THE outbreak began in the hamlets of Speedwell and Poolsbrook, in the parish of Staveley, in the Rural District of Chesterfield, in the first week of August, and has continued to the time of writing. Three of the medical men practising in Staveley then had cases of jaundice, which they regarded at first as being of the simple catarrhal variety, but these gradually increased in number, and they gradually became impressed with the idea that the disease was of an infectious nature.

Speedwell and Poolsbrook are colliery villages belonging to the Staveley Coal and Iron Company. They are well drained, and have roads paved with furnace slag and engine ashes. The conservancy system is in vogue, and the privy ashpits, 35 to 46 feet from the houses, are regularly emptied by the company's servant. Each house has a privy, two or four of which empty into each ashpit. The houses in Speedwell are 175 in number, are built of brick, and are in good repair, but have only one door, though through ventilation is obtainable by the back windows. Poolsbrook is more modern than Speedwell, and contains 213 houses, a church, school, public-house, and several shops. The houses are in excellent order, and have each two doors. The water supply of both is upland surface water from the Crowhole and Ramsley reservoirs, but owing to drought the supply has been almost entirely drawn from the latter. The water from Ramsley reservoir has a slightly peaty character and more colour than that from Crowhole. The greatest proportion of Ramsley to Crowhole water during the summer was supplied on and between July 28th and August 1st, when the water had a distinctly yellow colour, which fact gave rise to the popular, but erroneous, opinion that the colouring matter of the water was the cause of the outbreak of jaundice which followed soon after.

Dr. Arthur Court was the first to inform me of the outbreak. This he did on September 29th. This gentleman and

Drs. Badocock and Blight most obligingly furnished me with lists of their patients suffering from the malady, but unfortunately these were not complete, owing to the fact that many of the patients suffered so little inconvenience that they attended at the surgeries for treatment, where their names were not recorded. I supplemented these lists by my own inquiries at the houses of patients, who often informed me of cases of which I had not previously heard.

I visited altogether 53 houses where cases had occurred, and found 3 households with 3 patients each, 9 with 2, and 42 with 1 patient each; 50 of the patients were males and 19 females. The youngest patient was a child of 16 months, nursed by a jaundiced mother, and 2 other young patients were respectively aged $2\frac{1}{2}$ and 9 years. The oldest patient was 70 years of age, and 2 others 50 years and a few months. The following are the age-periods of the patients:

Between 1 and 5 years...	...	...	2
" 5 " 10 "	...	...	1
" 10 " 15 "	...	...	2
" 15 " 20 "	...	...	8
" 20 " 25 "	...	...	16
" 25 " 30 "	...	...	17
" 30 " 35 "	...	...	9
" 35 " 40 "	...	...	8
" 40 " 45 "	...	...	2
" 45 " 50 "	...	...	1
Upwards of 50 years "	...	...	3
Total ...	...	...	69

I think this number represents quite half the cases which have occurred up to the present.

SYMPTOMS.

The onset of the disease was usually gradual, with *malaise*, want of appetite, and in many cases headache. Drowsiness was a frequent symptom. There was almost always abdominal pain, sometimes very severe; and cramps in the calves of the legs occurred in many cases. After a variable period, usually two or three days, but sometimes longer, the patient vomited, but there were several exceptions to this. Diarrhoea was only occasionally present. When the temperature was taken at this stage, it was found to be elevated, being usually between 100° and 101.5° F., but a temperature of 104° was observed in at least one case. A distinct rigor occurred in several cases, and there was a history of chilliness and shivering in all.

Jaundice usually came on a day or two after the vomiting, but was delayed for a week in several cases, though, on the other hand, slight jaundice and yellowness of the conjunctivæ was sometimes observed before the vomiting, but became more marked after it. When the temperature was taken at this stage (jaundice), which was the one at which the greater part of the patients were first seen by a doctor, it was found to be normal. No elevation was observed after this in uncomplicated cases. Pruritus was a frequent symptom. Tenderness of the liver was present in most cases, but enlargement less frequently.

The spleen was frequently found enlarged. Nephritis was found in only one case. The only complications observed beyond the case of nephritis were two cases of acute croupous pneumonia, two of urticaria, and one of acute yellow atrophy. A *post-mortem* examination of the patients who died from the last-named disease was made by Dr. Court, and a microscopical examination of her liver, spleen, and kidneys, by Professor Arthur Hall, of University College, Sheffield. The conditions found pointed to the death being due to acute yellow atrophy, this was probably accelerated by epidemic catarrhal jaundice.

ILLUSTRATIVE CASES.

The following cases will illustrate the usual history:

CASE I.—J. M., aged 34. "I had a cold and thought I had influenza, for I ached all over—every bone in my body. This was on Wednesday, and I got worse next day and was jaundiced on Friday."

CASE II.—G. S., aged 27. "I had a cold and ached all over. Then I vomited and was jaundiced afterwards—two or three days afterwards."

CASES III AND IV.—E. R., aged 33, was ill and then jaundiced. He vomited on Saturday and was jaundiced on Monday. W. S., aged 39, living in the same house as the last, began to vomit on the following Sunday and was jaundiced on the Tuesday.

CASE V.—J. J., aged 30. "I started with influenza on September 20th and vomited; jaundice came on gradually a week after."

CASES VI AND VII.—S. S., aged 24. "I had a chill and perspiration in the pit on September 20th and was jaundiced the same day. I did not vomit." S. A. P., aged 13, living in the same house as the last, took ill