with these organisms in their nasal or aural discharge are capable of giving rise to diphtheria in others. On the other hand, it may be argued that diphtheria bacilli, as with other organisms, may be virulent to man and not to animals and *vice versa*, so that these 11 cases, which after all are in the majority, may have been virulent in their original state.

No. 20, though proved to be a virulent Klebs-Loeffler bacillus, gave no growth or acidity in sugar broth, which suggests that this test, on which so much stress has been laid by some writers, is not infallible.

CONCLUSIONS.

In conclusion, to enumerate a few points which this only too short investigation appears to indicate:

1. That it is advisable to cultivate all cases of rhinorrhœa—apart from that of the acute stage—and otorrhœas in scarlet fever cases, especially in hospital practice.

2. That bacilli when found at all resembling the diphtheria bacillus, must in the present stage of our bacteriological knowledge be regarded as a modified variety of that organism, bearing in mind that their staining properties are often the only means of diagnosis available, the clinical symptoms in this class of case being for the most part absent. For other proofs of the unsatisfactory state of our knowledge we can refer to the writings of Stamm,² Meyer,³ Freeman,⁴ Abbott,⁵ and others, who found organisms indistinguishable from that of diphtheria in rhinitis fibrinosa, similar results by J. W. Stephens and Kanthack in cancer oris, and again by Cautley⁶ in catarrhal pharyngitis.

3. That systematic isolation of these rhinorrhœas and otorrhœas is not only justified but advisable.

4. That such isolation may be reasonably expected to reduce the post-scarlatinal diphtheria incidence.

5. That it is an open question whether such mild cases require antitoxin treatment. The two cases, however, referred to in the early part of this paper, in which the rhinorrhœa was followed by the deposit of membrane upon the fauces, seems to indicate the use of the serum; at least it may do good, and can do no harm. Even assuming that many of these cases unassociated with general symptoms have not the disease themselves, but have had the bacillus simply grafted on to an ordinary coryza or cold, or on to an existing rhinorrhœa or otorrhœa, none the less it is highly important that they should at least be separated from the healthy. It is a generally accepted fact that healthy people may carry about diphtheria bacilli in their throats, and produce in others a severe attack of the disease; examples of this are given by Airy,⁷ the late Sir Richard Thorne,⁸ Newsholme,⁹ and others; similarly the otherwise healthy subjects of nose or ear discharge containing diphtheria bacilli may reasonably be held to be pathogenic to others.

6. That these discharges, unassociated with sore throats or symptoms, and therefore easily overlooked, may be the cause of the often unaccountable outbreaks, and the persistence of the disease amongst school children.

And, finally, I wish to thank Dr. Hume, the Medical Superintendent of the hospital, for giving me every possible facility for the conduct of my investigation; and Dr. Cartwright Wood for determining the pathogenic powers of the bacilli; without the facts elicited by his animal experiments my observations would have remained without the positive value which I venture to think they now possess.

REFERENCES.

- ¹ *Zeitschr. f. Hyg.*, xxiv, 1897, p. 443. ² *Archiv für Kinderheilkunde*, Band xiv, 1892, p. 157. ³ *Archiv für Laryngologie und Rhinologie*, Band iv, 1896, p. 249. ⁴ *Medical Record*, New York, vol. i, p. 618. ⁵ *Medical News*, May 13th, 1893. ⁶ Supplement to the Twenty-fourth Annual Report of Local Government Board, p. 455. ⁷ *J. Epidem. Soc.*, vol. iv. ⁸ *Diphtheria Natural History and Prevention*, p. 57. ⁹ *Epidemic Diphtheria*, p. 138.

PNEUMOCOCCUS ARTHRITIS WITH NOTES OF SEVEN CASES.

By NATHAN RAW, M.D., M.R.C.P.LOND.,
Physician, Mill Road Infirmary, Liverpool.

THE microbic origin of acute lobar pneumonia has met with such general acceptance by physicians in all parts of the world that it is unnecessary here to mention the many arguments in favour of it. Acute lobar pneumonia must now be

classified amongst the specific infectious diseases depending entirely on the presence of its specific micro-organism, the diplococcus pneumoniae of Fraenkel. Up to a short time ago, the action of this organism was thought to be confined to the lung tissue, but recent investigations have clearly demonstrated that the micrococcus pneumoniae is capable of being conveyed by the blood stream to various parts of the body, and setting up most acute and serious lesions.

These lesions may either precede the actual onset of pneumonia or may occur either during or after an attack. The lesions which have been accurately observed as being caused by the pneumococcus are as follows:

1. Fibrinous lobar pneumonia.
2. Pneumococcus arthritis.
3. Pneumococcus abscess.
4. Pneumococcus cellulitis.
5. Pneumococcus peritonitis.
6. Pneumococcus empyema.
7. Pneumococcus meningitis.
8. Pneumococcus pericarditis.
9. Pneumococcus otitis media.
10. Pneumococcus salpingitis.
11. Pneumococcus endocarditis.
12. Pneumococcus keratitis.

The exudation might be purulent from the first, sero-fibrinous throughout, followed by suppurative or hæmorrhagic.

In a later paper I hope accurately to describe many of the above-mentioned complications which have occurred in my own practice; but I want to confine my attention now to the action of the pneumococcus in the joints, setting up either a simple synovitis or a violent suppurative arthritis requiring immediate surgical treatment.

The occurrence of arthritis during an attack of pneumonia had been noted by several observers long before the specific organism had been isolated, but they were considered to be merely coincidences, and not in any way directly connected with the attack of pneumonia.

A very remarkable monograph was published by Dr. Leroux of Paris in 1899, in which he tabulated some 28 cases of pneumococcus arthritis occurring in the practice of Continental physicians. Dr. Cave of Bath also published an admirable paper in the *Lancet* of January 12th, 1901, recapitulating these cases and adding more, bringing the total of authentic cases up to 31, and minutely describing 2 cases which had occurred in his own practice.

It has been a matter of surprise to me during the last few years that more cases have not been recorded in this country, and I can only attribute it to the fact that general attention had not been directed to the subject. I feel sure that cases will now be properly classified and opportunities given for more extended observation. It would appear from the papers of Drs. Leroux and Cave that arthritis was a very rare complication of acute lobar pneumonia; but my experience is that it is not so rare as we would be led to believe, and those of us who have the opportunity of seeing a large number of cases of pneumonia will, I feel sure, be able to support this view.

During the last four years I have had under treatment in Mill Road Infirmary, Liverpool, an institution containing over 800 beds, no fewer than 817 cases of acute pneumonia. As the cases occur amongst the very poor population, many of whom are addicted to alcohol, the mortality has been high, averaging each year 24 per cent.

Of these 817 cases of pneumonia, arthritis as a complication showed itself in 7 cases: 6 of these were suppurative and 1 serous; 4 patients recovered and 3 died.

The following are very brief notes of my 7 cases:

CASE 1.—H. G., aged 28, was admitted into Mill Road Infirmary with acute pneumonia of the right lung. The whole lung was involved, and the attack was a very severe one, with considerable pleurisy. He had his crisis on the eighth day, and was apparently doing well, when on the eleventh day of his illness he had a rigor and was seized with intense pain in the right ear, accompanied by a sudden rise of temperature. Next day a swelling appeared over the second right costal cartilage with great pain and tenderness. Two days afterwards a large abscess rapidly formed over the right buttock. His condition became very critical, and his temperature high and erratic. An incision was made over the swelling on the chest, and was found to be an abscess of the right sternoclavicular joint pointing between the first and second ribs; the abscess of the buttock was drained, giving exit to a large quantity of sweet-smelling pus. The abscess in the middle ear burst the same day, giving great relief.

Very careful precautions were taken in examining microscopically and bacteriologically the pus from every source above described, as well as the sputum.

In all cases the apparently same organism the pneumococcus of Fraenkel was found in pure culture. The man made a good recovery and is well to this day. I have no doubt that in this case prompt surgical interference saved his life.

CASE II.—T. A., aged 52, was admitted on January 5th, 1900, with severe pain in right side of chest and profuse dark expectoration. He had also pain and swelling of right ankle and swelling with great tenderness over the right sternoclavicular joint. Next day he was found to have definite pneumonia at the right apex, with effusion at the base.

The joints seem to have swollen before the onset of his pneumonia and he had suffered from rheumatism on several occasions. As soon as pus was found in the joints they were freely incised and drained, both articulations being found roughened and bare. The pneumonia ran an ordinary course and his temperature fell, but later he developed an empyema which was also opened and drained. This man had a most severe general infection by pneumococci, and although repeated cultures were made from the pus and sputum no other pathogenic organisms were found except diplococci. After a long illness he made an excellent recovery, his chest quite healed, as also the sterno-clavicular joint; the ankle healed up but the joint remained stiff. I am greatly indebted to my senior assistant, Dr. John Hay, for the great care he bestowed in the treatment of this case.

CASE III.—K. F., aged 49, charwoman, admitted June 11th, 1900, with severe pain in right chest and abdomen, with all the symptoms of acute pneumonia. She was an alcoholic, and had a severe struggle until the crisis, which occurred on the seventh day. Two days after this a swelling appeared under the right deltoid muscle, with great pain on movement. A needle was inserted into the joint and pus withdrawn. I freely incised the joint, and let out 18 ounces of curdy, sweet-smelling pus. I noticed at the operation there was considerable infiltration between the muscles around the joint. Her temperature dropped and she appeared to be doing well for twelve days, when she developed empyema of the right side. This was opened and drained; then a large abscess developed in the abdominal wall, which was drained, but the patient sank into a condition of general pyæmia, and died on the twenty-eighth day. Pneumococci in pure culture were obtained in all the samples of pus and sputum.

CASE IV.—A man, aged 23, was admitted with an ordinary attack of acute lobar pneumonia of the right lung. His crisis occurred on the seventh day. Two days afterwards he developed a painless swelling of the right knee, with a temperature of 100°; a needle drew off clear serous fluid. Under the microscope, however, pneumococci were found in abundance. The joint was aspirated of 7 ounces of fluid and firmly strapped. After a slow and tedious convalescence he eventually recovered and walked out with a useful joint.

CASE V.—A man, aged 43, a confirmed alcoholic, was admitted in a collapsed condition with acute lobar pneumonia of the right lung. Two days after admission, being the sixth day of the disease, he developed a painless swelling of both knees. A needle showed sero-fibrinous effusion which on cultivation and examination was found to be a pure culture of

pneumococci. He died on the following day, and at the post-mortem examination he was found to have a most severe pneumonia with inflammation of both knee-joints. The organisms recovered from the lung and the knee-joints appeared to be identical.

CASE VI.—A man, aged 58, also a drunkard, was admitted on April 13th, 1900, with acute pneumonia of the right lung. He had his crisis on the seventh day, and appeared to be comfortable, when he was seized with great pain in the right knee and rapid distension of the joint. A needle showed sero-pus, which on being stained revealed pneumococci with well-defined capsules. Dr. Pitt-Taylor opened and drained the joint, which continued to discharge pus for two weeks and eventually healed, leaving the man a very useful joint. He was discharged recovered, but his knee was fairly stiff, although not in any way ankylosed.

CASE VII.—F. D., aged 42, tram driver, was admitted on July 12th, 1901, with acute pneumonia of the right lung. On admission he complained of great pain in his right shoulder, which was dusky-red and inflamed, but without any definite fluctuation. He died before any operation could be performed, and careful examinations of the fluid from the shoulder showed pure culture of pneumococci. In this case also an extensive intramuscular infiltration was found around the shoulder-joint.

CONCLUSIONS.

From an observation of a large number of cases of pneumonia I am convinced that the pneumococcus is capable of producing very serious lesions in other parts of the body than the lungs. It will be seen that arthritis as a complication occurred in 7 patients out of 817 cases of pneumonia—equal to about 1 per cent. Of these 7 cases 4 have recovered and 3 died. Curiously enough, my cases correspond almost exactly with those recorded by Continental observers, in that 6 were males and 1 female. In all 7 cases the pneumonia was right-sided, and the joints affected were all on the right side. This, I should say, was only a coincidence, although the fact is striking.

I have observed also in some cases of pneumonia a slight redness and pain in the shoulder-joint of the side affected which has subsided with the crisis. The joint affection may either precede the lung symptoms or follow the crisis, or develop intercurrently. In Liverpool alcohol seems to decide in a great measure the severity of the attack and the prospect of recovery, and I feel sure that one at least of the above cases would have recovered had he not been a confirmed drunkard.

The only treatment to my mind is early evacuation of the pus if it can be reached. It is just possible that many isolated cases of synovitis and arthritis occurring either during or after an attack of pneumonia—or even independent altogether of pneumonia—may be due to this organism, and I would suggest that the name, "Fraenkel's diplococcus," would be more suitable than "pneumococcus." Appended is a table of the above cases.

Table of Cases of Pneumococcus Arthritis.

No.	Date.	Sex.	Age.	Relation to Pneumonia.	Seat of Arthritis.	Nature.	Complications and Remarks.	Treatment.	Result.
1	1898	M.	28	Pneumonia 3 days before	Right sterno-clavicular joint	Suppurative	Right otitis media, abscess of thigh	Free incision and drainage	Recovered.
2	1899	M.	52	Pneumonia 2 days after	Right ankle, right sterno-clavicular	Suppurative	Right empyema, joint stiff	Free incision, arthrotomy	Recovered.
3	1900	F.	49	Pneumonia 2 days before	Right shoulder joint	Suppurative	Right empyema, extensive cellulitis of shoulder	Incision, drainage, arthrotomy, septicæmia	Died.
4	1897	M.	23	Pneumonia 2 days before	Right knee	Serous	None	Aspiration	Recovered.
5	1898	M.	51	Pneumonia 2 days before	Both knees	Serous	A very severe infection, toxæmia, alcoholism	General	Died.
6	1900	M.	58	Pneumonia 3 days before	Right knee	Suppurative	Stiff though fairly useful joint	Free incision and drainage	Recovered.
7	1901	M.	42	Pneumonia about same time	Right shoulder	Suppurative	Severe general infection, with extensive cellulitis of shoulder	General	Died.

CASE OF ANTHRAX IN WHICH THE INFECTION AROSE FROM A HITHERTO UNDESCRIBED SOURCE.

By J. CONNAL WILSON, L.R.C.P., L.R.C.S. EDIN.,
Haworth, Keighley, Yorkshire.

THE following case is, I think, of sufficient interest to have it placed on record.

J. W. W., aged 26, "picker" maker, consulted me in the surgery on the evening of July 1st, 1901, his chief complaint being directed towards a spot on his left cheek, slightly lower than the angle of the mouth. As there appeared more swell-

ing and induration than were consistent with such an apparently trivial cause, his temperature was taken and ascertained to be 101.8° F. He was ordered home to bed, placed upon Listerine internally, and when seen next morning had a slightly higher temperature, a greater amount of swelling involving all the glands of the head and neck, and extending down to the clavicles and top of the sternum. At my request he was seen by Dr. Bell of Bradford, who agreed with me that the case was one of malignant pustule, and advised speedy removal of the infected area, which was perhaps not markedly characteristic, but from which were obtained capillary tubes of serum, these being proved to contain the bacillus of anthrax when examined microscopically the same evening. The operation was performed in the course of a