3. *Enucleation of Muscle from its Sheath Outwards.*—The inner divided edge of the sheath is picked up with dissecting forceps, and, by means of the handle of the knife and the forefinger, the inner part of the muscle is readily enucleated from its attachments to the sheath anteriorly, internally, and posteriorly. Care must be taken that, at the level of the fold of Douglas, posteriorly and for an inch or so above, the deep epigastric vessels be carried outwards on the under surface of the muscle. In the separation anteriorly it will be found necessary, should the incision include in its length any of the transverse fibrous intersections, to use the point of the blade of the knife to free the sheath from these—a perfectly simple performance. The intersections do not penetrate to the posterior surface of the muscle.

4. *Division of Sheath (Posterior), together with Transversalis Fascia, Subperitoneal Fat and Peritoneum.*—The enucleated portion of the rectus being retracted outwards and the inner part of the anterior sheath inwards, the abdomen is opened in the same vertical plane as the previous incisions. As soon as the peritoneum has been incised a director or the forefinger is introduced, and the following structures divided, up and down, as far as necessary: in the whole length of the incision the transversalis fascia, subperitoneal fat and peritoneum, and above the level of the fold of Douglas the posterior layers of the muscle sheath.

5. *Suturing of Sheath (Posterior), together with Transversalis Fascia, Subperitoneal Fat and Peritoneum.*—To close the abdomen the above structures can be readily united by a continuous suture, the necessary parts being retracted to help the operator. A curved intestinal needle, a needle holder that will not allow the same to turn, and fine silk being used. Care must be taken to accurately adjust the divided edges of the semilunar fold of Douglas, which is easily recognised an inch or so below the umbilicus.

6. *Suture of Sheath (Anterior).*—The rectus, after the posterior structures have been united, is allowed to slide back into its natural position, thereby completely covering the line of suture. The anterior layers of the sheath are united by a similar continuous suture.

7. *Suture of Skin and Fat.*—This is performed in the ordinary way. Although this description may seem somewhat long and tedious, the actual performance, so far as the opening of the abdomen goes, will be found to be extremely simple, and to take little, if any, longer than any other method. The incision can readily be extended, if necessary, by slitting the anterior sheath with the extreme point of the knife, enucleating the muscle with the finger, and dividing the posterior structures with scissors. As regards closing the abdomen the parts come together so accurately and readily, and it is so easy to see what one wants to do, that the continuous suturing can be completed with great rapidity.

Directly the patient regains consciousness and begins to vomit the advantages of this method become apparent, for whereas the rectus under the influence of the anæsthetic has been flaccid, it now regains its normal tone and resumes its functions, unimpaired either by a row of sutures or by loss of nerve supply. In the act of vomiting it stands out, one may almost say, involuntarily, firm, and stiff, and so guards in the most efficient way possible the very area which has been injured. With every subsequent strain to which the abdominal wall is liable, whether in expelling the contents of the abdomen or pelvis, in coughing, or in other physical effort, the action of this muscle will be the surest preventive of hernia, forming at such times a tough and more or less rigid band, tending to approximate itself to its fellow of the opposite side, and so holding itself securely in its sheath, and more particularly into that part of it from which it has been separated. It must at the same time be remembered that the disturbed area presents for the formation of adhesions in the process of healing wide opposing surfaces some $2\frac{1}{2}$ inches broad (1 inch anteriorly between sheath and rectus, the same posteriorly plus the breadth of muscle internally). On several occasions I have opened the abdomen in this way *post mortem*, from pubes to ensiform cartilage, to right and left of the middle line, and closed it as described. To all appearances it was perfectly satisfactory. This was particularly noticeable when, on putting the hand into the abdomen from above through the diaphragm, pressure was exerted from behind the line of suture forwards. The thickness of the rectus could then be plainly felt in front of the fingers giving a sense of perfect security and strength to the area in question.

I used the method on the living subject for the first time on February 7th of this year, the case being one of semi-solid dermoid tumour of the ovary about the size of a fetal head. The patient was a small, thin woman, and the result so far has been entirely satisfactory; nor, although it is early days, do I see how she can develop a hernia through the scar in the future.

The operation was followed by vomiting, and one could see and feel the recti standing out equally right and left, with the skin incision over the left muscle. The patient was on a couch on the tenth, and allowed to walk on the twenty-first, day after the operation. Palpation of the parts now, five weeks from that date, when the recti are acting, assures one of the security which exists.

In looking up the subject after I had performed the opera-

tion; I found that the late Mr. Greig Smith, in his admirable book on abdominal surgery, advocates a somewhat similar procedure, but it is to take the place of incision in the semilunar line; and he recommends that the rectus be turned inwards, thereby injuring the motor nerve supply, which I particularly avoid. The sutures, also, are apparently put through the muscle substance.²

To sum up the advantages of this method, they appear to me to be:

1. Extreme simplicity.
2. Division of aponeurotic structures only, such being afterwards supported by a strong muscle entirely free from injury by sutures or division of nerves.
3. Extreme improbability of hernia.
4. Great reduction in time of the length of period at present necessary for convalescence after simple cases of laparotomy.

REFERENCES.

- ¹ *Annals of Surgery*, vol. xxxiii, No. 1, 1901. ² *Abdominal Surgery*, vol. 1, p. 109, 5th edition. Greig Smith.

CASE OF CONGENITAL HYDRONEPHROSIS : NEPHROTOMY AND DRAINAGE : RECOVERY.

By JOHN THOMSON SHIRLAW, M.B., C.M.,

Upholland, Wigan.

As cases of congenital hydronephrosis are somewhat rare, I venture to put this on record.

History and State on Examination.—On April 11th, 1900, I was called to see a male child, 6 weeks old, on account of a swelling of the abdomen, which had been present at birth, and been getting slowly larger. The child had been brought into the world by a midwife, who declared that the swollen abdomen had impeded the labour after the birth of the head. The child, when I saw it, was weakly and wasted, and had irregular attacks of diarrhoea. The abdomen presented the appearances of an abdomen with a large ovarian cyst; it was greatly distended, the distension being greater

on the left side; the skin was thin and tightly stretched, and large veins ramified over its surface. Palpation and percussion revealed a tumour exactly the shape of a kidney occupying the left lumbar, iliac, and hypochondriac regions, and left half of the umbilical region. It was dull all over, right up to the lumbar spine, and gave the vibratile thrill. It had a smooth surface and edge all round, except at the inner side, where a distinct notch could be felt. As time went on the child grew thinner, although it took nourishment greedily, and the tumour increased until it occupied the entire umbilical region, and entered the hypogastric region as well, so that the abdomen projected on to the thighs. The tumour seemed to have rotated somewhat to the right at its lower end, so that the notched inner surface lay at a slight angle to the median line. The bowels were crushed up under the liver, and lay between the tumour and that organ; one or two coils lay over the surface of the swelling. The descending colon could be felt and percussed at first over the surface of the tumour, but eventually slipped round and lay quite close to the lumbar spines. The tumour was most prominent and nearest the skin about one inch above the umbilicus, and half an inch to the left side; and into this point I inserted a hypodermic needle, and drew off some fluid, which was clear, non-albuminous, and showed nothing under the microscope beyond an occasional epithelial cell.

Operation.—The diagnosis having been made, I strongly advised operation, in the hope of establishing a fistula, and with a possible chance of complete cure. On May 2nd, accordingly, when the child was 9 weeks old, nephrotomy was done, with the assistance of my partner. There being no room for extraperitoneal drainage on account of the youth of the child and the position of the colon, but still desiring to have the drainage as far back as possible, I made a 2-inch incision midway between

the iliac crest and the last rib and parallel to the latter. I entered the peritoneum rather suddenly, being only aware of the fact when a coil of small intestine presented. To get more room the incision was carried forward another 2 inches. The sac was easily found, brought to the surface by two pairs of artery forceps, a nick made in the posterior peritoneum, and a small trocar and cannula inserted. The quantity of fluid was not measured, but there was fully a pint and a-half. It was at first watery, then mucilaginous and soapy, and streaked with white towards the end. The sac wall was exceedingly thin. A well-fitting drainage tube was then inserted, the tube invaginating a portion of the sac wall, and then fixed with a suture through the skin. A drain was also inserted into the peritoneum. The child was very collapsed after the operation, with a very slow and intermittent pulse, but it was wrapped in wool and given brandy and strychnine.

Progress.—After six hours the child was recovering from the shock, and had a pulse-rate of 170. From that time on the child did well. The temperature never rose beyond 100.5° until the fifth day, when the sinus became foul, and caused a temperature of 102°. For the first week the dressings were soaked through and through, but the drainage gradually lessened, and by October 13th, five months and a half after the operation, the sinus closed, there being no further drainage. The cavity was well washed out daily with weak carbolic lotion. Three weeks after the operation, the drainage tube was removed altogether, as a permanent fistula seemed to have been established; about this time, also, the child began to waste rapidly, and was given cod-liver oil inunctions and small doses of an oil emulsion internally. This had the desired effect.

There seems to be some confusion as to the true meaning of the word "congenital" as applied to hydronephrosis. According to Morris¹, congenital hydronephrosis is "hydronephrosis in the foetus and newborn, not hydronephrosis which occurs some time after birth, and is due to a congenital cause." Bland-Sutton,² however, has reported a case of congenital hydronephrosis in a woman, aged 44. I think there can be little doubt that my case was a true congenital one. The cause of the obstruction was not sought for. The operation had to be done hurriedly on account of the weak state of the little patient. There were no other congenital deformities or malformations, with which the condition is sometimes associated.

As regards treatment, in my case aspiration was not tried. Mr. Thornton³ thinks the chance of cure by such means very slight, but that there is a good chance of cure by nephrotomy. In my case the kidney was reached through the peritoneum for the simple reason that it was much easier, and the fluid in the sac aseptic.

The cure, in my opinion, was due to the fact that the sac had reached the utmost limit of distension, and that thereby the secreting structure had been destroyed, or almost entirely destroyed. Would aspiration, then, alone have been quite as effectual? I hardly think so, considering the very free escape of fluid for some time after the operation, followed by the discharge of purulent fluid, which meant granulation of tissue and the disintegration of the few remaining more highly-developed secreting cells.

REFERENCES.

¹ *Surgical Diseases of the Kidney*. ² *Lancet*, vol. i, 1896. ³ *Ibid.* vol. ii, 1889.

BILOCULAR INTRAPELVIC AND SCROTAL HYDROCELE.

By J. LACY FIRTH, M.S., F.R.C.S.,

Assistant Surgeon, Bristol General Hospital.

A HYDROCELE may have an hour-glass shape, constituting what the French term a hydrocele *en bissac*, either because a localised inflammatory thickening of the tunica vaginalis renders one part more unyielding than the rest, or because the tunica becomes pouched at some part from local weakness. Curling pointed out that such a pouch might form in the digital fossa between the epididymis and the back of the testicle, and might push its way towards the opposite side of the scrotum. Similarly pouches may form on the front or sides of the tunica vaginalis, and the pouches may, by reason of the relative weakness of their walls, become larger than the tunica vaginalis itself. Such a pouch may bulge far into the opposite side of the scrotum and cause a single hydrocele to resemble superficially a double one. These varieties of bilocular hydrocele are contained in the scrotum. The case of bilocular hydrocele described below is of the rarer form in which one loculus is in the scrotum and the other within the abdomen, and the orifice or channel of communication between the two lies in the inguinal canal.

W. L., a labourer, aged 20, came to the out-patient department of the Bristol General Hospital in October, 1899, complaining chiefly of a swelling in the left side of the scrotum, but also of a swelling of the prepuce and of a discharge from the penis. The latter was due to a chancre which in due course proved to be syphilitic. On questioning him closely, it was ascertained that he had had some degree of swelling in the left side of the scrotum as long as he could remember, though it had given him no serious trouble until about a month previously. Further, his father had told him that when an infant he had had a swelling there, and had been turned head downwards on two or three different occasions in order to reduce the swelling when it had become tense. The swelling always became smaller when he lay down. About a month before coming to the hospital he had had a blow with a plank on the lower part of the scrotum, and the effects had kept him in bed a fortnight, during which time a black bruise mark formed where he had been struck. At the end of the fortnight he returned to work and had to carry heavy sacks of cement.

After three days he had to give up work again, owing to pain in the left groin and scrotum. The swelling at this time was of the same size as it had been before the accident. It was then tapped by a practitioner, and a bowl of fluid, said to be like urine, was drawn off. The swelling soon came back when he returned to work, and it was again tapped in the hospital casualty room. It was after this second tapping that he came under my care. No active treatment was given to the hydrocele until March 20th, nearly five months later. By that time the scrotal swelling had increased in size, and was painful whenever he attempted to work, and these facts induced him to come into the hospital and submit to an operation. The following note of the patient's condition was made on March 3rd: As the patient lies upon his back in bed, the left side of the scrotum is seen to be enlarged to the size of a big lemon. When he coughs or strains the swelling immediately expands, returning again to its former size as soon as the cough ceases. It is very translucent. It feels like a rather flaccid sac of fluid. The testicle is felt lying in the posterior part of the swelling, and is very sensitive, so that examination of it is hardly possible. The globus minor and body of the epididymis feel indurated and nodular. The swelling can be almost entirely reduced through the inguinal canal, but quickly returns again when the pressure is relaxed. When the scrotal swelling is suddenly and smartly pressed upon, a slight bulging of the abdominal wall in the left iliac region is produced, and on laying one hand on that region and the other on the scrotum fluctuation is readily perceived. Similarly a thrill is transmitted by gently tapping either part. The area of the abdominal parietes through which fluctuation can be felt is roughly pyriform in shape. The inner border is the middle line, and extends 5 inches above the symphysis pubis and to within an inch of the umbilicus; the outer edge is at Poupart's ligament, and extends to a point 2 inches from the anterior superior iliac spine, and the upper border is a curved line, with its convexity upwards, joining the distal ends of the two borders mentioned.

A sudden tap made upon any part of the abdominal wall, even in the right hypochondriac region, immediately produces a visible expansion of the scrotal swelling. There is no visible prominence of the abdominal wall in the fluctuating area either when the patient is lying or standing. Both flanks are resonant in all positions, showing the absence of ascites. A rectal examination was made, and an attempt to obtain fluctuation between the examining finger and the other hand placed over the iliac fossa or in the scrotum, but without success. This examination was very unsatisfactory, owing to the patient's intolerance. The patient had a small and very flaccid hydrocele of the right tunica vaginalis.

Operation.—On March 20th an incision was made over the external abdominal ring and inner part of the inguinal canal, and after a little dissection the wall of the hydrocele sac was exposed and incised. Ordinary-looking hydrocele fluid flowed freely away. Through the opening and through the neck of the sac the forefinger was easily passed backwards into a large cystic cavity lying in the iliac fossa. The neck of the sac and inguinal canal would easily have allowed two fingers to pass. The intra-abdominal part of the hydrocele sac was extraperitoneal. The scrotal portion, which contained the testicle, was separated from the spermatic cord a short distance above that organ and cut across there, and the lower part closed by a few points of suture, so as to form a tunica vaginalis for the testicle. The rest of the scrotal sac was freed from the structures of the spermatic cord which lay behind it, and from other structures up to the inguinal canal, just as a hernial sac is separated in the operation for the radical cure of hernia. Gentle traction was made upon the free portion, and the remaining portion of the sac, the intra-abdominal portion, gradually came down, and was separated easily from the surrounding structures, until in a very short time it was found, rather to my surprise, that the whole of it had been removed, and the cavity where it had been had become simultaneously obliterated. The inguinal canal was closed after Macewen's method. The dressing was removed for the first time eight days after the operation. The testicle then felt enlarged and soft, and it was clear that some fluid had collected in the newly-formed tunica vaginalis. A week later, at the second dressing, the wound had healed by first intention except in a minute area at the upper end. The hydrocele was the same as at the last dressing, and extending from its upper end to the external abdominal ring the spermatic cord was felt considerably thickened and indurated. Also the hydrocele on the right side was as before. In November these hydroceles remained, the left one being the size of a hen's egg, and tense and irreducible, but giving no trouble. There was no inguinal weakness manifest.

The existence of this type of bilocular hydrocele with an intrapelvic and a scrotal sac probably depends upon a faulty obliteration of the funiculo-vaginal process of the peritoneum. In the infantile form of hydrocele, not uncommonly seen in infants and young children, its obliteration has taken place near the external abdominal ring, and the whole of the rest of the process below that level, including the tunica vaginalis, has become distended with fluid. In bilocular hydrocele there is a sac of fluid in the scrotum and inguinal region like that in the infantile form, and in addition an intra-abdominal