OBSERVATIONS ON THE RATE OF VIBRATION
IN ANKLE CLONUS.

By J. A. MACWILLIAM, M.D.,

Professor of Physiology in the University of Aberdeen.

MANY observers have found the rate of vibration in ankle clonus to be usually about seven or eight per second. Gowers¹ says: "The movement is very uniform, from six to nine contractions per second." He recorded the movement by attaching a writing point to the foot, and causing it to inscribe its movement on a recording drum. Waller² described rates of 8, 9, and 10 in 3 cases of paraplegia, and a rate of 10 per second in a case of spinal syphiloma. Horsley³ found the rate to be 10 per second in a man suffering from caries of the spine. Herringham⁴ observed a frequency a about 7 per second in 4 cases from which he made tracings.

The upper limit of the vibration frequently seems, as far as has been described, to be sharply defined. I have not been able to find any record of a rate above 10 per second, but the following observations show that this limit is by no means an absolute one, that it can be overpassed to a very marked extent, and that different modes of registration may yield different results in some instances.

A man, aged 40, suffering from hemiplegia of ten months duration, showed greatly exaggerated "myotatic" reactions in the lower limb on the paralysed side. The knee-jerk was greatly increased; a single tap on the patellar tendon elicited several contractions in the quadriceps muscle—the condition called "tremospasm" by some observers. When continued tension was applied to the quadriceps by traction upon the patella a well-marked clonus was set up in the muscle and continued as long as the tension was maintained. A tap upon the tendo Achillis caused a multiple response from the calf muscles, two or three twitches following the tap. When ankle clonus was elicited in the usual way the vibration of the calf muscles looked unusually rapid, and tracings made showed that it was so. A tambour (button cardiograph, etc.) was strapped on the leg, so as to press on the calf muscles and record changes in them; this tambour was connected in the usual way with a recording tambour, while a time tracing showing half seconds was simultaneously inscribed. The rate of vibration was found to be as a rule 13.5 or 14 per second—that is about double the rate of ankle clonus as ordinarily observed. (See Fig. 1). In Fig. 2 a similar rate of

Fig. 1.

Fig. 2.

vibration (14 per second) is seen, while alternate vibrations exhibit a marked difference in character, each alternate curve being broader at the base and less sharp at the apex than the intervening curves. Indications of such difference, though only slight, may be detected in Fig. 1 also.

Fig. 3.

With somewhat different conditions in situation of tambour, amount of tension, etc., the differences in alternate vibrations may become very striking, so that the tracing obtained may look like a pulse tracing with a well-marked dicrotic wave (Fig. 3). Fig. 4 reproduces a portion of a tracing in which

Fig. 4.

the "dicrotic" type gradually merged into the kind of tracing shown in Figs. 1 and 2. At the right hand side of Fig. 4 there are alternate large and small curves; towards the left side the general characters of Fig. 2 develop; it is seen that the small alternate waves grade into the broader curves of Figs. 2 and 1. In the early part of the tracing the rate is about 12 per second; in the later part 13.

The effects of two single taps on the tendo Achillis are shown in Fig. 5; two well-marked twitches occur after each

Fig. 5.

tap; these twitches occupy somewhat longer time than two contractions of the ankle clonus. After the two twitches (in Fig. 5) there come one or two slighter oscillations of longer duration. (Upward movement of the lever = contraction).

Fig. 6 shows a tracing of the multiple knee-jerks caused by

Fig. 6.

two separate taps on the patellar tendon, recorded by a tambour strapped on over the quadriceps muscle. Three large contractions resulted in each case, coming at the rate of about 10 per second.

Fig. 7 shows the clonus excited in the quadriceps by

Fig. 7.

traction on the patella; it exhibits a rate of about 7.5 per second. It is noticeable that this clonus is decidedly slower than the rate of succession in the three well-marked contractions excited by a tap on the patellar tendon; this is in contrast to what has been stated above in regard to the calf muscles, where the rate of ankle clonus was somewhat higher than the rate of succession in the contractions caused by striking the tendo Achillis.

It is important to note that while a rate of 13 to 14 per second was obtained during ankle clonus when the receiving tambour was strapped on the calf muscles, a very different result was got when the movement of the foot was recorded, the button of the tambour being brought into contact with the ball of the foot while the clonus was going on. The vibration shown was only half as fast (see Fig. 8). In some

instances simple large curves at this rate occurred; in other instances the large curves described by the foot movement were notched at the top, indicating a probable fusion of two vibrations into one large curve. Thus, it is evident that the large movements executed by the foot are by no means necessarily a true indication of the actual vibration in the calf muscles.

Fig. 8.

The tendency to fusion which sometimes exists in cases of tremor was pointed out by Dawson Williams and Wolfenden,⁵ who examined cases of paralysis agitans, disseminated sclerosis, etc., by placing the hand (fingers or palm) on an india-rubber bag connected with a recording tambour. In a case of paralysis agitans they obtained large curves at the rate of 5.1 per second; these were notched at the summit, indicating a true rate of 10.2. So in disseminated sclerosis giving 5.2 large waves per second there was evidence of secondary waves, pointing to a true rate of 10.4. In some tracings they noted alternate smaller waves, recalling the appearances seen in a tracing of a "dicrotic" pulse. Herringham⁶ also noted the tendency to fusion in some tremors.

It is hardly necessary to remark that I do not affirm the occurrence of fusion of ankle clonus generally, but only that such may occur, as in the case described above, and that the foot movement may only show half the real rate of vibration in the muscle.

So high a rate of vibration in ankle clonus as 14 per second is interesting in regard to its relation to the character of the discharge, tonic influence, etc., from the motor cells of the spinal cord. Horsley and Schäfer⁷ state that the automatic action of the nerve cells of the cord appears to be never capable of originating a rhythm of greater frequency than 10 per second. But, while we know that the integrity of the spinal cord is essential for the occurrence of ankle clonus and cognate phenomena, we have no definite knowledge as to what relation the cord bears to the rate of vibration.

REFERENCES.

- ¹ *Diseases of the Nervous System*, London, 1886, vol. i, p. 13. ² *Brain*, iii, p. 179. ³ *BRITISH MEDICAL JOURNAL*, 1885, i, p. 112. ⁴ *Journal of Physiology*, 1890, xi, p. 482. ⁵ *BRITISH MEDICAL JOURNAL*, 1888, i, p. 1049. ⁶ *Loc. cit.* ⁷ *Ibid.*, vii, p. 106.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

SUPRARENAL EXTRACT IN GASTRO-INTESTINAL HÆMORRHAGE.

Apropos of Mr. Thelwall Thomas's interesting article upon the hæmostatic properties of suprarenal extract which appeared in the *BRITISH MEDICAL JOURNAL* of November 23rd, it may interest some of your readers to know that the drug is often of the greatest value in the treatment of severe hæmatemesis. During the past eighteen months the results I have obtained from its employment have been so uniformly satisfactory that I have quite abandoned all other remedies in its favour, and believe that in at least five cases death would certainly have ensued from loss of blood had preference been given to the more orthodox methods of treatment. The plan usually adopted was to administer 10 fluid ounces of a freshly-prepared decoction containing 2 grs. of the desiccated gland to the ounce, as soon as possible after an attack of

hæmatemesis, and to repeat the dose at the end of two hours. In two instances, where melæna was the sole symptom, three doses were given in four hours. In no case were any unpleasant symptoms observed, though occasionally the first dose produced an inclination to vomit. The cases which afford the most immediate and striking results are those in which the ulcer is comparatively recent, and is situated near the cardiac end of the stomach, as, for example, in the acute disease of young adults, and in that which sometimes complicates cirrhosis of the liver and valvular disease of the heart. In chronic ulcers of the pyloric region, on the other hand, where the eroded vessel is often of large size and so firmly embedded in fibrous tissue as to be incapable of contraction, experience will probably show that suprarenal tissue, in common with other hæmostatics, is of little value: but even in such cases an effort should be made to irrigate the stomach with a strong solution through a soft tube.

I might also mention that the constant slight hæmorrhage of gastric cancer often ceases entirely with one dose a day, while enemata of a similar character will often control the bleeding from an inoperable cancer of the rectum.

London, N.

W. SOLTAU FENWICK, M.D.

A BREEDING PLACE OF MOSQUITOS.

The accompanying photograph of a plantain illustrates what I believe to be, hitherto, an unobserved source of mosquitos. Each year the old leaves wither, and are cut down, leaving cup-shaped hollows, marked *c* in the photograph. These are

filled with a watery fluid, partly the exudation of the plantain and partly due to rain, and I have found myriads of mosquitos, together with eggs and larvæ, in these cavities. These plantains are cultivated to a large extent in Rhodesia for their ornamental value, but I feel sure not wisely.

Enkeldoorn, Rhodesia, South Africa.

A. DUNLEY-OWEN.