mentioned. In the case detailed below, gastro-enterostomy was preferred on account of the indurated condition of the pylorus. This was judged to be non-malignant at the time; but lest subsequently this estimate should prove to have been erroneous, the operation of gastro-enterostomy appeared the safer course to adopt.

G. W., aged 43, had been a dyspeptic for twenty years, but he did not begin to suffer acutely till twelve months before operation. Previously occasional bilious attacks with vomiting and diarrhoea had occurred. After an attack of sea-sickness, however, frequent gastric pain, relieved by vomiting, became a more constant feature during the next year. The site of the pain was some 3 or 4 inches from the pylorus, showing, as many other cases of gastric ulcer have done, that pain is a most fallacious guide to the situation of an ulcer. Doubtless this was the period of active ulceration, though no hæmatemesis occurred at any time to confirm this opinion. No palpable pyloric tumour could be made out. Increasing weakness and emaciation followed, and the signs of dilatation of the stomach developed rapidly. The organ when distended reached $2\frac{1}{2}$ inches below the umbilicus. Everything in the way of gastric medicines and digestive aids had been employed by Dr. Austin, Dr. Eustace Smith, and other advisers with little effect, including one-tenth grain doses of bichromate of potassium. Washing out the stomach with the siphon tube gave most relief, though this lavage exhausted him much. Nevertheless he steadily lost strength, flesh, and colour, and gave up his employment after losing 3 stones in weight. An operation was determined upon for his relief, upon the assumption that cicatrization of an ulcer was responsible for the dilatation.

On opening the abdomen, gastro-enterostomy was selected in preference to pyloroplasty on account of the hard cartilaginous feel of the pylorus, and Senn's decalcified bone plates were employed. As the visceral wounds were made very ample, the plate sutures were supplemented by a row of Lembert's sutures all round. The stomach was incised about $2\frac{1}{2}$ inches from the greater curvature and about 7 inches from the pylorus. The jejunum was opened about 1 foot from the duodeno-jejunal junction. The intestine was doubled back upon itself, as suggested by Rockwitz, before applying it to the stomach in order to secure correspondence in the direction of the peristaltic waves in the two organs.

The operation was aseptic rather than antiseptic; and the course afterwards was uneventful, neither pulse nor temperature being disturbed. Neither opium nor morphine was given. For three days absolutely nothing by the mouth was permitted. Four nutrient injections were administered each day of the formula recommended by Dr. Knut Hoegh.¹ These injections were well borne, and gave more satisfaction than other forms of nutrient enema that had been used in other cases. On the third day tea, wine and water, and meat juices were allowed by the mouth. By the fifth day soup, beef-tea, and white foods were given; and on the eighth fish and fowl were added. Finally, on the twelfth day ordinary solid food was ordered.

The patient rose on the fifteenth day, and on the seventeenth was exhibited to the members of the North of Ireland Branch of the British Medical Association. A fortnight later he reported himself as feeling well, having gained 2 pounds in weight, and proposed resuming his duties as an engineer to a railway.

PYLORECTOMY FOR SUPPOSED CANCER: PATIENT IN GOOD HEALTH THREE YEARS AFTER OPERATION.²

By T. R. JESSOP, F.R.C.S.,
Consulting Surgeon, Leeds General Infirmary.

SEEING that the patient whose history I am about to narrate is now in the enjoyment of good digestion and excellent general health at the expiration of nearly three years since

¹ *Annals of Surgery*, Part 18: "One ounce of glucose boiled with one drachm of flour and five ounces of water; after this has cooled there is added half an ounce of claret and two eggs beaten up with half an ounce of water."

² Read before the Yorkshire Branch of the British Medical Association at Scarborough, November 7th, 1894.

the removal of her pylorus for new growth, the time has arrived when the facts may be placed on record with the confidence that the charge of hasty publication will not be brought against me. I had once previously performed pylorotomy at the Leeds Infirmary, on a patient whose disease proved to have made more extensive inroads upon both stomach and duodenum than had been divined, or than appeared even on handling the parts when exposed for removal, until too late for the operation to be abandoned, and in whom, therefore, owing chiefly to the prolonged nature of the operation, the shock proved fatal within three days. Since meeting with these two I have not seen another example in which the circumstances seemed favourable for pylorotomy.

Mrs. R., aged 49, first came under observation on November 26th, 1890. She was then complaining of abdominal pain and vomiting, and had lost upwards of 2 stones in weight. It was noted at the time that the stomach stood out in the epigastrium, suggesting retention, and a distinct splashy sound could be elicited. About midway between the ensiform cartilage and the umbilicus, and about 2 inches to the right of the middle line, a hard, irregular nodule could be felt. A diagnosis of pyloric cancer was made. Up to January, 1891, under a rigid dietary and some medicinal treatment, the symptoms seem to have abated. The patient gained a few pounds in weight, and felt better in every way, and the lump in the abdomen could not be found.

In September sickness returned, and gradually increased in severity up to December, 1891. She then looked fairly healthy in face, but was wasted. The abdomen was decidedly thin; the stomach dilated and splashy. A swelling the size of a tangerine orange could be felt at the pyloric end of the stomach.

Satisfied that the disease was malignant, that she was surely losing ground, whilst her strength was by no means exhausted, and that the physical signs in her abdomen indicated an absence of adhesions, I advised her to consider the question of having the pylorus removed. The proposal was favourably entertained by herself and her husband, but before giving their consent they naturally desired to consult the physician upon whose judgment many members of their family had been accustomed to rely. She accordingly went to Glasgow, and there saw Dr. S. Gemmell, who, after a careful examination, wrote to me confirming the diagnosis, and approving the proposed operation.

From December 24th to 28th the stomach was washed out daily, and the patient fed by nutrient enemata and suppositories.

On December 28th, 1891, pylorotomy was performed. An incision $3\frac{1}{2}$ inches long was made in the middle line, starting at the apex of the ensiform cartilage. The stomach at once presented, and the lump at the pylorus was easily found. The stomach and duodenum were secured by means of Hahn's clamps and about $1\frac{1}{2}$ inch of bowel removed. The cut ends of stomach and duodenum were then united by fine silk sutures, the serous surfaces posteriorly being united first, then the mucous membrane all round, and finally the serous surfaces anteriorly. Lembert's sutures were used for the peritoneum. The incision through the stomach wall bled freely, but the vessels were readily seen and ligatured. Once during the operation the patient attempted to vomit, and forced a little fluid through the opening in the stomach. The peritoneum of the median incision was secured separately with fine catgut sutures, and the remaining layers of abdominal wall stitched with silkworm gut. No drain was employed.

The patient vomited once or twice during the afternoon after the operation. The vomit contained a little blood. She was fed by nutrient enemata every three hours. Morphine gr. $\frac{1}{4}$ was injected hypodermically in the evening, after which she slept five hours.

On December 29th, at 6 p.m., the patient commenced to take a drachm of milk with two of water every two hours, by the mouth, but as after the third time she vomited a little, it was discontinued. Morphine gr. $\frac{1}{4}$ was injected for the night.

On December 30th the note made was: Slept five hours; given ice during the night, vomited a little in the morning and two or three times this afternoon; given morphine gr. $\frac{1}{4}$ at night. She passed a good night without vomiting, and on

December 31st she was allowed to have a little barley water. She vomited, at 11 P.M. on that day, a little greenish fluid; and on January 1st was ordered to have milk and soda, barley water, and koumiss. The general condition good. The abdomen was a little distended; flatus was passing freely. She was given a turpentine enema on this night. No faeces passed, but some flatus and the remains of the nutrient enemata. After this the patient went on steadily towards recovery. There was no further vomiting. On January 9th she had a sponge bun. On January 19th some fish, and on January 21st some fowl. On January 25th she got up, and on February 9th went for a drive.

From time to time I have seen Mrs. R. since she passed from my direct observation. In the course of a few months she resumed her ordinary duties, and gradually she found herself able to enjoy unconscious digestion. At the present time she is in very good health, and shows no sign of having suffered from any serious disease.

EXAMINATION OF TUMOUR.

Length $1\frac{1}{2}$ inch; circumference 6 inches. The pyloric opening barely admitted the tip of the little finger; the stomach wall was much thickened, the duodenal wall less so. At the lower part of the pyloric opening was a saucer-shaped ulcer of the size of a shilling, with thickened base and edges.

Microscopically.—The chief bulk of the tumour consists of densely packed fibrous tissue, with intervening small round cells. Here and there in the interstices of the fibrous tissue are the remains of degenerated glands. In the deeper layers of the section, outside the fibrous tissue these glands are numerous.

FÆCAL FISTULA OF TWO YEARS AND TWO MONTHS' STANDING FOLLOWING HERNIOTOMY FOR STRANGULATED HERNIA: OPERATION: CURE.

By ANTHONY BOWLBY, F.R.C.S.,
Assistant Surgeon to St. Bartholomew's Hospital.

THE following case presents several interesting features. In the first place it is very rare to find the intestine give way as long as a month after herniotomy. I have myself seen it perforate as late as the ninth day, but not later. It is evident that the precaution of leaving the damaged gut near to the femoral ring allowed it to contract adhesions in a position which was a favourable one for the escape of its contents, and it is fairly certain that had it been placed in the midst of the peritoneal cavity acute peritonitis would have ensued.

E. D., aged 66, was admitted into St. Bartholomew's Hospital on March 3rd, 1893, suffering from a faecal fistula in the right groin of two years and two months' standing.

His history was that in January, 1890, he was admitted into St. Mary's Hospital suffering from a strangulated right femoral hernia, and that after operation he made a good recovery, and left the hospital with the wound closed at the end of three weeks. The registrar's note at St. Mary's Hospital recorded the following observation: "Gut deeply congested; sulci where nipped; returned into abdomen; sac removed but not ligatured; drainage tube passed into abdomen and removed next day."

A week after leaving hospital an abscess formed at the site of the operation and discharged spontaneously. The discharge of the pus was followed by escape of faecal matter, and this steadily increased in quantity. At the end of a year, however, the wound closed for three weeks, and then again reopened. For the next year he gradually passed more and more faecal matter by the wound, and for the last three months before admission had passed nothing by the rectum, except when an aperient was taken, the whole of the faeces passing for several weeks at a time by the fistulous opening. He also began to lose flesh quickly, and had gone down 2 stone in weight. It was under these circumstances that Dr. E. J. Lewis advised him to submit to an operation for relief.

Condition on Admission.—Fairly well nourished. In the right groin is a sinus in the situation of the crural canal. It

admits with difficulty the tip of the little finger, but soon narrows so as only to allow the passage of a director. The sinus constantly discharges liquid faeces and the skin around is excoriated and sore. On the left side is a scrotal hernia with an undescended testis. The viscera are generally healthy.

He was kept in bed and fed on light diet, and it was observed that his statement as to the absence of evacuation by the rectum was true, the bowels being only opened *per rectum* by aperients. A few hard, scybalous masses could sometimes be removed by enemata. Charcoal taken by the mouth was observed at the fistula forty-eight hours later. After consultation, it was decided to operate at the seat of the fistula.

On March 20th a vertical incision was made, opening up the fistula and dividing in part Poupart's ligament. It was found that a piece of small intestine occupied the crural ring and was drawn by the scar tissue a little way into the canal. It was dissected from the surrounding parts with some difficulty, owing to the very narrow space available, and was then discovered to have an opening in it half an inch in length. Into this aperture the opposing mucous membrane bulged. This opening was closed by a series of Lembert's sutures of fine silk, and the gut was placed in the abdomen just clear of the crural ring. It did not appear that the general peritoneal cavity was exposed during the operation, the neighbouring coils of intestine being adherent. The skin incision was not closed. Two days later faeces again passed by the wound, and continued to do so in considerable quantities for the next week. After that time the discharge rapidly lessened, so that by April 18th it was noticed that only about half a teaspoonful of discharge escaped in twenty-four hours. The bowels also acted regularly, with the assistance of a slight laxative, and sometimes for a day or two no flatus passed by the fistulous opening.

For the next month progress was not so satisfactory; the parts around the wound were a good deal inflamed, and he also suffered from an attack of inflammation of the right testis. The quantity of faecal discharge varied much, but on the whole it diminished.

By June 3rd the discharge was simply purulent, and contained no faeces. On this day he was put under an anaesthetic, and as the escape of pus did not seem quite free, the aperture in the skin was enlarged, and the mouth of the sinus erased. A little rough exposed bone was discovered near the pubic crest. From this time improvement was rapid. For a week at a time no faeces escaped, and after June 20th he never passed flatus by the wound.

On July 21st he was discharged, the sinus still being open, but no faecal matter escaping.

On October 30th the patient came to show himself. He said that the sinus had healed finally a week or two after leaving the hospital, and an examination of the parts showed a sound, firm cicatrix. The man had gained a stone and a half in weight, and his bowels were open naturally. He had returned to his work as a bootmaker. A year later—that is, in November, 1894, he remained in good health and with a sound scar.

The result of the small plastic operation performed was most satisfactory. At first I was disposed to treat the artificial anus by resection of intestine and suture; but, considering the very considerable risks of such a proceeding in an old man, I came to the conclusion that it was better to see what could be done by less severe measures. I have chiefly brought the case forward as one which encourages the performance of such a comparatively trivial operation, for there is a tendency at present to consider that long-standing faecal fistula after hernia requires resection of intestine for its cure; and seeing that in my patient the fistula was of more than two years' standing, it is evident that mere duration does not call for the performance of the more serious of the two operations which may be employed.

At a meeting of the Berlin Medical Society on January 9th Professor Virchow was re-elected President for the ensuing year.

JOHN MCPHAIL DOUGALL, M.D., M.B., C.M., was unanimously elected Chairman (from outside) of the Parish Council of the parish of Welburn, near York.