REMARKS ON TWO CASES OF RECTAL DISEASE TREATED BY MODIFIED OPERATIVE METHODS.¹

By G. H. MAKINS, F.R.C.S.,
Assistant Surgeon to St. Thomas's Hospital.

I.—RECTO-URETHRAL FISTULA.

W. F., aged 23, a collier, a strong healthy young man, with a good family and personal history, who when 7 years of age underwent the operation of left lateral lithotomy for vesical calculus. Since that date he had always passed a portion of his urine *per anum*; at times the whole had been passed in this manner, but more frequently the greater part *per urethram*. Gas was passed *per urethram*, and occasionally faecal particles, but as a rule he suffered no pain. Rectal control was complete, but he had not complete control over the urethra, and incontinence was noted almost daily. Examination showed the old perineal scar firmly healed; on the passage of a catheter about 1 inch could be felt bare in the membranous urethra when the finger was placed in the rectum. The recto urethral opening lay about $1\frac{1}{2}$ inch from the anus, admitted the tip of the forefinger readily, and had the usual sharp knife-edged margins. The urine was acid, specific gravity 1025, and contained a trace of albumen.

Operation.—On October 7th, 1893, the following operation was performed: The patient was placed in the lithotomy position, the perineum shaved, and an H-shaped incision made. The horizontal arm was 2 inches in length and about 1 inch anterior to the anal margin. Each vertical limb was about $1\frac{1}{2}$ inch in length. The skin having been divided, the wound was rapidly deepened with scissors, the left forefinger being kept in the rectum as a guide until the communication between the urethra and bowel was reached. The membranous urethra was found large enough to admit the forefinger readily, and was separated from the rectum only by a sharp marginal ridge.

A circular incision was now carried around the ridge and the urethral mucous membrane dissected up for half an inch on either side. Lembert's stitches were passed on the raw surface of the flaps so raised, and the mucous membrane was turned in towards a catheter placed in the urethral canal. The urethra was thus closed.

Attention was then turned to the rectum. The rectal wall, already free below, was somewhat separated from the back of the prostate above; the fistulous opening was continued down to the anus, dividing both sphincters. A set of Lembert's sutures were employed on the raw surface of the bowel flaps, and the canal was closed, as had already been done with the urethra.

A conical wound, extending to the prostate, with the closed canals in front and behind, now existed, the lateral vertical incisions allowing of considerable separation of them. Deep silkworm gut sutures were now carried through the raw sides of the wound, as is done in Mr. Lawson Tait's ruptured perineum operation, and the whole perineum closed, except a small cleft at the anterior extremity, which was plugged with gauze.

A No. 10 gum elastic catheter was left in the urethra to drain the bladder. There was remarkably little bleeding during the operation. Two and a-half hours afterwards, however, an untoward event occurred which threatened to wreck the whole operation. The patient asked for the bedpan, and evacuated about 16 to 20 ounces of blood, partly fluid, partly in clots. On examination blood flowed with some freedom from the anus, and Mr. Stabb, who saw the patient in my absence, plugged the rectum for me. I saw the patient about an hour later, and found him pale and with a feeble pulse. No blood passed *per urethram*, or by the catheter.

On October 8th the patient was improving; no further hæmorrhage had occurred. On the third day the rectal plug was removed; this operation, by distending the anus, caused some disturbance of the wound; the catheter also got blocked, and urine began to escape by the drain opening left at the front of the wound. The catheter was therefore removed, but it was not thought advisable to pass a fresh one

for fear of disturbing the urethral suture. For some days the urine passed almost entirely by the perineum, and the deep stitches were therefore removed to prevent the risk of extravasation taking place in the deep part of the wound. The margins did not gape, and although a few drachms were passed *per anum* on one or two occasions, no appreciable quantity passed into the rectum. On the eleventh day he began to pass his water *per urethram*, and the perineal wound rapidly healed, so that on the nineteenth day he was practically well. He remained in the hospital another twelve days, and has since resumed his occupation as a collier.

I received a note regarding him from his doctor some weeks ago, in which he tells me the man remains well, and passes his water naturally, the only trouble consisting in the fact that at times he needs to pass his urine at once when the bladder is full. This slight defect has apparently existed ever since the lithotomy operation, and is, I think, probably due to the scar in the prostate.

As will at once be recognised, the operation performed in this case is a modification of Mr. Lawson Tait's operation of perineorrhaphy, the H incision allowing of lateral approximation of sufficient perineal tissue between the rectum and urethra to get an efficient barrier between the two canals. Although some complications arose, and the primary union I hoped for was not obtained, yet the result was, in my opinion, a gratifying one, as I think anyone will admit who has had experience of dealing with these very troublesome cases.

The advantages which I would claim for the method are the following:

1. The production of a sufficient mass of perineal tissue between the rectum and urethra, gained by the lateral vertical incisions.
2. The opportunity given of widely separating the edges of the cut mucous surfaces meeting at the fistula.
3. The possibility of remoulding the two canals.

As regards my own experience of treating this condition, I may say that I have twice had the opportunity of operating by other methods on another patient. On the first occasion, I performed a suprapubic cystotomy to remove a recurrent phosphatic calculus. I kept the wound open as a drain, and afterwards performed a plastic operation by paring the edges of the fistula and suturing through the dilated anus. The operation much reduced the size of the fistula, but it did not entirely close. A second attempt at suture by the same method was also unsuccessful. I attributed this to the extreme tenuity of the edge of the fistula, and the impossibility of bringing a sufficiently deep raw surface into contact, a condition which is satisfactorily met by the method I now advocate.

It may be remarked that the new method, when the case does not heal by first intention, is not widely different to the old plan of slitting up the fistula, and laying the rectum and wound into one, and then plugging while the wound heals by granulation. I would submit, however, that there is an essential difference, namely, that by the new method the separated mucous-lined canals are much more widely separated, that the lining membrane is inverted, and that the possibility of the two mucous surfaces again becoming continuous is reduced to a minimum. The latter result also is much the less likely from the much more rapid healing which can take place. Again the prompt suture of the divided sphincter and is a manifest advantage and comfort to the patient.

II.—GROWTH OF RECTUM.

The second case which I have to bring to your notice is also one depending on the modification of some of the recognised operations for removal of growths of the rectum. Only a certain number of cases suitable to the method are likely to be met with, but for such as may occur I think the operation has much to recommend it, as I hope the recital of the following case will show.

M. A. C., aged 49, a housekeeper. Her father died at the age of 72; her mother was alive, aged 76. One sister died of consumption, and a brother from some complaint of the bowels at the age of 50. The patient had never been very strong, but had only been under regular medical treatment on two occasions, once for varicose veins, and once for hæmorrhoids. She had not suffered from habitual constipation. Her present illness commenced eight months before

¹ Read at a meeting of the South Eastern Branch of the British Medical Association held at St. Thomas's Hospital.

she came under treatment with relaxation of the bowels, and since then she had habitually passed watery motions. She has occasionally passed blood; this she has attributed to the hæmorrhoids. She has suffered from prolapse on and off for years. She has been troubled with sickness after food more frequently of late, but she seems to have been a chronic dyspeptic. These symptoms have been attended by progressive weakness, inability to do her work, and slight loss of flesh.

The patient was a thin, sallow woman, and complained of extreme weakness, frequent sickness, and constant diarrhoea. On examination *per rectum*, under an anæsthetic, well-marked mixed piles were found, and on the anterior wall, about 2 inches from the anus, a definite sessile papillary growth with broad base, which could be drawn down to the anus with a clamp. There were no signs of other visceral disease.

Operation.—On August 29th, 1893, the patient was anæsthetised, and the following operation was performed. The anus was forcibly dilated, a circular incision made at the junction of the skin and mucous membrane, and the mucous membrane and piles were freed from the external and internal sphincter muscles by dissection with scissors. Thus far the operation was the typical one performed by Mr. Whitehead for hæmorrhoids. The sphincters were now divided towards the vaginal wall to obtain increased space, the muscular and areolar coats of the rectum divided by a circular incision, and the dissection was carried upwards around the whole rectum for another $2\frac{1}{2}$ inches until a point above the limits of attachment of the growth was reached. The peritoneal cavity was not opened. The freed portion was then divided, and the bowel brought down and sutured to the cut margin of the anal skin with silk. The incision into the perineum was closed with two silkworm gut sutures, which also retained the anterior ends of the sphincters in apposition. A large tube was placed in the rectum to prevent the accumulation of flatus, and the wound was dressed with bichloride gauze.

The after-course was uneventful. Retention of urine necessitated the use of the catheter for a few days. The wound healed steadily. On the tenth day the deep perineal sutures were removed. After the eighteenth day complete control of the motions returned, and she got up. She steadily improved, and left the hospital well one month after the operation.

After a short holiday she resumed her work, which she has continued to do, and she and her medical attendant consider that she is stronger and better than she had been for years.

The growth removed was papillary in character with a broad base, occupying an area of 2 by $2\frac{1}{2}$ inches on the anterior wall of the rectum. Careful examination showed that it had not infiltrated the whole thickness of the rectal wall; in fact the muscle was scarcely affected, hence the prognosis is a comparatively good one.

As to the nature of the growth, microscopical examination shows it to be identical in structure with the ordinary polypoid growths or adenomata. Experience has shown, however, that such growths are common precursors of typical columnar epitheliomata, and therefore a guarded prognosis only can be made. In this particular instance the free removal of the tumour should, however, give the patient a good chance, whatever the actual nature of the growth may be.

The mode of operation adopted, however, is the point of interest in this case, and I think it offers the following advantages:

1. It leaves the sphincters as nearly as possible intact. If the growth were lateral or posterior, the posterior extremity of the sphincter might be divided towards the coccyx.

2. The union of the rectum to the skin is preferable to uniting it to a strip of mucous membrane left covering the sphincter flaps made when the anus is divided both anteriorly and posteriorly; and any existing piles are radically cured by the same operation.

As to the indications for such a procedure I take it that they consist in limited extent of the growth, and in the pre-existence of rectal prolapse after defæcation, this the cardinal indication for the complete Whitehead's operation for

hæmorrhoids, allowing the portion of the bowel brought down to the anus to be sutured there without undesirable tension.

CASE OF PERFORATION OF THE INTESTINE DURING ENTERIC FEVER TREATED BY LAPAROTOMY.

By ALFRED PARKIN, M.S., M.D. LOND., F.R.C.S.,
Senior Surgeon to the Victoria Hospital, Hull.

This case is of interest on account of the discussion on the operative treatment of perforations of the stomach and intestines at the Bristol meeting of the British Medical Association.

A female, aged 32, came under the care of my partner, Dr. Sherburn, on August 19th, 1894. She had been ill about a week with indefinite symptoms, but a few days previous to being seen diarrhoea came on, and continued throughout the illness. She was wildly delirious on the 20th and 21st, and on the 26th distinct rose spots appeared on the abdomen; the temperature remained at the level of 103° , the pulse 130, and the respiration varied from 26 to 30.

The condition of the patient remained the same until September 3rd, which would correspond to the end of the third week, when she was suddenly seized at 10 A.M. with acute abdominal pain, continuous, and referred to the umbilicus. At 10.15 A.M. she was sick, and when seen half an hour later she was in a collapsed condition, with face drawn, lips blue, and extremities cold. The abdomen was no longer doughy but hard, less distended, and tender to touch. The bowels were opened twice involuntarily, the motions being formed, and it seemed as though the contraction of the abdominal muscles had been the cause of this. The liver dullness, which before had extended from the sixth rib to the margin of the thorax in the upper right, had now disappeared. Pulse imperceptible, respirations quick and shallow, temperature 101° . An injection of morphine and atropine was given at once, and every effort made to overcome the collapse. These efforts were so successful, that on returning at noon the patient seemed as well as before the perforation, only the absence of liver dullness and slight abdominal rigidity indicating what had happened.

Chloroform was given followed by ether, and a small incision made below the umbilicus. On opening the peritoneal cavity a considerable quantity of gas escaped, without much smell, followed by a few ounces of turbid yellow fluid. The small intestine was traced throughout its whole length, the position of many ulcers being indicated by the brilliant injection and the surrounding thickening of the intestinal walls. About 2 feet from the cæcum there was a large ulcer, the floor of which displayed a small perforation not bigger than a pin's head, through which the intestinal contents were oozing. The ulcer was folded longitudinally on itself and sewn up with a Lembert's continuous silk suture, without difficulty on account of the congested state of the gut, which would scarcely hold the silk. No other perforation being found, the peritoneal cavity was well irrigated with weak carbolic, a Keith's tube inserted into the pelvis, and the wound sewn up.

On recovering from the anæsthetic the patient said she had no pain, and was quite comfortable; in every way her condition was better than before the operation. She was sick twice during the night, and the next morning her pulse had fallen to 120, the temperature was normal, the abdomen flaccid, not distended, and moving readily with respiration. She was fed with nutrient enemata every four hours, and with peptonised milk, beef-tea, and brandy by the mouth. Her condition remained quite satisfactory, with no symptoms of peritonitis until the third night after the operation, when she again became delirious, the temperature rising to 102° . At 1 A.M. on the 6th she tried to get out of bed, and then complained of the abdominal pain having suddenly returned. The abdomen became rapidly distended, collapse supervened, and death took place ten hours after the return of the pain, exactly three days after the operation. Before death the temperature rose to 106° .

A very limited *post-mortem* examination was allowed. There was no peritonitis around the sutured gut, but close to the cæcum there was a large ulcer with an extremely thin floor devoid of peritoneum, and around this there was evident peritonitis. The portion of the gut operated upon was quite tight, and withstood considerable pressure from injected fluid. The ileum for 3 inches close to the cæcum was ulcerated all round, so that only islets of mucous membrane were left. It was not possible to remove the whole of the intestine, large and small, to make certain that no second perforation had taken place, but none could be found by pulling out various portions of the intestine through the wound.

This case adds one more to the list of 20 such cases which have been operated upon, with so far only 1 recovery. There is no doubt that perforation in the course of enteric fever is almost invariably fatal, and any recovery from such a complication can only be regarded in the light of an accident, unless surgical treatment be adopted. The obscurity in the diagnosis is often great, but in this case the symptoms were pronounced.

Although the patient was in the height of a severe attack perforation was evidenced by sudden collapse, alteration in the character of the abdomen, and by complete disappearance of the liver dullness. Curiously enough, the perforation causing these symptoms could only be regarded as of the smallest possible size. The treatment adopted in the first