practitioner has to do is to withdraw the plug with the attached brush, to which it forms an excellent handle, while the mouth of the tube is held downwards, to scrape off some of the discharge from the affected parts with the brush, to replace the cotton plug and attached brush in the tube, and send off the box with enclosed tube by post.

In starting a culture on blood serum or other basis the brush which contains the secretion is rubbed over the surface of the medium.

The advantages which the method seems to possess are: (1) That the discharge is removed by and received in sterilised instruments; (2) that the brush is an excellent means of collecting the discharge, and does not alarm the patient by its appearance; (3) that everything being ready for sending off by post no time is lost in transmission; (4) that the brush is again an excellent instrument for spreading the discharge over the surface of the culture basis; and (5) that the cost of the apparatus is trifling. These sterilised tubes could be kept by local apothecaries, and be had on a moment's notice. They are of small bulk, so that they can readily be carried about when required.

REMARKS ON MILK DIET IN THE SEVERE INTESTINAL DISEASES ACQUIRED IN THE EAST.

By GEORGE THIN, M.D.

DURING a long experience of the management of certain climatic disorders of the intestine by exclusive milk diet, my attention has been much directed towards minimising the difficulties inseparable from the treatment. Many persons have a strong dislike to milk, and I am often met by the statement that the patient "has never been able to take milk in his life." One sufferer from a severe form of psilosis prefaced her statement of her case by telling me: "It is no use prescribing milk for me. I can neither take it nor digest it. All the medical men whom I have consulted have soon satisfied themselves that this is correct, and have been obliged to give me something else." For cases like that one, I have found the use of agitated milk¹ of inestimable value. So far, I have only twice failed to succeed in getting a patient to go on exclusive milk diet. In one case, which was neither severe nor well marked, the diarrhoea did not rapidly yield, and the treatment was abandoned after a short trial; and another patient, who was doing extremely well, gave it up for, I am afraid, no better reason than caprice, with disastrous results.

Another and sometimes a serious difficulty in connection with milk diet is the quantity of fluid which the patient must swallow. Many patients, especially when convalescing, take as much as 6 to 8 pints daily and even more without any difficulty, but there are others in whom the necessary quantity of milk suited to the case produces a constant feeling of distension accompanied by troublesome flatulence. There are also cases in which the patient is very ill, and his life in great danger, in which it is impossible to get him to swallow sufficient milk to sustain life. During the past summer I had a case of this kind under my care. A man who had been for two years suffering from chronic diarrhoea acquired in Bombay, complicated with fistula and an unhealthy condition of the mucous membrane of the rectum, on which Mr. Herbert Allingham found it necessary to operate, fell into a low typhoid condition. The difficulty of getting him to swallow sufficient milk was so great, and his repugnance to it apparently so invincible, that the treatment was abandoned for three days, during which time an endeavour was made to support him on beef-juice. His condition became rapidly worse; there was incessant diarrhoea, the motions being passed involuntarily; the tongue was hard, dry, and cracked, the teeth covered with sordes, and the pulse quick and thready. We considered the chances of recovery to be very small; he seemed, indeed, to be rapidly sinking, but I was satisfied that if he were to be saved at all it could only be by

milk, and accordingly exclusive milk treatment was resumed. He could swallow only such small quantities at a time that it was found impossible to get him to take more than 2½ pints in twenty-four hours. The typhoid condition and incessant diarrhoea continued without improvement for forty-eight hours, and it became evident that unless more milk could be assimilated it was hopeless to look for recovery.

I then directed part of the milk to be evaporated to one half its bulk, and by giving him this evaporated milk he was able to take at once the equivalent of 3½ pints in twenty-four hours, which proved sufficient to arrest the threatened collapse. Gradually, after a time, his condition began to improve, and he eventually consumed in natural and evaporated milk the equivalent of 9 pints daily. He made a perfect recovery, and in five months was shooting grouse in Scotland.

I afterwards prescribed evaporated milk with excellent results in another serious case of wasting from long-standing chronic diarrhoea acquired in India. The patient, who had found it extremely difficult to swallow more than 3 pints of fluid a day, was able, by taking a certain proportion of his milk evaporated, to consume the equivalent of 5 pints, gradually lost his diarrhoea, and was able to digest solid food and gained strength proportionately. Other patients, not in the same immediate danger, have been allowed the option of taking part of their milk evaporated, and some of them have thriven on the concession. Others have had no difficulty in taking, and have preferred to take, the milk in its natural state.

The two severe cases alluded to above were examples of the chronic intestinal catarrh (not dysentery) which is common in India, and which is often described under the name of "tropical diarrhoea." At the present time I have under my care a typical case of psilosis (*vulgo* "sprue") of long standing, which is steadily progressing towards recovery by the help of evaporated milk. By using part of the milk prepared in this way the patient is now able to take sufficient nourishment to enable the diseased bowel to recover. He has been under medical treatment more or less continuously for a number of years, but I gather from what he tells me that his occasional attempts to get well by milk diet were abandoned because he could not take sufficient milk to maintain the required minimum of strength. This induced him to supplement his diet by other articles of "invalid" food, and these invariably after a time brought on a relapse.

To evaporate milk properly for patients like these is a matter of some care. A simple apparatus has been made for me by James Allen and Son, 21, Marylebone Lane, W., consisting of a small regulating spirit lamp, with a wire stand for holding an enamelled pan, containing the milk, at a convenient height from the flame. The heat may be increased or diminished at will by the regulator on the lamp. The milk should be heated rapidly, but must not be allowed to reach the boiling point; and in this pan 10 ounces can be reduced to 5 in half an hour. It must be stirred continuously after it gets warm and until it has become cold to prevent a skin forming. If the cream is allowed to come to the top the milk takes longer to evaporate, and the taste is not so good. When the evaporation is properly effected the cream remains mixed in the milk as usual, but, of course, rises on standing. The milk should therefore be stirred before being drunk. Evaporated in this way it has a richer taste and looks richer than ordinary milk, but is not unpalatable. It gets sour more quickly than fresh milk. The process consumes a good deal of time and requires constant attention.

Whilst writing about the treatment by milk, I may make a few remarks in connection with one of the diseases for which it is the remedy. There is a note in the *Times of India* of September 12th which states "that Surgeon-Captain Dyson has traced that the disease known at Darjeeling as 'hill diarrhoea' and in the Straits and China by the peculiar name of 'sprue,' is due to the presence in the water of small particles of mica." Psilosis, or "sprue," has been often written about as identical with "hill diarrhoea" of India, and I am not sure that, under the influence of Indian medical literature, I have not myself done so at one time.

The affections known by these names have a feature in common, namely, the passing of frequent motions of a light

¹ See papers by Surgeon-Major Goldsmith, *BRITISH MEDICAL JOURNAL*, vol. ii, 1893, p. 12, and the writer, *BRITISH MEDICAL JOURNAL*, vol. ii, 1893, p. 64.

yellow colour, which, after all, is the usual colour of motions which are rapidly driven along the small and large intestine. I had, however, at one time a patient suffering from typical psilosis, whom I sent to spend a month with her brother, an army surgeon of long experience in India. When he brought her back to me he said: "The disease from which my sister is suffering is new to me. I have not observed it in the parts of India in which I have served, and it is not 'hill diarrhoea,' of which I have seen hundreds of cases." Typical psilosis occurs in Ceylon, and I have seen well marked examples in patients who have returned from that island. I have not met with evidence to show that it occurs at all commonly in Bombay or Calcutta. Cases of severe wasting diarrhoea from these parts of India about which I have been asked to give an opinion have so far been none of them cases of psilosis. Typical examples of the two distinct affections—the common chronic diarrhoea of India, and psilosis of the tongue and small intestine—are related in a paper presented by me to the Royal Medical and Chirurgical Society, and published in the seventy-fifth volume of the *Transactions* (1892).

As regards the presence of mica in the drinking water as a cause of this disease, I should regard the theory as improbable. I am not familiar with the distribution of mica in the world, but to prove it to be the cause of psilosis it would have to be found not only in Ceylon, but in the drinking water in the Straits Settlements, Hong Kong, the whole China coast, and at least several widely distant parts in the interior of China, in Java, and the Philippine Islands, in all of which places psilosis is very common. In the *Encyclopædia Britannica* I find it stated that mica is found in Finland, New York, Canada, Shetland, Sutherland, Ross-shire, Inverness, Skye, Fifeshire, New York, and Greenland, from all of which parts no case of psilosis has yet been reported. The only place where psilosis is common, and in which the presence of mica is noted, is Ceylon. Besides there are facts connected with the clinical features of this disease which cannot be reconciled with the mica theory. Patients who have been for a long time quite well are subject to relapses from chill and error of diet, although they have returned to England, where no case of this disease has ever been known to arise. Nay more, the disease may show itself in an unmistakable form for the first time after a patient has returned to England. An extraordinary instance of this occurred in the case of a patient who consulted me last summer. A lady, who had returned from China seventeen years before, developed unmistakable symptoms of psilosis, which had existed for nearly two years to a slight extent before its nature was detected. The development of all the characteristic symptoms to a degree of considerable severity led to the disease being diagnosed, and under the treatment appropriate to the cure of psilosis she has made a good recovery. Although she had not enjoyed robust health after her return from China, suffering particularly from anemia, she had never had symptoms of "sprue" until two years ago.

The proximate cause of psilosis is not known, but the probability that it consists in some organic ferment, very possibly a bacterium, is strong, and I think it likely that the curative action of milk diet is not due to any special therapeutic virtue of milk, but to the fact that it does not contain the pabulum in which this ferment can live. It is impossible otherwise to account for the fact that not only do the soothing foods which are so useful in other forms of diarrhoea not cure this form, but they will often bring on at once a severe relapse. I have often known arrowroot bring back diarrhoea and sore mouth after the patient had not had either symptom for several weeks. The same thing applies, in my experience, to beef-tea and bread, and severe and persistent relapses were produced in two patients by a raw, new-laid egg. It seems to me impossible to account for these facts otherwise than by assuming that in the egg, beef-tea, arrowroot, and I may add wheaten bread, a ferment can multiply with a rapidity that cannot be attributed to any inorganic substance.

UNIVERSITY OF JURJEFF.—The report of the University of Jurjeff (still better known probably by its old name of Dorpat) for 1894 show that on December 1st the total number of students was 1,546, of whom 754 belonged to the medical faculty.

CANNABIS SATIVA SEU INDICA : INDIAN HEMP.

By R. COWAN LEES, M.B., C.M., F.F.P.S.G.,
Glasgow.

It has always been difficult to understand why the resin of this plant should alone be recognised in the *British Pharmacopæia*, more especially when we find it stated in works on the physiological action of this drug that in India several preparations are used by the natives to produce its stimulating and exhilarating effects, amongst which watery infusions are specially mentioned.

During a short visit to India some years ago my attention was drawn to the fact that several modes of using the herb were employed by the natives—modes not capable of extracting much, if even any, of the resin. So far as I could observe, watery infusions were commonly used, but whether in combination with other substances or not I am not in a position to state. This fact, however, led me to try what benefits might be derived from the use of a preparation of the plant, not depending on the resin alone.

Messrs. T. and H. Smith, who first obtained the resin in a state of comparative purity, state that "it is a brown amorphous solid, burning with a bright white flame, and leaving no ash; powerful in its action when taken internally, and that two-thirds of a grain act as an active narcotic, whilst 1 grain produces complete intoxication"; but the question might be advisedly asked, Is it completely freed from its essential oil? As a matter of fact, it is found that when the extract is kept for some time it becomes hard and brittle, and less potent in its action, a circumstance which goes a long way to prove that such a condition is the result of loss of volatile oil from the resin, and pharmacists are advised to "lay aside and not employ for medicinal use that which has become old."¹

We are told by Bently and Trimen that "both Hindus and Mohammedans use this herb, either by smoking—with or without tobacco in combination with other substances—or by simple infusion in water."² Gunjah—guaza of our London market—has but a faint taste, with a peculiar but not unpleasant narcotic odour. These properties depend in a great measure on the volatile oil and resin. The latter some consider the more important constituent of the plant.

In the watery infusion employed by the Hindus and Mohammedans as mentioned above, we can conceive of little, if any, of the resin being dissolved and held in solution, whilst, on the other hand, much of the volatile oil might be dissolved by the water, together with other constituents of the plant. Dr. Personne regards the volatile oil as the sole active principle, and in proof of this he states that "when the volatile oil is inhaled, a distinct sensation of shuddering with motor excitement, followed by prostration and syncope, is experienced." Again, Dr. Preobraschensky has found a volatile alkaloid—most plentiful in the flowering tops—and which he considers somewhat similar in its action to nicotia or nicotine.

Feeling somewhat satisfied that water was capable of dissolving at least a portion of this volatile oil, and knowing that watery infusions of the drug were used for intoxicating and stimulating purposes in India, I had prepared for me a strong aqueous extract of the flowering tops of the female plant of the usual strength of liquid extracts, and from its use I have obtained good and satisfactory results. It possesses the anodyne and soporific action generally ascribed to the resinous extract, although in a modified degree. It has the characteristic odour of the hemp, has a beautiful deep amber colour, is miscible with water, and hence there is no difficulty in combining it with other liquids, and it presents no unseemly immiscible mixture repellent to a patient.

Liquor cannabis indicæ in my experience gives all the beneficial effects without the drawbacks of the tincture, avoiding those extreme exhilarating conditions bordering on intoxication, which are sometimes met with even when using a medium dose of the latter. It does not seem to in-

¹ *Squire's Companion to the British Pharmacopæia*.

² *Medicinal Plants*, vol. iv, page 231.