carbolic acid for this purpose in all cases of infantile diarrhoea, but its hot taste and unpleasant smell, as well as the occasional occurrence of carboloria, has made me transfer my allegiance to resorcin. This drug has the advantage of being extremely palatable to children, devoid of toxic properties when given in ordinary doses, and very inexpensive. According to the *Pharmacopœia*, the dose of resorcin is 1 to 5 gr., and it is probably on account of this insufficient dosage that the value of the remedy has been so much overlooked. As a matter of fact, the drug produces no ill effects in an adult unless the dose exceeds a drachm, and I have long been accustomed to prescribe 3 gr. every four hours to infants only a few weeks old without the least ill effect.

In cases of diarrhoea the first effect of the drug is usually noticeable after the third or fourth dose, when the motions decrease in frequency and in amount, the dejecta at the same time acquiring a more natural appearance and losing their excessive fetor. At the end of the second day the diarrhoea has generally ceased, and is not infrequently replaced by obstinate constipation. I have before me the notes of 120 cases of intestinal dyspepsia in infants and young children which have been treated in the manner described. Of these, in 53 per cent. the disorder had lasted from one to two weeks, in 34 per cent. from two to four weeks, in 10 per cent. from four to eight weeks, and in the remaining 3 per cent. for a period of more than two months. Out of the entire number in only nine instances did the diarrhoea continue after the treatment had been pursued for a week, the majority ceasing within three days. Of these nine refractory cases two were instances of cholera infantum, both of which succumbed within a short time of being brought to the hospital; in one case tuberculous ulceration of the intestine was found to exist after death, while the remaining six, most of which were very chronic in their nature, were rapidly cured as soon as benzonaphthol or salicylate of bismuth were added to the original mixture. These results go to prove that when given in sufficient quantity resorcin constitutes a reliable remedy in cases of infantile diarrhoea arising from fermentative processes in the stomach and intestine. When, however, the disorder has lasted for a long time and follicular ulceration of the large intestine exists, the drug may be advantageously supplemented by those remedies which exert their antiseptic properties in the intestine. Benzol naphthol is particularly valuable in this connection since it is but slightly toxic, but in order to prove effective it must be given in full doses at short intervals. Thus in some cases where the symptoms were obstinate I have prescribed as much as 40 gr. to a child during the course of twenty-four hours with only the most excellent results. My experience with the most recent antiseptic, the salicylate of strontium, has not been altogether satisfactory, and I have been unable to convince myself that it possesses any superior advantages over the corresponding salt of bismuth.

THE ATROPINE GROUP:

AN EXPERIMENTAL AND CRITICAL NOTE.¹

By GORDON SHARP, M.B. EDIN.,

Late Research Student in Pharmacology, the Owens College, Victoria University.

SOME twelve months ago, I had under my care for a considerable period, a lady of 25, who suffered from repeated attacks of renal colic in whom a $\frac{1}{2}$ gr. of atropine sulphate (along with $\frac{1}{4}$ gr. morphine hydrochlorate) although always relieving the distressing conditions, produced unpleasant flushing of the face, dryness of the throat, and confusion of the senses. Having read of scopolamine being free from the disagreeable properties of atropine just named, I thought I might be able to get all the advantages of atropine with none of its disadvantages. I resolved to try the drug for myself, but after trial I found it differed little or not at all from atropine. Previous acquaintance with hyoscine taught me that it was little different from atropine, and after some experiments with daturine and duboisine I was led to extend my studies of the whole atropine group, and to read up the literature for the past twenty years, hence this paper.

¹ Read in the Section of Pharmacology and Therapeutics at the Annual Meeting of the British Medical Association, held in London, July-August, 1895.

To the student of pharmacology no group of bodies is more confusing than the so-called mydriatic alkaloids known by the names of hyoscyamine, hyoscine, daturine, duboisine, scopolamine, and atropine, etc., all of which possess the property of dilating the pupil. A vague examination of them might lead one to the belief that each had some peculiar action of its own, but closer examination and increased knowledge show them to be very closely related if not identical one with another. True their sources are varied, but reasoning by analogy we know that tea, coffee, guarana, Paraguay tea and kola-nut are all very widely different plants, and yet they all contain one common alkaloid, theine, and not each a different base as at one time supposed.

The literature of the group of the atropine bodies is most interesting, because it shows the conflicting views of authorities; but as one comes to the present period it further shows a tendency to the view that we have only one or two bases to deal with. One other point it teaches us is that we cannot clear up the difficulty from the chemical side alone, but must call in the aid of pharmacology and therapeutics. With regard to Gerrard's test for the mydriatic group, I find after repeated trials that we do not, as is often supposed, get a yellow colour with one and a red colour with another member of the group, but with a trace of any one of them an alcoholic solution of mercuric chloride heated gives a yellow, while an appreciable quantity of any one of them gives a bright red colour of red oxide of mercury. The test is difficult to apply, and failures to obtain colour reactions with certain members may be accounted for in this way.

HYOSCYAMINE.

Whether hyoscyamine be identical with atropine or no one thing is certain—that they are very closely related; for in 1880 A. Ladenburg converted hyoscyamine into atropine, and in the following year the same authority found that both bases split up into tropic acid and tropine. Further comparisons show both to give a bitter almond odour with potassium bichromate and monohydrated sulphuric acid, and both, when heated with nitric acid and afterwards treated with alcoholic solution of caustic potash, become of a violet colour. Chemically so far, then, they would appear to be identical, and pharmacologically I do not know if we can distinguish between them.

DUBOISINE.

The introduction of duboisine about the year 1878 by Bancroft, Holmes, Gerrard, Ringer, and Tweedy stimulated research, and soon it was found that it was not a new base, but only an impure atropine. At this time A. Ladenburg gave it as his opinion that we had only three mydriatic alkaloids—namely, atropine, hyoscyamine, and hyoscine agreeing in formula and in decomposition products. Later Ladenburg so far modified his views as to believe only two bases exist—namely, atropine and hyoscyamine, and these are isomeric and very closely related.

DATURINE.

The introduction of daturine led to disputes amongst authorities. Some believed it to be identical with hyoscyamine, while A. Ladenburg considered daturine, duboisine, and hyoscyamine as identical. In the same year (1880) E. Schmidt said he failed to detect the difference between daturine and atropine. Deduction from this makes atropine and hyoscyamine identical. Passing to the year 1885 we find E. Schmidt saying atropine and daturine are identical, and in this view he is joined by Ladenburg.

HYOSCINE.

Coming to hyoscine we have seen that A. Ladenburg had so modified his views as to regard this body as not a different base from atropine or hyoscyamine. Pavloff in 1889, in summing up the pharmacology of hyoscine, regarded it as differing from atropine only in that it (hyoscine) diminished the irritability of the brain. Bender in 1892 looked upon hyoscine and scopolamine as identical, while in 1893 A. Ladenburg disputed the identity of the two. In the same year E. Schmidt said the so-called Ladenburg's hyoscine was really scopolamine, and in this view he was supported by O. Hesse.

SCOPOLAMINE.

In 1889 the alkaloids from both the Austrian plant and from

Japanese belladonna were looked upon as either a mixture of hyoscyamine with other alkaloids or as a mixture of hyoscyamine, atropine, and hyoscyne; but in 1892 E. Schmidt named the supposed pure base scopolamine, and said it occurred in belladonna root, stramonium seeds, and occasionally in *duboisia myoporoides*. O. Hesse looked upon scopolamine as identical with hyoscyne. Last year, however, O. Hesse, as the result of experience, came to look upon scopolamine as made up of two bases: one identical with hyoscyne, and the other isomeric with it, and he proposes to name it atrosyne. Kaehlman quotes Kobert as saying scopolamine has a paralyzing action on the brain, and does not accelerate the pulse, differing in this respect from atropine. He believes as a mydriatic, analgesic, and antiphlogistic it is superior to atropine, and is free from the objection of causing dryness of the throat, congestion of the head, and acceleration of the heart's action. Bokenham in 1894, as the result of his experience, said in some animals the pulse was slowed where it would have been rendered rapid by atropine, and in himself it caused no dryness of the throat. As an offset to these opinions we have the experience of Thos. K. Pooley, who found that six instillations of the conjunctival sacs of a one-fifth per cent. solution of hydrobromate of scopolamine produced in a girl of 13 irregularity in the heart's action, with a pulse of 120, staggering gait, complaint of dryness of the throat, and of "pins and needles" sensation in the soles of the feet, and there was continual working of the muscles of the face and lips. Two adult women in whom the drug was employed were similarly affected.

The conflicting views held as to the identity or non-identity of the various named bases makes it certain that our knowledge of them is yet far from exact. When we come to search for a cause of all this confusion, we have to bear in mind the following points: (1) the difficulty of obtaining a pure alkaloid; (2) the great readiness with which the members of the atropine group decompose, heat above a certain degree, acids, and alkalies readily breaking them up; (3) in certain animals atropine slows, while in other animals it makes the pulse rapid. Temperature of the surroundings, too, alters its effects, and in small doses it has one action, in large doses another, and individual peculiarities must not be forgotten, hence the difficulty of deducing from clinical experience. I have known atropine produce no dryness of the throat and no flushing in one patient, while in another these symptoms were very marked.

In favour of there being only one base we have the following considerations: (1) All authorities are agreed that the one base readily passes into the other, as, for example, atropine can be changed into hyoscyamine and hyoscyamine into atropine. (2) In experiments on the frog's heart every one can replace the other; stop the heart with muscarine and any one of them will start it; stop the heart by simple exposure to the air and any one will start it—under certain limits, of course. (3) They all dilate the pupil and all increase the intraocular tension, and, although this latter property is denied some of them, it is well-nigh impossible to prove the negative. (4) They all possess hypnotic action. (5) They all react in the manner of caustic soda or potash towards alcoholic solution of mercuric chloride.

Each new mydriatic base, when introduced to the notice of the profession, has had some special property ascribed to it, but time and trial have generally proved it to be in most cases only another name for atropine. As well-known examples we have daturine and duboisine. In this country we now regard them as impure atropine. Neither has been employed extensively internally, although it must be mentioned that recently Belmondo found hypodermic injections of duboisine equal to hyoscyne, and Mazzochi and Antonini believe it to be superior to atropine and morphine. Hyoscyamine in some respects has been less studied than the other members of the group, and most people seem afraid to hazard an opinion whether it is or is not identical with atropine. All are agreed that at least it is closely related to atropine. A curious point is that the source does not determine whether the product be hyoscyamine or atropine but rather the manner of manufacture. All agree that at least one is a conversion product of the other, but it does not matter which you start with; if you have atropine and wish hyoscyamine you have only to convert it, and if you have

hyoscyamine and wish atropine conversion is easily accomplished. Another interesting point must be mentioned. Samples of supposed atropine, employed as such and giving atropine results, have by experts been declared to be hyoscyamine. A further point of interest is that daturine has by authorities been declared identical with both atropine and hyoscyamine; if this be so we must infer that atropine and hyoscyamine are identical with one another, that is if expert opinion is worth anything at all. We must here mention that we mean the white hyoscyamine, not the brown amorphous substance which is a calumative in delirium.

Hyoscyne has not been sufficiently long before the medical world for a definite opinion to be formed regarding its effects and therapeutic value. Different samples may vary and account for the varying results obtained. I have used the drug in the form of hydrobromate both clinically and pharmacologically, and am unable to think it differs in any way from atropine. Its hypnotic action may be more marked, but I do not know but what I could have obtained the same results by means of atropine. It certainly produces dryness of the throat, rapid pulse, flushing of the face, and delirium (in fairly large doses).

Scopolamine is the newest member of the atropine group, and is only now on its trial, so that all the evidence one can collect is important. My own experiments with scopolamine hydrobromate were repeated again and again, and every effort made to eliminate error, and I gave the results in detail, so far as is necessary, to illustrate every point. From the hydrobromate the pure alkaloid was obtained and compared with pure atropine in its behaviour towards alcoholic solution of mercuric chloride, and no difference could be detected. One-hundredth of a grain of hydrobromate placed on the frog's heart, the pericardial sac being opened, caused diminution of the ventricular contractions. Another frog's heart was stopped by the application of muscarine nitrate. It commenced to beat strong and regularly after $\frac{1}{100}$ of a grain of the hydrobromate had been applied, and the subsequent exhibition of muscarine in the same heart failed to produce stoppage. A frog's heart, allowed to stop for some hours, did not start when $\frac{1}{100}$ of a grain of the hydrobromate was applied, but did beat when the same quantity of atropine sulphate was exhibited. (See later trials.) At another time a frog's heart poisoned by muscarine applied well over it failed to beat when scopolamine hydrobromate increased to the $\frac{1}{2}$ of a grain was applied. Atropine sulphate freely applied also failed to bring about heating of the organ. On the heart-rate the following notes were made. A frog's heart beating regularly 38 in the minute was brought down to 29 in five minutes by $\frac{1}{100}$ of a grain. In an hour another $\frac{1}{100}$ of a grain was instilled, and the rate became 28. Atropine sulphate in 1 per cent. solution was afterwards applied, and the rate fell to 26.

Some weeks later than the foregoing the following trials were made.

As before a frog was pithed, the cord destroyed, the pericardial sac opened, and the animal allowed to lie till the heart had ceased to beat spontaneously. The $\frac{1}{100}$ of a grain of scopolamine hydrobromate applied in five minutes caused it to beat afresh. The same heart standing overnight had ceased beating, and another $\frac{1}{100}$ of a grain started the heart, the contractions being chiefly in the sinus and auricles, while still another $\frac{1}{100}$ of a grain caused the ventricle to contract more powerfully, at the same time increasing the contractions in the auricles and venous sinus. In three hours atropine sulphate was applied freely, but the contractions remained the same in character.

One-hundredth of a grain instilled in my own eye at 10 one night produced complete dilatation by 7 the next morning. In twenty-five hours after complete dilatation the diameter had become reduced by half; and in sixty hours after complete dilatation the pupil had nearly, but not quite, returned to normal. The $\frac{1}{100}$ of a grain taken by the mouth on myself gave dryness of the throat, some slight hypnosis, and some slight slowing of the pulse. In a month later $\frac{1}{2}$ of a grain was taken, and there was some acceleration of the pulse, distinct mydriasis, hypnotic tendency, very dry throat, and some confusion of the senses. Some ten days later the $\frac{1}{2}$ of a grain was swallowed, and the noticeable symptoms were: the pulse was at first more rapid, then slower

than normal, but full and bounding all along; throbbing was felt in the temples; the throat and nostrils were disagreeably dry; the sight was blurred and the pupils markedly dilated; the gait was staggering, and there was confusion of the senses; the face was flushed, and there was a tendency towards sleep, but the uncomfortable conditions prevented rest for any considerable time. In less than twelve hours all discomfort had passed off, and no ill effects remained. (Results have been detailed as they occurred, and without comment.)

All my experiments with scopolamine were conducted with the salt supplied to the profession by Messrs. Burroughs, Wellcome, and Co. The temperature at which the experiments on frogs were carried on was 65° F. In the cases where the salt was applied the heart muscle was moistened by normal saline solution, and a tablet of the named strength laid on the surface of the heart and allowed to slowly dissolve.

To gather up the various threads of our essay. Daturine and duboisine we might call impure atropine. The varying effects obtained by different observers in the case of hyoscyne and scopolamine may be accounted for on the supposition that neither is a definite base. Hyoscyamine we might further suppose to be identical with atropine. Atropine is the alkaloid of which we know most, and we might call it the fundamental or representative base; and our knowledge permits us to say at least that in hyoscyne and scopolamine we have a certain proportion of atropine along with a varying proportion of a conversion or decomposition product of atropine, and which one might call a tropeine.

Instead, then, of multiplying names, we should call atropine what is usually atropine, and get hyoscyne and scopolamine free from any admixture, and name the resulting products by names corresponding to hematropeine. For my own part, as I have already said, my experience of hyoscyne, scopolamine, daturine, and duboisine has led me to think they differ little from atropine but other observers think different, and for this reason I have endeavoured to indicate the lines upon which investigations might be carried on, and by means of which we might arrive at the truth.

REFERENCES.

Y. B. P.=Year Book of Pharmacy. For hyoscyamine, Y. B. P. for 1880 and 1881. For duboisine, *Lancet*, 1878; *Pharm. Journ.*, 1878; Y. B. P. for 1881, 1885, and 1894; *Nouv. Rem.* 1893; EPITOME, BRITISH MEDICAL JOURNAL, March 23rd, 1895. For daturine, Y. B. P. for 1880, 1881, 1882, and 1885. For hyoscyne, Pavloff in *St. Petersburg Med.-Chir.-Dissert.* for 1889 and 1890; Y. B. P. for 1892 and 1893; O. Hesse, in *Liebig's Annalen*, cclxxvii, 304, 308. Sharp, in *Practitioner* for January, 1894. For scopolamine, Y. B. P. for 1890, 1892, and 1893; O. Hesse, in *Liebig's Annalen*, as under hyoscyne; O. Hesse, in *Apotheker Zeitung* for 1895; Raehlmann, in *Semaine Médicale* for 1893; Bokenham, in BRITISH MEDICAL JOURNAL for 1894, vol. ii, p. 597; T. R. Pooley, in *Canadian Lancet* for January, 1895. (The abstracts of the Year Book of Pharmacy are so full and so exact that, in order to lessen the labours of those who wish to examine the literature of the subject, I have largely quoted that work.)

LORETIN: A NEW ANTISEPTIC.¹

By HERBERT SNOW, M.D.LOND., ETC.,
Surgeon, Cancer Hospital.

LORETIN is an organic iodine compound, discovered by Professor Claus, of Freiburg. Its formula is $C_9H_4IN.OH.SO_3H$, and its euphonious proper title in chemical nomenclature meta-iodo-ortho-oxyquinoline-ana-sulphonic acid. It is a bright yellow odourless crystalline powder, very slightly soluble in water and alcohol; cold water takes up 1 to 2 parts per 1,000, boiling water 5 to 6; it is insoluble in ether and oils, forms emulsions with the latter and with collodion. Being an acid it forms neutral salts with sodium and potassium, with ammonium and magnesium, which are readily soluble in water, forming solutions of a deep orange-yellow colour; the neutral calcium and barium salts are only slightly soluble. The manufacture process consists in boiling together in water equal parts of potassium iodide, potassium carbonate, and oxyquinoline-sulphonic acid with chloride of lime, representing one atomic equivalent of chlorine. After cooling a little hydrochloric acid is added. Thus is formed the calcium salt of loretin, which is again decomposed with hydrochloric acid to obtain the crude drug,

subsequently carefully purified. The pure acid is extremely stable for an iodine compound, being unaffected by prolonged exposure to air or even to direct sunlight; it is easily decomposed only by oxynitrogen compounds, by free chlorine and bromine, or by certain organic compounds already undergoing decomposition.

Ammelburg administered, for several weeks in succession, loretin to dogs in doses up to 10 g., rabbits to 5 g. Hypodermic injections amounting to 5 c.cm. of a 5 per cent. solution of the sodium salt of loretin, were made daily into guinea-pigs for long periods. In no case did any objectionable symptoms follow; and in the urine, which was carefully collected by means of specially constructed cages, no iodine, blood, sugar, or albumen. These physiological experiments were continued by Professor Albrecht in the Veterinary High School at Munich, the results being published in the *Deutsche Zeitschrift für Thermedizin und vergleichender Pathologie*, 1894, vol. xx., p. 353.

Ammelburg's bacteriological investigations proved loretin to be a powerful microbicide, much superior to iodoform, with which comparison more particularly holds weight. They deal with the micro-organisms of cholera, anthrax, suppurative, typhus, etc. Some of the more significant tables are appended; the experiments are being still continued.

Schinsinger has extensively used loretin for burns and operations of all kinds, including empyema, herniotomy, resection of the upper jaw, arthrotomies, carious and tuberculous processes, etc.; in six months he had not a single instance of toxic symptoms, much less of death. Feussling has largely employed it in veterinary practice, with the most favourable consequences. The former remarks: "The absolute absence of any irritant effect upon the skin is a very important advantage. Artificial erythema or eczema I have never yet observed caused by it; but, on the other hand, very persistent eczema has been cured by loretin." Extensive burns were healed by dusting with loretin, and left very slight scars.

Loretin is recommended for use as a dusting powder, either alone or mixed with calcined magnesia, starch, or French chalk; as collodion (2 to 10 per cent.); in pencils of cocoa butter (5 to 10 per cent.); in ointment, 5 to 10 per cent., with vaseline or lanoline; in solution of 0.1 to 0.2 per cent. of the free acid, or 1 to 2 per cent. of the soluble sodium salt; lastly, as gauze impregnated with precipitated calcium salt. My own experience is confined to the powder, which I have never found occasion to mix with any other substance. Dusted on the skin, or over a granulated wound, this causes not the slightest irritation or unpleasant sensation. It immediately destroys the malodour of the most fetid cancerous sore, controlling this in a manner which no other agent I have yet tried will do. Copiously puffed with an insufflator into the deep cavity formed by evacuating the axilla of carcinomatous glands, it efficiently precludes suppuration, even when free hæmorrhage has taken place after the closing of the wound, an occurrence almost inseparable from anæsthetic vomiting when the patient has been removed from the operating table. Not the slightest bad symptom from its employment in this way has so far been detected. When there is no deep cavity a wound dusted with loretin heals rapidly by first intention. I have had recourse to loretin in some 60 cases, mainly operations on the breast and axilla, notoriously a test region for antiseptics. In my hands it has proved an ideal antiseptic and deodorant, with no single drawback; and I am sure that no surgeon who has once tried it will ever again resort to the noisome and toxic iodoform, from the free use of which I have seen more than one death. Though whenever old-established agents answer sufficiently the purpose I have a strong prejudice against novelties, yet this substance—non-poisonous, devoid of smell, and absolutely preventive of suppuration—seemed to me so marked an advance upon anything previously brought forward, that I felt constrained to direct thereto the notice of the Section.

APPENDIX (Abstract of Dr. Ammelburg's Bacteriological Experiments).

In each 20 c.cm. of a solution or mixture containing 2 per cent. of one of the following disinfectants was employed: beta-cresol, lysol, carbolic acid, loretin, iodoform, as well as a control experiment. In some cases 20 c.cm. of a 2 per mille

¹ Read in the Section of Pharmacology and Therapeutics at the Annual Meeting of the British Medical Association held in London, July-August, 1895.