April 4th. Since the above was written the patient has gained strength and has left the hospital. He is very emaciated and the malignant growth is rapidly increasing, but he has no abdominal pain and no sickness, and takes ordinary food. The button has probably dropped back into the disused portion of the intestine.

REMARKS ON THE USE OF THE BUTTON AND THE TREATMENT OF INTESTINAL OBSTRUCTION.

Mr. Mitchell Banks, in the *BRITISH MEDICAL JOURNAL* of February 23rd, has referred to Murphy's paper in the *New York Medical Record*, and to the few recorded cases in which the button has been used in this country, and pointed out its advantages, so that it only remains for me to add my testimony in its favour as a mechanical appliance. The case which I now record shows that even during intestinal obstruction an anastomosis can be successfully performed by its aid. Murphy advises an end-to-end approximation wherever possible rather than a lateral. In this case I chose the latter, as it avoided the necessity of suturing the divided ends of both the large and small intestine, a proceeding which would have made the operation a lengthy one. It is interesting to note that my patient is even younger than Dr. Hamilton's, one of the cases recorded in the *JOURNAL* of February 23rd, in which Murphy's button was used after excision of cancer of the colon, in which the patient was 33 years of age. I showed at the Pathological Society in 1893¹ a columnar epithelioma of the small intestine from a young woman only 23 years of age.

The importance of not masking the symptoms of intestinal obstruction with opium, or at any rate of not being misled by the effect of opium on the symptoms to which Mr. Mitchell Banks calls attention,² is again and again overlooked, and patients die because, until they are practically moribund, they are not considered bad enough to be operated on. Then, when without pain or vomiting, prostration reaches its climax, the death is called a sudden one, and the practitioner asks, "Who could have expected it?" The symptoms were not urgent, because subdued by the morphine, although the abdominal distension may have been increasing all the time. Dr. Hamilton³ quotes Murphy's statement that in cases of intestinal obstruction waiting for indications for operation generally means "criminal procrastination." It is a strong term, but it may perhaps call attention to the fact that an acute intestinal obstruction needs surgical interference at once, as surely as a strangulated hernia, whether the pulse seems good or not, and that waiting for indications of advancing prostration, or increase of symptoms is in most cases putting aside the one chance left to the patient—early operation.

NOTES OF A CASE OF ENTERECTOMY FOR GANGRENOUS INTESTINE IN WHICH THE BUTTON OF MURPHY, OF CHICAGO, WAS SUCCESSFULLY EMPLOYED.

By JAMES MURPHY, M.A., M.D.,
Surgeon to the Sunderland Infirmary.

Mrs. E. C., aged 27, married seven years, nullipara. was sent in to the Sunderland Infirmary by Dr. Spencer, of Hetton-le-Hole on October 31st, 1894, having had symptoms of acute intestinal obstruction for six days. She had obstinately refused operation.

Her previous history was to the effect that she was confined to bed three years previously for three weeks with pain in the left loin, and for seven weeks more she was unfit for her household work. Twelve months later she was again confined to bed for three weeks, owing to pain in the front and left side of the abdomen, and, after getting up, her legs swelled for some time. In neither attack had she any symptom of obstruction. Three weeks previous to her admission to the Sunderland Infirmary she had acute pain about the umbilicus, which lasted for a few minutes and was relieved by fomentation. On October 25th, at 9 P.M., severe pain set in

in the umbilical region, vomiting occurred freely, she was much collapsed, and there was slight action of the bowels. The latter had acted naturally on the previous day. After this there was no further passage of fæces or flatus; there was constant vomiting of brown fluid, with an offensive but not faecal odour. The collapse continued and the pain was severe, being acerbated by paroxysms every now and then. Her abdomen was distended, but not distressingly so; there was no rigidity of the abdominal muscles and no visible coils; the pulse was fairly good; the tongue was dry and brown; the temperature slightly subnormal. No tumour could be felt on palpation, and percussion gave a resonant note all over, including the flanks.

Shortly after admission an incision 2 inches long was made in the middle line of the abdomen midway between the umbilicus and pubes, and exit was given to some 30 ounces of blood-stained serum. On introducing two fingers, a loop of small intestine about 6 inches long was found strangulated by a tight band. It was impossible to bring the band to the surface, and it was gently torn through with the finger-nail. The strangulated loop was then brought outside the abdomen, when it was found to be gangrenous and perforated in three places, the largest perforation being at the point of strangulation. It was also found that about a tablespoonful of fluid fæces escaped through the perforation into the abdomen after the separation of the band. The fæces were immediately sponged out and the abdomen irrigated. Lane's clamps were introduced 2 inches above and below the gangrenous portion. An end-to-end anastomosis was then formed by means of Murphy's button, the bleeding points in the mesentery were secured, and its divided portions were brought together by means of a few sutures. The intestine thus united, the clamps having been removed, was returned into the abdomen, which was again thoroughly washed out, a glass drainage tube introduced, and the abdomen closed. The time occupied in excising the gangrenous portion of the intestine, inserting and fastening the button, sewing the mesentery, and securing bleeding points was four minutes and a-half. She was in such a collapsed condition that she could not be removed from the operating table for some time. Brandy was administered by enema and she passed a restless night.

Next day she was collapsed and suffered a good deal of pain. She passed flatus on the second day, the tube was removed on November 4th and all went well till November 9th, when she was seized with intense abdominal pain and collapse, and looked as if she were about to die. After a full dose of morphine she got relief; there was nothing to record till November 18th, that is, the nineteenth day after operation, when she passed the button, which she had felt in the rectum for several hours. The button was passed sideways with an adherent mass of fæces. All the holes of the button were blocked by fæces, and the two chromicised catgut ligatures were still *in situ*, but there was no portion of intestinal slough. She was allowed out of bed in a few days, and in a week's time she left the infirmary, the wound, which was less than 2 inches, had healed firmly.

The button used was the largest size for the small intestine. This is the only case in which I have so far used the button, but I was exceedingly pleased with the easy and rapid manner in which it can be used, and its efficiency is shown by the result. The great saving of time is the most important feature of this exceedingly ingenious instrument. Owing to the state of collapse this woman was in she could not have stood an enterectomy by any of the older methods, and therefore she owes her life, not to Murphy of Sunderland, but to his distinguished namesake of Chicago.

CASE OF ENTERECTOMY IN WHICH MURPHY'S BUTTON WAS USED.

By DONALD D. DAY, B.S., F.R.C.S.,
Assistant Surgeon, Norfolk and Norwich Hospital.

MURPHY'S button is being widely recommended, and cases in which it has been used should be laid before the profession, whether wholly or partially successful. In the *BRITISH MEDICAL JOURNAL* of February 23rd two cases are

¹ *Trans. of Path. Soc.*, 1893.

² *Loc. cit.*

³ *Loc. cit.*

reported by Mr. Mitchell Banks and one by Dr. Hamilton, who both so fully describe the procedure that I need not recapitulate it, but will merely describe a case in which I used it:

G. B., aged 27, a healthy stoker, consulted me on July 7th, 1894; five weeks previously he had been seized with acute colic and sickness whilst at work, without diarrhoea or jaundice. He had always been rather constipated, and was in the habit of taking Epsom salts. Since the first attack he had had several similar ones. The day previous to my visit he had a very severe one, with vomiting and constipation and severe pain in the lower abdomen, aggravated by movement of the legs and by micturition. He was lying on his back with legs extended; the tongue was dry and brown, and there were sordes on the teeth and lips. His expression was anxious; respiration was entirely thoracic, the pulse was 120, and the temperature 101.5°; the skin was dry, the abdomen rigid, generally tympanitic, and excessively tender, more especially below the umbilicus, but not more on the right side than the left. By the rectum a painful swelling could be felt in Douglas's pouch. The bowels had acted slightly at 1 P.M. The urine was scanty and concentrated.

As he was evidently suffering from peritonitis, I sent him to the Norfolk and Norwich Hospital, and, in the absence of the surgeon of the week, he came under my care.

During the evening the abdomen was opened in the linea alba, and some thin sero-pus escaped; a small band connecting the intestine to the bladder was divided. The bowel 3 feet above the cæcum was found adherent deeply in Douglas's pouch. In breaking down the dense adhesions two or three distinct pockets of thick pus were evacuated, and the loop of bowel was drawn up to the wound. It was for 1 inch much thickened and contracted, apparently from dense organised lymph, and the surface was ragged from the broken down adhesions. This, perhaps, might have been left alone; but, with the possibility of its being a malignant stricture and the uncertainty as to the size of the lumen of the bowel, I decided to remove it. Accordingly, about 10 inches of bowel were cut away with a small portion of mesentery, and the ends were quickly and easily joined with a medium-sized Murphy button, the bowel being compressed meanwhile by an assistant's finger. The mesentery was united on each side with a continuous silk suture, which was carried up to the periphery of the bowel so as to support the line of union. The peritoneum was well flushed with hot water and the abdomen closed, a glass tube being inserted.

The after-progress of the case was good (the bowels acting on the fourth day) until August 7th, when some offensive pus escaped through the track of the tube, and on August 19th faecal matter flowed freely. This continued at intervals, and evidently the button was causing partial obstruction. It was dislodged by a dose of castor oil, and passed *per rectum* on September 12th. On the next day a ligature escaped through the wound. No more faecal matter passed, and the sinus contracted though it did not finally heal until January. There has been no obstruction since; the appetite is very good and the bowels act regularly.

The bowel removed was found to be contracted for about half an inch to the size of a No. 10 catheter by dense fibroid material, which could be stripped off the peritoneal coat. The mucous membrane was apparently quite healthy.

This case was communicated to Dr. Murphy privately, and has been entered as No. 65 in his last published list. The point of special interest is the long delay in passing the button—sixty-six days—which is, I believe, the longest on record. Dr. Murphy specifies that cases of existing peritonitis are unfavourable for satisfactory union of the bowel, which one would quite anticipate.

What was the cause of the partial failure in this case? Probably it was the peritonitis. It may have been from the button fitting a little too tightly to pass easily along the remainder of the ileum; it may have been from the supporting suture allowing of infection from within the bowel. Another time I shall certainly trust to the button without additional suture.

One more point of interest. What was the cause of the peritonitis? The vermiform appendix was free from reproach. It is possible that a small gland may have suppurated and

caused local peritonitis, but no history suggestive of it could be obtained; the mucous membrane was intact, and no trace of primary gland abscess can be found in the specimen. Can peritonitis be idiopathic? We know of purulent deposits in other serous membranes—the pleura and pericardium—which are apparently idiopathic. We are not taught that the pelvic peritoneum may behave in a similar way, but I see no reason why it should be exempt from trouble which may befall other serous membranes.

SOME REMARKS ON THE TREATMENT OF FRACTURES.

By W. ARBUTHNOT LANE, M.S.,

Assistant-Surgeon to Guy's Hospital and the Hospital for Children, Great Ormond Street.

IN considering the influence which a fracture may exert on the subsequent life-history of an individual, we must recognise one great difference between the results of fractures of the long bones of the upper and lower extremity.

In the case of the upper, the superjacent weight of the body is not transmitted through the joints as it is in the leg, therefore an alteration of the mechanics of a joint in the arm in consequence of fracture of one or more bones does not determine the development of such extensive changes as take place in the leg. Yet in the arm very considerable alterations may occasionally develop, and are more marked and depreciating to the value of the individual as a machine in proportion as changes have already taken place in the particular joint or joints from the prolonged pursuit of a laborious occupation.

We may conclude that in fractures of the long bones of the upper extremity it is not of such vital importance to reconstitute the damaged skeleton in its original form as it is in the case of the leg.

In fractures especially of the femur, tibia, or fibula, nothing plays a more important part in the prognosis as regards the presence or absence of subsequent pain and depreciation of the physical capacity than does "age."

In the comparatively young adult, deformity, with the consequent alteration in the physiology of the limb, brings about what may in the absence of a better term be called "adaptive" changes in the anatomy of the joints influenced. By this term I mean an alteration in the form and perhaps in the plane of the articular surfaces, with a variation in the extent of the synovial membrane and in the attachments of the ligaments, and these are usually unassociated with any pain or discomfort at any period subsequent to the receipt of the fracture. After a certain age, which varies within wide limits with the habits, occupation, surroundings, etc., of the individual, which may be roughly averaged as about 40 years, this being probably an excessive estimate for labouring people, a different series of changes takes place. These may for convenience sake be called "passive" changes, and are included under my description of mechanical or traumatic arthritis.¹ They consist in the removal of articular cartilage at the seat of abnormally great pressure, with subsequent eburnation and destruction of opposing bony surfaces, the alteration in the extent of the articular surfaces by the marginal deposit of cartilage and bone, and increase in the thickness and character of the synovial membrane with the secretion of an excessive amount of synovial fluid—in fact, just such changes as used to be regarded by the pathologist as indicative of the presence of a disease which he called by various names, that, perhaps, most commonly used being rheumatoid arthritis.

Clinically this means that in the slightest forms the patient experiences a feeling of insecurity and discomfort in the affected part, and in its marked forms a degree of pain that sooner or later absolutely incapacitates the sufferer from transmitting the weight of the body through the joint or joints of the leg for any useful purposes.

I propose to limit my remarks to fractures of the lower extremities. I would point out that as regards the mechanical treatment of fractures other than those that are transverse, and in which there is no displacement of fragments, or in

¹ Chronic Traumatic Arthritis, *Lancet*, January 30th, 1892.