indispensable. Baths of weak solutions of liq. carbonis deterg. were also most grateful to the patient. He had seen many cases occur in those who engaged in polo and cycling.

VI.—F. H. BARENDT, M.D.Lon¹, F.R.C.S.Eng.,
Liverpool.

DR. BARENDT considered that local applications were most important, especially in those cases where pruritus alone was present, and there was no skin lesion that could explain it. He extolled the application of blisters over the spinal region which supplied the pruritic area. After having tried several local remedies—resorcin, menthol, salicylic acid in ointment and paste form—he had obtained great relief by thoroughly rubbing into the affected area hot olive oil containing 2 per cent. of carbolic oil, for five minutes night and morning.

VII.—ELIZABETH GARRETT ANDERSON, M.D.Paris,
Physician, New Hospital for Women, London.

MRS. GARRETT ANDERSON said that in treating neurotic pruritus in women, and men leading feminine lives, she had found that an increase in the quantity of food which suited the patient gave good results. Rest, especially before meals, was important. Bathing was ill-borne in these cases. She also recommended complete silk underclothing. Cod-liver oil should be added to the food if necessary.

PROFESSOR MCCALL ANDERSON'S REPLY.

IN his reply, Dr. McCall Anderson said that the treatment of the cause was the important point, but he did not undervalue local treatment. He considered that the good influence of Harrogate, for example, was not due to the baths, but to the diet, and also to the healthiness of the surroundings. He agreed with Mrs. Anderson's remarks as to the importance of keeping up the nutrition.

AFFECTIONS OF THE SKIN OCCURRING IN THE COURSE OF BRIGHT'S DISEASE.¹

By P. H. PYE-SMITH, M.D., F.R.S.,
Physician to Guy's Hospital.

THE only cutaneous disorder hitherto generally recognised as occurring in Bright's disease is the erythema leve of Willan. His original description in 1808² was followed by Bateman and subsequent writers. He notices its uniformly smooth and shiny surface, the red patches becoming confluent, and its association with anasarca. He describes it, however, as occurring both in young and old patients, particularly in those who are intemperate, and as not always accompanied by dropsy. He also observes that the apparently slight inflammation may end in local gangrene.

Erythema leve is a superficial dermatitis which owes its characteristic smooth, shiny appearance to the fact that it appears on surfaces made thin by dropsy. It may be the result of local irritation from pressure, and the term has been applied to the first stage of a bedsore. But as it appears on the dropsical limbs there is no evidence of external irritation in its locality or spread, and it is seldom seen in the anasarca of cardiac or hepatic disease. It is also quite distinct from the local recurrent erythematous or erysipelatos dermatitis which leads to elephantiasis of the legs. So restricted, the term may still be used to describe the inflammatory blush which sometimes appears on the legs and thighs swollen by renal dropsy. In its pathology it should, however, be separated from other forms of true erythema—E. multiforme, E. iris, E. bullosum, E. nodosum, and urticaria—and regarded as a dermatitis readily excited on a dropsical skin, just as pleurisy, inflammatory oedema of the lungs, and sometimes inflammation of the pericardium, the larynx, or the peritoneum are observed in Bright's disease. There are, however, other forms of dermatitis which not infrequently occur in the course of chronic Bright's disease:

1. There is a bright red diffused rash which appears chiefly

on the trunk, less extensively on the neck, arms, and thighs, and very seldom on the face, hands, or feet. It is distinguished from the somewhat similar rash produced by natural or artificial diaphoresis, by its locality, by the absence of sudamina, and by its appearing when no hot air baths, or other means have been used to produce sweating, and when the skin is harsh and dry. It does not, as a rule, either itch or smart, and only lasts a few days. I have most often seen this rash in cases of chronic tubal nephritis.

2. There is a papular eruption with large discrete rather dark red pimples seated on a dry, rough, and sometimes scaly surface. This I have more often seen on the outer side of the thighs and legs, the shoulders and extensor surface of the forearms, but it also may affect the loins and the abdomen. I have never seen it on the face or on the hands and feet.

3. Apart from the mere coincidence of eczema with Bright's disease, there may be observed in some cases a moist dermatitis resembling eczema in its respect, but occupying the arms or the legs, without affecting the flexures of the joints, the face or the ears, without the irritation commonly present, and without having previously appeared.

4. On two occasions I have seen a very extensive and profuse dermatitis closely resembling the universal exfoliative dermatitis of Wilson, very red, very scaly, occupying the scalp, palms, soles, and genitals, as well as the trunk, face, and limbs. It has come on after the symptoms of Bright's disease have appeared, in cases of chronic interstitial nephritis, with little dropsy, and cardio-vascular changes already apparent.

Different as these forms of dermatitis are in their anatomy, they have certain common characters apart from their occurring in patients suffering from Bright's disease. They are superficial, and leave no trace behind when they disappear or after death. They run an acute or subacute, not a chronic, course, and seldom recur after they have disappeared. They are never pustular. They are occasionally purpuric, but this is rare, offering a contrast with the hæmorrhagic erythema of rheumatism and erythema nodosum. The subjective symptoms are usually slight. With respect to prognosis, they usually occur in the later stages of the renal disease, and therefore may be not infrequently present at the time of death. But they often disappear long before this, and their appearance does not coincide with any special aggravation of the other symptoms. As to their pathology, I am not inclined to connect them with uræmia except by a natural coincidence, for we constantly see this condition in its most characteristic and fatal form without any cutaneous eruption, and each of the forms of dermatitis I have described has been seen where there were no other symptoms of uræmia. Rather, I think, should these inflammations of the skin be associated pathologically with the retinitis, the serous inflammations, and the pulmonary oedema to which patients with Bright's disease are liable. With respect to the form of renal disease with which they are associated, I do not remember seeing any of them in the acute cases of nephritis with dropsy, either in children or adults; nor in cases of lardaceous disease of the kidneys. They have either occurred in the later stages of chronic tubal nephritis, or at any stage in the course of chronic granular degeneration with shrinking of the renal cortex.

The treatment I have found useful when local discomfort has called for it has been entirely external, and the vehicle an ointment rather than a lotion. Carbonate of lead, oxide of zinc, calamine or bismuth, have appeared to be the most useful applications, and sometimes inunction with olive oil or vaseline has been more efficient than any more active treatment.

In conclusion, I would remark that none of these affections (except erythema leve) are common, and my chief object in drawing attention to the subject is to elicit the experience of those who, I doubt not, will be able to fill up and perhaps to correct these imperfect outlines.

I must, however, particularly refer to a valuable paper by Dr. Le Cronier Lancaster, which was read before the Clinical Society.³ It had escaped my notice until the above account had been drawn up, and I have not modified any of the statements which differ from those made by Dr. Lancaster, because

¹ Read in the Section of Dermatology at the Annual Meeting of the British Medical Association held in London, July-August, 1895.

² *Cutaneous Diseases*, vol. i, p. 475.

³ *Transactions*, vol. xxv, p. 49, 1892.

by comparing our individual experience we shall best arrive at just clinical and pathological conclusions.

Dr. RADCLIFFE CROCKER, President of the Section, insisted on the importance of drawing attention to forms of dermatitis due to disease of the internal organs. In his experience he differed from Dr. Pye-Smith as to prognosis, which was gloomy in advanced cases of Bright's disease. He agreed with Dr. Le Cronier Lancaster.

Dr. SAVILL (London), referring to the exfoliative skin diseases mentioned by Dr. Pye-Smith, said that his subsequent experience of the epidemic skin disease led him to believe that renal disease formed a predisposition to contract the former. Out of five or six cases of such complication which he had seen, all were fatal. He regarded it as a very serious sign.

Dr. BRADBURY (Cambridge) had seen several cases of skin diseases occurring towards the end of morbus Brightii. He had observed erythemata, and one case was markedly urticarial. In cases of interstitial nephritis he thought the element gout, *plus* poisonous matter retained in the blood through imperfect elimination by the kidneys, might be a factor. He admitted this did not explain all cases, especially those of chronic tubal nephritis. He was of opinion that the prognosis in these cases was unfavourable—that the end was not far distant.

Dr. WALDO (Clifton) asked Dr. Pye-Smith whether he had often noticed that the serum after blistering cases of chronic Bright's disease was hæmorrhagic.

Dr. GRANGE had noticed urticaria.

In his reply, Dr. PYE-SMITH did not think there was any evidence of uramic origin; he had never seen skin affections in very bad cases of Bright's disease. Eruptions did not give him that grave augury as to prognosis which had been mentioned by other speakers. He had never noticed urticaria as seen by Dr. Grange. He was extremely sceptical as to the influence of gout. In reply to Dr. Waldo, he had had no experience of blistering leading to hæmorrhagic serum; in this respect he would point out that in Bright's disease one should never give opium or mercury, or apply a blister.

ON THE GUINEA-WORM.

By PATRICK MANSON, M.D. ABERD., F.R.C.P. LOND., ETC.,
Physician to the Seamen's Hospital, Greenwich.

WHEN mature this singular integumental parasite—*dracunculus medinensis* or guinea-worm, of which the female form alone is known, attains an enormous length—2, 3, or 4 feet. Though long, she is very slender, having a diameter of about the tenth of an inch. Her anatomy is very simple; practically she may be described as a musculo-cutaneous tube, enclosing an exceedingly delicate uterus, which, packed with millions of coiled up, long-tailed embryos, extends from mouth to tail. The head of the adult worm is rounded off, and is provided with a punctiform mouth and a crown of minute papillæ; the tail end is furnished with a sort of hook which possibly functions as a holdfast. The alimentary canal is a minute tube compressed and thrust to one side by the uterus. Vagina and anus are wanting—at all events they have not been seen; probably they are obliterated by the pressure of the enormous uterus.

When the embryos in her uterus attain a certain degree of maturity, the worm travels slowly through the connective tissues of her host, proceeding almost invariably in a downward direction towards the lower extremities. She generally gets as far as the ankle or foot. Arrived there she drills a minute hole in the derma, the superjacent epidermis being raised up as a bleb. In a day or two the bleb bursts or is broken, disclosing a superficial ulcer with the little hole, leading to, or, more rarely, occupied by, the head of the parasite, at its centre.

The prevailing idea is that the worm now creeps out, or is pulled out, and that her body, being cast away, decomposes, and so liberates the embryos. Another idea is that the worm breaking down in the tissues, her young escapes in the purulent discharge which the irritation of her presence excites. It

can easily be shown, however, that neither of these views is correct. The worm gives rise to no purulent discharge so long as she is alive; nor, left alone, does she quit the body until she has first got rid of her young. This can be shown by the following experiment:

Squeeze a little cold water from a sponge so that the stream shall fall on the sound skin within an inch or two of a guinea-worm ulcer; at the same time watch the little hole alluded to at the centre of the ulcer. In a few seconds a droplet of a whitish fluid will be seen to well up in the little hole, or a delicate tube will be protruded from it for an inch or more, and then suddenly rupture. Collect a little of the fluid that has welled up from the hole or escaped from the ruptured tube, and place it under the microscope. It will be seen to contain hundreds of coiled-up, apparently dead, embryo guinea-worms. Instil a little water below the cover-glass and watch. In a few minutes the embryos seem to wake up, stretch themselves, move, and in a very short time they are swimming about vigorously. Manifestly they are in their proper element—water.

The explanation of these facts, which I have frequently witnessed and shown to others, is this. The parent guinea-worm instinctively makes her way to the foot or ankle because her young are most likely to get to water in that situation. In guinea-worm countries the natives wear neither shoes nor stockings, and their feet especially are frequently in puddles of water. In some strange way the guinea-worm recognises this fact, and also recognises when the foot of her host is actually in contact with the water. So she makes her way to the foot or ankle, and at the right moment, when the foot touches water, expels her young just as when we experimentally douche the limb she is lying in. I find she takes about a fortnight to empty her uterus in such intermittent efforts at parturition. When all her young have been expelled she begins to come out herself, and can then be more or less readily withdrawn, or she may break down and excite suppuration. I may mention that as the parasite has no vagina she gets rid of her young through her mouth; the contractions of the musculo-cutaneous wall, in response to the stimulus of water applied to the skin of her host, cause the uterus to prolapse through the mouth and burst. An inch or two of uterus prolapses each time the stimulus is applied; becoming very tense under the sustained contractions of the body of the worm, it ruptures and shrivels up and so on until the entire organ is in time expelled.

The object of all these curious arrangements is to give the guinea-worm embryo the best possible chance to get at its intermediate host—a fresh water crustacean, cyclops quadricornis. Fedtschenko, now a good many years ago, discovered that cyclops was the intermediate host of the *dracunculus*, and convinced Leuckart and other helminthologists of this. But, so far as I know, his experiments had not been confirmed or repeated until last year. Having a case of guinea-worm under my care at the Seamen's Hospital, I bethought me of testing Fedtschenko's statements, and also of seeing if the guinea-worm could be raised in England. I procured a large supply of embryos by the douching process I have described, and placed them in watch glasses and little bottles, together with a number of cyclops collected from ponds in the neighbourhood of London. They were placed together one evening. On the following morning, on looking at my cyclops with the microscope, I found that nearly every one of them had some ten or twenty guinea-worm embryos coiled up and wriggling about in their body cavities. The parasite had penetrated the armour of the cyclops by attacking it between the joints of the ventral plates; it does not get to the body cavity through the alimentary canal after being swallowed. I kept the cyclops alive for many weeks, and watched the slow metamorphosis of the incarcerated embryos. I repeated the experiment with the same success this year, as the preparations (for most of which I am indebted to Mr. Rousselot) and Mr. Andrew Pringle's very telling photographs show.

In this last set of experiments the parasites and the crustaceans were placed together on May 10th. I dissected, or rather crushed, a few cyclops from time to time, and so could follow the metamorphosis step by step. I found that soon after it enters cyclops the young guinea-worm changes its shape. When a free swimming animal the little embryos body is flattened from side to side, something like a very

¹ Read in the Section of Dermatology at the Annual Meeting of the British Medical Association held in London, July-August, 1895.