MALIGNANT DISEASE OF THE PANCREAS.

By GUTHRIE RANKIN, M.D. GLASG., F.R.C.P. EDIN.,
Physician to Warwick School.

FROM recorded observations of pancreatic lesions it would seem that primary disease is the exception rather than the rule, and that when the pancreas is attacked by morbid change of a malignant nature its implication is usually secondary to pre-existing disease in the stomach, duodenum, liver, or gall bladder. The following cases are therefore doubly interesting, because in both the disease was primary; in both it gave rise to a train of symptoms which *post-mortem* examination fully explained; and in both these symptoms failed to comply, in some essentials, with the classical descriptions to be met with in our best known textbooks.

CASE I.—C. L., aged 50, a painter, of small stature, ill-nourished, and dejected in appearance; declared he had never been ill in his life till towards the end of 1891, when he first came to me complaining of paroxysmal abdominal pain and of obstinate constipation. There was no evidence of organic disease anywhere, with the exception of some increase in the area of percussion dullness of the liver; but this was not accompanied by tenderness on pressure. There was a very decided blue line along the margin of his gums, and all the circumstances of his case seemed to point to his occupation as the most likely cause of his discomfort. His habits of life had been regular, but he was accustomed to a liberal daily allowance of beer. Nothing remarkable was revealed in his family history; both parents lived to well over 70, but the actual cause of death was unknown. Careful dietary and ordinary treatment soon relieved his colic, and after about three weeks' care he was able to resume his work. Four months later he again came under my observation with a similar succession of symptoms *plus* slight jaundice. The liver enlargement continued, and there was now tenderness along the whole of its lower edge. The pain was more intense, but again, after about three weeks of suitable treatment, it disappeared; the jaundice likewise vanished; the liver shrank almost to normal size; and the man returned to his trade. He remained well for two months, but was then obliged to give up work. The pain was acute and paroxysmal, he had lost all appetite, had become much emaciated, and exceedingly feeble. At first his condition improved under treatment, but at the end of a fortnight weakness increased to such an extent that he took to his bed. His liver was again increased in size and tender, but the surface was uniform and smooth. He had a rapid, soft, and shabby pulse, but there was no cardiac lesion. The lungs were not the seat of active disease, though there was evidence of previous inflammatory trouble. The other organs were healthy. The urine was copious, full of bile, and contained an intermittent haze of albumen, but no sugar. Shortly after taking to his bed he developed a most distressing cough, which, though less or more constant, periodically increased to paroxysms, which brought about extreme exhaustion. The expectoration was scanty, and muco-purulent in character. Occasionally he suffered from drenching perspirations. The blue line on his gums was much less distinct than eight months previously, and there was no return of the jaundice. At intervals he was attacked by the most violent pain, which, beginning at the pit of his stomach, rapidly radiated over the abdomen, and was only alleviated by large hypodermic doses of morphia. So severe were those paroxysms of pain, that the man's cries could be heard the length of the street in which he lived, and if much time elapsed before the administration of morphia he was brought to the verge of frenzy, no less painful to witness than dangerous to endure. In the course of his illness these acute seizures recurred five or six times only, but on each succeeding occasion the resulting exhaustion became more extreme, ending twice in syncope which threatened to prove fatal. There was no vomiting, not even nausea, to contend with, but diarrhoea was an increasing trouble during the last fortnight of his life. The evacuations were numerous—four to eight in twenty-four hours—but not persistently colourless, for the most part watery, but sometimes pultaceous and sticky, scanty in amount, and offensive; they never contained blood or mucus, and did not look fatty. Tenderness continued over the liver, and deep pressure in the epigastric region elicited expressions of special discomfort, while he often complained of a deep burning pain which he referred to his stomach. No nodule or lump was perceptible to the touch even at the end of the deepest expiration when the abdominal walls were well relaxed. He emaciated rapidly, was persistently despondent, and manifested the most remarkable prostration. He eventually died, in November, 1892, from asthenia. The usual remedies for the relief of symptoms were resorted to, but morphia was the only drug from which he derived comfort.

At the *post-mortem* examination on the day following the patient's death, it was found that the heart was thin in its walls, friable in its tissue, but without valvular defect. On both sides there was some congestion, evidently hypostatic, of the lower lobes of the lungs, and there were old pleuritic adhesions to the left parietal wall. The pyloric end of the stomach presented some thickening and infiltration, but the passage into the duodenum was patent. The liver was enlarged and full of dark blood; it completely overlapped the pancreas. The gall bladder contained some bile but no calculi. The pancreas was the seat of extensive disease; the head was enlarged, nodular, hard, leathery to the knife, and of a grey colour on section. Scattered here and there throughout the body of the gland were numerous deposits of varying size bearing exactly the same characteristics. There were firm adhesions to all the neighbouring structures, from which the pancreas could with difficulty be separated. The other organs were without pathological change.

CASE II.—E. H., aged 67, a farmer, of spare habit, but healthy in appearance, and of a ruddy complexion, had suffered for years from subacute attacks of typhilitis. Early in 1893 he was, for the first time, the victim of an attack of colic. This seizure, which was followed during the rest

of his life by many others of a similar character, seemed pathognomonic of gall stone. It was ushered in by sudden acute pain over the upper part of the abdomen, accompanied by vomiting, and followed by jaundice. From the date of the first attack of this colic he never again experienced trouble in the region of the caecal valve. His health otherwise remained tolerably good, and for about nine months immediately preceding the commencement of his last illness he was particularly well and able to look after the usual business of his farm, with occasional days off because of slight attacks of pain. In October of 1894 he had an unusually bad seizure of colic, and when the jaundice cleared off it was remarked that his appetite did not return as on previous occasions, and that there remained a certain amount of tenderness deep in the epigastrium. He suffered from imperfect digestion, and complained of such feebleness that he felt unable for more than the most moderate amount of exertion. He emaciated rapidly, and towards the beginning of December a sharp attack of pain compelled him to keep his bed on account of the debility which ensued. This seizure was not of long duration, but he believed it to be more severe than any preceding one, and, though it was followed by a marked increase of bile in the urine, there was no consecutive jaundice. The temperature had never risen till now, when for some days it reached 101°, and was accompanied by flushings, sweatings, flatulent distension, and violent throbbings of the blood vessels, more especially of the abdominal aorta. There was also rapid and excited action of the heart, together with some irregularity of the pulse. The tenderness in the epigastric region had become so intense that he could not bear to be touched, and for the first time the defined lower limit of a hard substance could, on favourable opportunities, be felt in the middle line below the ensiform cartilage. The lump was pulsatile, and on auscultation a loud systolic murmur could be heard, but the pulsation was not expansile, and gave the impression of a solid mass, through which the beats of the aorta were conveyed to the surface. When the febrile symptoms, which were attributed by the patient to his having been chilled through getting out of bed in the middle of a cold night, had abated, the tenderness became less and admitted of a more thorough examination of the abdomen. A distinct hard mass could then be felt in the middle line, extending towards the right side, where it became lost behind the edge of the liver. The liver itself was increased in size, but was tender only over the region of the gall bladder. The lungs were healthy. The heart was feeble, generally quiet and regular, but occasionally intermittent and very rapid in action, and there was a systolic bruit whose greatest intensity was situated in the pulmonary area. The urine at no time contained either albumen or sugar, but it was frequently thick from the presence of urates and deeply stained from an excess of bile. The bowels were uniformly constipated, and the motions, though usually normal in appearance, were twice or three times colourless and offensive, but never oily. There was no jaundice during the later stages of the illness. Once only, towards the end of life, was there a tendency to diarrhoea, but this was evidently due to faecal accumulation, and was speedily relieved by emptying the rectum. He had several slight attacks of sickness, but he was able to take a fair amount of liquid nourishment, which he both retained and absorbed perfectly.

In spite of these favourable conditions, his weakness increased daily until he could hardly lift his arm from the bed, his voice sank to a whisper, his emaciation became more and more pronounced, and his countenance assumed a pinched and cachectic look. He suffered no actual pain between the times of the paroxysms, but still disliked much handling of the upper part of his abdomen. In the middle of January, 1895, unexpectedly one morning, just after his nurse had finished putting him straight for the day, he complained of a feeling of cold and faintness, and, before his family could be called, he died.

His family history was unimportant; there was a record of several sudden deaths from "heart disease," but no case of malignancy was known in the connection. His own habits had always been careful and regular, and his whole life had been quiet and uneventful. The treatment of his case was carried out on general principles, and was successful in relieving his urgent symptoms as they presented themselves.

At the *post-mortem* examination the abdomen only could be examined, when it was found that the whole pancreas was enlarged and infiltrated with a hard nodular growth, the head of the gland being conspicuously the seat of morbid change. The liver was somewhat increased in size, but contained no pathological deposit, and the gall bladder was half empty, with a few small calculi lying loose within its cavity. There was evidence of recent localised peritonitis in the neighbourhood of the liver and duodenum, but there was no fluid in the cavity of the abdomen. The aorta was not dilated nor aneurysmal. The stomach was larger than normal, and its walls were unusually thin; the pyloric end was thick and indurated, but there was no obstruction to the onward passage of the stomach contents. There were some secondary hard deposits in the upper part of the duodenum a few inches below the pylorus, together with constriction of the calibre of the gut in this situation, and the pancreas was glued to the adjacent bowel. The appendix was dilated and full of thick, sticky, dark-coloured fluid, and the neighbourhood of the caecum showed evidences of previous inflammatory disturbance. The other organs were healthy.

The symptoms presented by both these cases—many of them most interesting—were, in the light of *post-mortem* experience, fully explained, but during life they gave rise to many doubts and difficulties in diagnosis. The leading positive symptoms in both cases were great emaciation, excessive weakness, paroxysmal pain, and in the later stages of the illness deep-seated epigastric tenderness, while in Case II there was the further evidence of a lump in the epigastrium; on the negative side there was an absence of fatty stools, glycosuria, prominent digestive disturbances, persistent jaundice, or malignancy in the family history.

So much stress has been laid on the presence of fat in the alvine evacuations as characteristic of pancreatic disease

that these two cases are instructive as tending to prove that, even in advanced conditions of structural disease of the organ, that symptom may be entirely wanting. The absence of colour from the motions has also been insisted on by some writers as indicative, in the absence of readily recognisable hepatic disease, of some interference with the pancreatic secretion. Claude Bernard taught that the coloration of the faeces was indirectly due to the action of the pancreatic juice, and Dr. Walker, of Peterborough, likewise came to the conclusion that a deficiency of pancreatic fluid would, as readily as a deficiency of bile, cause clay-coloured evacuations. If this were entirely true it is difficult to understand why in both these cases—and particularly in Case II, where the whole substance of the gland had undergone extensive change—the motions were not always colourless. Further, it would seem doubtful whether reliance can be placed on the theory of the formation by the pancreas of a glycolytic ferment, because sugar was invariably absent from the urine during the whole course of both illnesses. The absence of notable digestive disturbance and of persistent jaundice is specially striking in such pronounced types of pancreatic disease. The probable explanation is that, the fats being chiefly administered in the form of milk, that is, practically as an emulsion, were capable of direct absorption; and that the common bile duct escaped such pressure as would cause its complete occlusion.

In Case I the original history pointed to lead poisoning as the likely etiological factor, but *post-mortem* evidence suggests the suspicion that the attacks of colic, from the beginning, may have been due to pressure or inflammatory disturbance of the solar plexus. At any rate, there can be little doubt that in the later stages this was the explanation of the terrible paroxysms of acute pain which the unfortunate man endured. The enlargement of the liver might be explained by temporary vascular changes caused by the progress of the morbid growth. Another instructive symptom in this case was the distressing and fitful cough, which was probably due to irritation of the vagus nerve where it terminates in the solar plexus, while the drenching perspirations might also be the result of sympathetic nerve disturbance conveyed through the splanchnics. Intermittent pressure on the common bile duct would account for the jaundice, the emaciation and exhaustion might be looked upon as direct results of the dyscrasia, and the repeated syncopal attacks were either likewise due to this cause, or to interference with the cardiac nerve supply through the vagus or sympathetic nerves.

In Case II the early seizures were so apparently characteristic of the passage of gall stones that little doubt was entertained as to their nature; but again, *post-mortem* evidence makes one hesitate to ascribe them to this cause alone, even though a few calculi were found in the gall bladder; for might not the paroxysms be equally explained by the same cause which produced them in Case I? Gall stones were never actually found in the evacuations, and though their presence in the bladder made it likely that the patient suffered during life from the passage of one or more into the duodenum, it is equally possible that to some extent the painful seizures were due to changes going on in the pancreas, and producing pressure effects on the neighbouring nerve plexus; and in this connection it is to be observed that the later attacks of colic were not followed by jaundice. The abdominal distension, copious sweatings, flushings, and unusual rapidity of cardiac action would all be produced by disturbance of nerve equilibrium emanating from the site of pathological lesion. The hard lump felt in the epigastric region towards the end of the illness proved to be the lower edge of the pancreas, through which were transmitted the pulsations of the abdominal aorta, the pressure of the hard and enlarged gland against the large vessel being the cause of the loud systolic murmur; lump and murmur together, however, were strongly suspicious of aneurysm. The rapid emaciation and pronounced prostration must again be referred to the direct effect of the malignant growth on the vitality of the blood, because it was specially remarkable in this case that, throughout the whole course of the illness, nourishment in ample quantity was not only swallowed and retained, but perfectly assimilated.

The leading characteristic of both cases was the rapidity of the emaciation when it had once commenced, together with the degree and early manifestation of prostration, greater than that met with in any other form of malignant disease, and it suggests that the pancreas ought to be suspected when in a patient over 40—particularly in a male patient in whom the disease is said to occur most commonly— indefinite epigastric discomfort is accompanied by rapid emaciation and overwhelming weakness. If to these symptoms are added nerve disturbances which can be referred to the vagus or sympathetic, the probability of the pancreas being the *fons et origo mali* is strengthened; and if further corroborative testimony is afforded by a hard mass in the epigastrium, by glycosuria, by persistent jaundice without liver symptoms, by stools which are greasy in appearance, or clay-coloured without an obstructed bile duct to be found in explanation, then the diagnosis may be looked upon as practically certain, while the prognosis unfortunately is absolutely hopeless.

PULSUS PARADOXUS IN ACUTE LARYNGITIS.

By EDWARD M. BROCKBANK, M.D. VICT., M.R.C.P.,
Resident Medical Officer, Royal Infirmary, Manchester.

IN the BRITISH MEDICAL JOURNAL for June, 1893, I recorded some observations on the occurrence of *pulsus paradoxus* in cases of acute laryngitis. At the time of writing the paper, although I had seen several cases in children in which this physical sign was present, I had not been able to secure a pulse tracing showing it.

In February, 1894, however, a case of acute laryngitis came to the Royal Infirmary, Manchester, with the irregular pulse well marked, and from it I managed to take some good tracings. The following are a few brief notes of the case:

L. H., aged 6, was brought, on February 24th, 1894, to the accident room, suffering from severe dyspnoea, which the mother said had come on rather suddenly about thirty-six hours previously. The boy looked very ill, tossing about from time to time as he lay on his back on his mother's knee, throwing up his arms and gasping for breath. His face was dusky and cyanosed; the breathing was very laboured, especially during inspiration, and the extrinsic respiratory muscles were in action. The lower ribs were markedly drawn in with each inspiration. The respirations were about 45 per minute. The cardiac impulse could be seen through the chest wall and the heart was felt to be beating in a very heavy and laboured way. The pulse at the wrist was about 90 per minute, and was easily felt to be very irregular in the volume of the individual beats. It was almost imperceptible during inspiration. No membrane could be seen on the fauces. As the child was very ill, and from the mother's account getting rapidly worse, it was decided to perform tracheotomy.

The case was sent up to the theatre at once, and a small quantity of chloroform administered. Just before the trachea was opened I took two pulse tracings, one of which (Fig. 1) is reproduced here. In this the irregularity of the pulse wave is plainly seen, the smaller beats occurring synchronously with inspiration, and the larger ones with expiration.

Fig. 1.

The trachea was opened within twenty minutes of the patient's arrival at the hospital, and immediate relief to the urgent symptoms was obtained. The dyspnoea and cyanosis passed off rapidly, and the pulse became more regular, less laboured, and quicker. Figs. 2 and 3 are copies of tracings taken from the pulse one minute and two minutes respectively after the trachea was opened. In the former it will be seen that the character of the pulse is much altered from that in Fig. 1, being much more regular in the height of the upstroke. There still, however, occur in it some feeble beats, indicating that the heart had not recovered itself completely,